



FLY-NO-MORE[®]
16:16 MINERAL
WITH RADON-ORAL LARVICIDE

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

FOR CATTLE ONLY

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Mix or blend Nutrena Fly-No-More 16:16 Mineral into the daily ration and grain ration. Each pound of Nutrena Fly-No-More 16:16 Mineral contains 490 mg of Radon Oral Larvicide. Feed at a rate to provide an average daily intake of 1.0 mg of Radon Oral Larvicide per 100 lbs body weight. Use the following feeding table as a guide.

Body Weight of Animal Lbs	Ounces per head Nutrena Fly-No-More 16:16 Mineral to feed daily
1000	1.4
1100	1.5
1200	1.7
1300	1.8
1400	2.0

Do not feed other minerals after this program is started. Start feeding Nutrena Fly-No-More 16:16 Mineral daily in the spring when flies begin to appear and continue feeding through the summer until the fall until cold weather restricts fly activity. Nutrena Fly-No-More 16:16 Mineral may be fed up to slaughter and to marketing dairy cows. Nutrena Fly-No-More 16:16 Mineral prevents the development of horn flies, face flies, house flies and stable flies in the course of treated flies but it is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other methods to reduce the population of existing adult flies. In some cases, supplemental fly control measures may be needed in the around cattle lots and buildings to control house flies and stable flies which can breed and rear in cattle manure but to other decaying vegetable matter or silage in the premises. In order to achieve optimum fly control, Nutrena Fly-No-More 16:16 Mineral should be used in conjunction with other good management and sanitation practices.

Nutrena is the registered trademark of Nutrena Feeders Corp.
 Patent No. 3,132,842

242