NOTICE:

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee. expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or lestabilished sale practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combinations with other materials

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Evershield D Seed Protectant is for use by professional seed treaters only. Observe all cautions and limitations on labeling of products used in mixtures.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by st-, disposal, or by cleaning of equipment. Do not empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used or chemical reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies

CONTAINER DISPOSAL

Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL

Consult federal, state or local disposal authorities for -approved alternative procedures such as limited open burning.

STORAGE

Store in a cool dry place.

Contents: U.S. 5 Gallons

ers SEED PROTECTANT

Active Ingredients:

Chloroneb (1,4-dichloro-2,5-	
dimethoxybenzene) 29.5%	
Inert Ingredients	
Chloroneb (Demosan*) 2.8 lbs./gal.	

U.S. Patent No. 3,113,399 E.P.A. Reg. No. 1352-30 E P.A. Est. No. 36626-TN-1

المراف ومعود فالمراف والمراف والمعادية



ال.

Manufactured by **Chemical Products Division** Agricultural Products Dept. Cargill, Incorporated P.O. Box 9300 Minneapolis, Minnesota 55440

CAUTION

KEEP OUT OF REACH OF PRECAUTIONARY STAT

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Swallowed: Harmful if swallo cal attention

inhaled. May be harmful medical attention. Avoid inhali a respirator.

Skin: Avoid contact with sole Wash thoroughly with soap in hands and face thoroughly a water after use and before ear r Eyes: Avoid contact with $e \in$ contact with eyes, flush in n tunning water and get media unitation persists.

ENVIRONMENTAL HAZARD > This product is toxic to finn lakes, streams or ponds, Do 11 water by cleaning of equip: € of wastes





SEED PROTECTANT

tive Ingredients:

loroneb (1,4-dichloro-2,5-
nethoxybenzene) 29.5%
rt Ingredients
loroneb (Demosan*)2.8 lbs./gal.

3. Patent No. 3,113,399 7.A. Reg. No. 1352-30 7.A. Est. No. 36626-TN-1

inniel in livinegistiered trademark of Carg in-



Manufactured by Chemical Products Division Agricultural Products Dept. Cargill, Incorporated P.O. Box 9300 Minneapolis, Minnesota 55440

CAUTION

KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Swallowed, Harmful if swallowed, get medical attention.

Inhaled: May be harmful if inhaled, get medical attention. Avoid inhalation by using a respirator

Skin: Avoid contact with skin or clothing Wash thoroughly with soap and water. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Eyes. Avoid contact with eyes, In case of contact with eyes, flush immediately with running water and get medical attention it irritation persist.

ENVIRONMENTAL HAZARDS

ŪŪ

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS

- Keep product away from food or food products. Bags containing treated seed should be labeled "Treated seed do not use for feed, food or oil purposes."
- Apply with commercial seed treating equipment such as a slurry treater.
- 1. Use at Evershield concentrate ratios given in the table below. Add Evershield Seed Protectant concentrate to cool tap water and mix thoroughly for 15 minutes to prepare the Evershield dilution.
- Add Evershield Seed Protectant liquid dye** to each batch of Evershield dilution and mix for an additional 10 minutes.

•		Fluid Ounces of Concentrate per 100 pounds of seed				
	Seed	Evershield CM	Evershield T	Evershield D	Disease Controlled	
	Cotton	2.34	3.32 1.65	5.6-10.2* 5.6-10.2* 5.6-10.2*	1, 2, 3, 4	
	Beans and Soybeans	1.65 1.60	1 65 1.60	1.7-4.15* 1.7-4.15* 1.7-4.15*	1 2 4	
	Sugar beets	3 36 1 65	2,49 1.65	2.54-6 22* 2 54-6 22* 2 54-6 22*	1, 2, 4	
	Grasses (non-pastered)	4.57 2. 4 5	4 37 2 45	3 68-14 69* 3 68-14 69* 3 68-14 69* 3 68-14 69*	2, 4 2	

- Rhizoctonia solani, 2. Pythium Spp., 3. Scierotium rolfsi,
 Seed & seedling diseases.
- *Use high rate on pathologically intested seed or tor soils such as in California.

DO NOT GRAZE TREATED BEANS OR SOYBEANS WITHIN 45 DAYS AFTER PLANTING.

**21CFR Chapter 1 Section 3.13 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably polored to prevent than subsequent indivertent use as food for man or feed for animals.

31 800