

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Causes skin and eye irritation. Do not get in eyes, on skin, or on clothing. Avoid contamination of feed and foodstuffs. During exposure in seed treatment and loading operation, wear goggles, clean synthetic rubber gloves and protective clothing. Wash face, hands, and arms thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash work clothes with soap and hot water before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and wildlife. Do not apply directly to any body of water. Exposed treated seeds may be hazardous to birds or other wildlife. Dispose of all excess treated seed by burial away from bodies of water.

STORAGE AND DISPOSAL

Prohibitions:


Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal:

Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state and local procedures under the Resource Conservation and Recovery Act.

Container Disposal:

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other approved state or local procedures.



RTU
LINDANE

seed prote

Active Ingredient:
 Lindane (Gamma isomer of benzenehexachloride)
Inert Ingredients
Total
 This product contains 2.67 lbs. of Lindane per

KEEP OUT OF REACH OF CHILDREN
WARNING

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or poison control center immediately. Do not induce vomiting. Give small sips of water and induce vomiting by touching back of throat. Do not give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply respiration if necessary.

If on Skin: Wash the affected area with soap and warm water.

If in Eyes: Immediately flush eyes with plenty of running water.

See side panel for additional precautions.



Manufactured by
 Chemical Products Division
 Agricultural Products Dept.
 Cargill, Incorporated
 P.O. Box 9300
 Minneapolis, Minnesota 55440

RTU-LINDANE

Seed protectant

Ingredient:

γ (Gamma isomer of benzenehexachloride)	28.05%
Other ingredients	71.95%
	100.00%

Product contains 2.67 lbs. of Lindane per gallon.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

STATEMENT OF PRACTICAL TREATMENT

Call a physician or poison control center immediately. Drink 1 or 2 glasses of water to induce vomiting by touching back of throat. Do not induce vomiting or give anything to an unconscious person.

Move the victim to fresh air. Apply respiration if indicated.

Wash the affected area with soap and warm water.

Immediately flush eyes with plenty of running water. Call a physician.

See the back panel for additional precautionary statements.

Manufactured by
Cargill Chemical Products Division
Agricultural Products Dept.
Cargill, Incorporated
P.O. Box 9300
Minneapolis, Minnesota 55440

EPA Reg. No. 1352-40
EPA Est. No. 36626-TN-1

RTU is a Trademark of Cargill, Inc.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Using suitable liquid or slurry seed treating equipment such as Panogen or Mist-O-Matic, apply RTU-Lindane at rates

Seed should be sound, well cured and clean prior to treatment. Only the recommended rates of product per 100 pounds

RTU-Lindane will protect seeds and seedlings from white grub and seed corn maggots.

Seed

Corn
Peas
Sorghum
Wheat, rye, oats
Cucumbers, watermelons, canteloupes, squash
Sudan grass
Cabbage, cauliflower, broccoli, brussel sprouts
Carrots, onions
Barley
Sugar beets, beets
Beans
Sunflowers
Cotton:	
Acid delinted or reginned
Fuzzy - not delinted
Lentils
Safflower
Radishes

At the recommended rates, apply RTU-Lindane undiluted directly to the drum to the seed. If desired, add one part water to five parts Lindane and recalibrate equipment so that recommended rates still be applied. Slight agitation may be required to mix. Keep container tightly closed when not in use.

Treated seed should not be used for or mixed with food for human or animal feed, or processed for oil. Seed commercially treated with Lindane must be labeled in accordance to all applicable regulations of the Federal Seed Act.

Labels attached to treated seed must read:
TREATED SEED SHOULD NOT BE USED FOR FOOD OR ANIMAL FEED, OR PROCESSED FOR OIL. EXPOSED TREATED SEEDS MAY BE HAZARDOUS TO BIRDS AND OTHER WILDLIFE. DISPOSE OF ALL EXCESS TREATED SEED BY BURNING AWAY FROM BODIES OF WATER.

MIX CONTENTS THOROUGHLY BEFORE USE

NOTICE TO BUYER

The product conforms to its chemical description and is fit for use when used in accordance with label directions. Cargill will not be responsible for special, consequential or contingent damages arising from injury to property.