



# BHC EMULSIFIABLE CONCENTRATE

## INGREDIENTS

<b>ACTIVE INGREDIENTS:</b>	
Gamma isomer of benzene hexachloride .....	11%
Other isomers of benzene hexachloride and related compounds .....	13%
Aromatic petroleum derivative solvent .....	70%
<b>INERT INGREDIENTS</b> .....	6%
<b>Total</b> .....	<b>100%</b>

CONTAINS 1 POUND GAMMA ISOMER OF BHC PER GALLON

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

USDA Reg. No. 1386-32

Net Volume

FC-24B-6-69

1 U.S. Gallon Liquid

PROD. 205

MANUFACTURED BY

## CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. May produce irritation of the skin or eyes. Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

Do not apply under conditions which might result in residual odor or taste remaining on feed and foodstuffs or crops. Do not use, spill, pour, or store near heat or open flame, or any product which will absorb odors. Keep container closed when not in use. Do not use on animals of any kind.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of containers, rinse solutions, and unused chemical in a pit on non-crop land located well away from water supplies.

This product is poisonous to bees exposed to direct treatment or to residues in sprayed areas. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

## NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

ACCEPTED  
Jul 28, 1969  
UNDER THE FEDERAL INSECTICIDE  
ACT AND THE FEDERAL TOXIC  
SUBSTANCE ACT REGISTERED  
UNDER NO. 1386-32

UNICO BHC Emulsifi

DIRECTIONS FOR

1336 PINE TREES—To control Southern pine beetle, mix

and apply this mixture as a uniformly thorough spray pine trees. Begin applications early in the spring (March) and repeat at 2-3 week intervals through September. Apply also to injured and

To control turpentine beetles, mix 1 part of this product with 9 parts of water. Apply mixture as a uniformly thorough spray to the lower 1/3 of the trunk of living pine trees. Begin applications early in the spring (usually April and May) and repeat at 2-3 week intervals through September. Apply also to injured and felled pine trees, and to stumps.

DO NOT GRAZE LIVESTOCK IN TREATED AREAS. DO NOT APPLY TO PINE TREES IN AREAS TO LIVESTOCK. DO NOT APPLY TO PINE TREES IN AREAS WITH ROOT CROPS OR PEANUTS. BIRDS AND ANIMALS ARE KILLED.

**CAUTION**

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. May produce irritation of the skin or eyes. Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

Do not apply under conditions which might result in residual odor or taste remaining on feed and foodstuffs or crops. Do not use, spill, pour, or store near heat or open flame, or any product which will absorb odors. Keep container closed when not in use. Do not use on animals of any kind.

ACCEPTED  
Jul 28, 1969  
UNDER FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. ~~1386~~ 32

**UNICO BHC Emulsifiable Concentrate**

**DIRECTIONS FOR USE**

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of containers, rinse solutions, and unused chemical in a pit on non-crop land located well away from water supplies.

This product is poisonous to bees exposed to direct treatment or to residues in sprayed areas. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

**NOTICE**

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

**PINE TREES**—To control Southern pine beetle, mix 1 part of this product with 50 parts of water, and apply this mixture as a uniformly thorough spray to the bark area of infested and surrounding pine trees. Begin applications early in the spring (March or April) and repeat at approximately 30-day intervals through September. Apply also to injured and felled pine trees.

To control turpentine beetles, mix 1 part of this product with 14 parts of water, and apply this mixture as a uniformly thorough spray to the lower 12 feet of the trunks of pine trees. Begin applications early in the spring (usually April and May) and repeat within 3 to 4 weeks. Apply also to injured and felled pine trees, and to stumps.

**DO NOT GRAZE LIVESTOCK IN TREATED AREAS. DO NOT FEED ANY FOLIAGE FROM TREATED AREAS TO LIVESTOCK. DO NOT APPLY TO PINE TREES WHICH ARE TO BE FOLLOWED IN A ROTATION WITH ROOT CROPS OR PEANUTS. BIRDS AND OTHER WILDLIFE IN TREATED AREAS MAY BE KILLED.**