

ACCO ALGICIDE 15-01

Percent By Weight

ACTIVE INGREDIENTS:

Poly[oxyethylene(dimethyliminio)ethylene-
(dimethyliminio)-ethylene dichloride] 15.0 percent

INERT INGREDIENTS 85.0 percent

DIRECTIONS

ACCO ALGICIDE 15-01 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits. An initial slug addition of 4.0 to 10.0 fluid ounces of ACCO ALGICIDE 15-01 per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of ACCO ALGICIDE 15-01 based on the total weight of water in the system is recommended. Repeat initial dosage until control is evident. Subsequent slug additions of 1.0 to 10.0 fluid ounces of ACCO ALGICIDE 15-01 per 1000 gallons of water (8 to 80 parts per million of ACCO ALGICIDE 15-01) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

CAUTION! KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not get in eyes. Avoid contact with skin. In case of contact, flush with plenty of water for at least 15 minutes. If eye irritation persists, get medical attention. Avoid contamination of food.

Rinse empty container thoroughly with water and discard.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. Use only as directed on label.



Manufactured by
ACME CHEMICAL COMPANY
MILWAUKEE, WISCONSIN 53210

Form 4523-1

EPA Est. No. 1363 WI 1
EPA Reg. No. 1363-39

Printed in U.S.A.