

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

INDOORS APPLICATIONS: The use of this product in food processing establishments should be confined to time periods when the plants are not in operation. Food should be removed or covered during treatment. All food processing surfaces should be thoroughly cleaned before spraying.

VACUATE: In order to achieve the best results, vacuuming for maximum effectiveness, a combination of spot and surface treatments is recommended.

SURFACE SPRAYING: For control of roaches, ants, silverfish spiders, crickets, clover mites, cheese mites, granary weevils, flour weevils, confused flour beetles, rust red flour beetles, smutted grain beetles, spider beetles, cigarette beetles, drugstore beetles, mealworms, grain mites and cadellids, use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around corners of closets, etc., and at the base of machinery behind shelves and drawers. For silverfish spray the legs. For ants spray the nests and points of entry. If a face application only is to be used, spray floors and other surfaces applying at a rate of 1.5 gallons per 1000 cubic feet of space.

For control of carpet beetles, spray types of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray thoroughly into cracks. Floors are infested areas of shoring. Repeat treatment as needed.

To control bedbugs, spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all points. Treat boxes, and in all trunks and trunks. Repeat treatment as needed.

To control fleas and brown dog ticks, thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, in all dogs, windows and door frames, and on the areas of floor and floor covering.

IMPORTANT: Remove old bedding. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths, remove any infested articles from storage, brush thoroughly and air for several days in bright light, if possible. Apply spray thoroughly to empty chests, closets, bureaus and other storage places. Direct the spray into cracks, joints and crevices. After airing infested articles may be treated lightly, keeping in mind at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 1.5 gallons per 1000 cubic feet of space. For wasps and hornets control, spraying of these insects should be done late in the afternoon when swarms are at rest.

Mellocraft®

BEST DOCUMENT AVAILABLE COMMERCIAL SPRA-TOX AQUEOUS LIQUID INSECTICIDE

A Concentrated PYRENONE* Micro Emulsion
Clear — Stable — Leaves No Oily or Objectionable Residues
In view of National Energy Shortage, this product is extremely low
in petroleum distillate.
• Effective, Fast Acting Insecticide •
For Use In Homes, Institutions, Restaurants, Food Processing Plants,
Dairies, Livestock and Outdoors for Mosquito Control.
Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS

Pyrethrin
Pyrethrin, made in France
Petroleum distillate

INERT INGREDIENTS

Equivalent to 98% water and 2% pyrethrin concentrate
67% treated oil

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional cautions
DO NOT TRANSPORT OR STORE UNDER 32 F.

PYRENONE — Registered Trade Name of ICI, Ltd., England

FOGGING: For an enclosed or accessible space of the above mentioned height, use a rate of one gallon to 2000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult animal, which is sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. To control blood sucking lice, apply to the infested areas of the animal using a stiff bristle petting spray, the rest of the hair. Repeat every 2 to 3 weeks if required.

OUTDOOR APPLICATION:
ADULT MOSQUITO CONTROL: Apply as a fine mist. Do not spray the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Use marshy places, stagnant pools or ponds and storm drains. Apply at a rate of 1.5 gallons per acre (4.5 ounces per 1000 square feet) using truck mounted conventional misting equipment such as H. model 510-91 or 100 H. traveling at 5 mph to treat a 100 foot swath. Calibrate equipment to deliver 4.5 gallons per minute.

Precautionary Statements HAZARD TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply where food processing is in progress. After space spraying in food processing plants, wash all equipment, benches, storage etc., where exposed food will be handled with potable water. In the home, all food processing surfaces and utensils should be covered during treatment and thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARD

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local provisions. Refer to the Agency Registration and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling, reconditioning or disposal in a sanitary and safe manner, in accordance with state and local authorities.

NET CONTENTS: _____ Gallons

© 1989 T.M. Co.

Printed in U.S.A.

ANOTHER "QUALITY ASSURED PRODUCT" MANUFACTURED BY THE MELLOCRAFT COMPANY • TOLEDO, OHIO 43693