138 1 10-5-89 PM 17

(Front Panel)

LOL LOGOTM

PYRENONE® DAIRY CATTLE SPRAY

Protects Against Hornflies, Stableflies, Houseflies, Gnats, and Mosquitoes

FOR THE PROTECTION OF DAIRY ANIMALS FROM BITING FLIES AND AS AN AID IN THE CONTROL OF HOUSEFLIES IN BARNS, HOUSES AND MILK PARLORS

**Equivalent to 0.240% (butylcarbityl) (6-propylpiperonyl) ether and 0.060% related compounds.

^e Registered U.S. Patent Office (FMC)

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 1381-5 EPA Est. 407-MN-1 G-74-84-1

Distributed By Land O'Lakes, Inc. Fort Dodge, Iowa 50501

NET CONTENTS

TH - Trademark of Land O'Lakes, Inc.

(Side Panels)

PYRENONE DAIRY CATTLE SPRAY

PRECLUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid wetting the hides of animals. Protect food, milk, milking utensils from spray.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AN: DISPOSAL

STORACE: Store in a secure area, in original container only, away from fertilizers, food or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

CONTAINER DISPOTAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Pyrenone Dairy Cattle Spray is a special insecticide formulation compounded for the dairy industry. Pyrenone Dairy Cattle Spray is especially effective for rapid control of hornflies on dairy cattle and for the protection of cows from other biting flies, mosquitoes and gnats.

To protect dairy stock from hornflies, houseflies, stableflies, gnats, mosquitoes, spray once daily at milking time, after cattle enter the barn, using 1 to 2 ounces per head. The spray should be applied as a light rist primarily to envelop or hit the flies as the attempt to leave the cows. May be used twice daily at the rate of 1 ounce per animal. Do not use more than 2 ounces per day per animal. Avoid wetting the hides of the animals.

Calves may be sprayed very lightly once a day instead of attempting to build up a residual repellancy by spraying more heavily.

Use Pyrenone Dairy Cattle Spray as a space spray treatment in barns, houses and milk parlors for an aid in the control of flies. Direct mist toward the upper portions of the enclosure, filling the air with fine spray particles, using approximately 2 ounces of material for each 1,000 cubic feet of space. Whenever possible, keep dcors and windows closed for 10 minutes after application.

Food should be removed or covered during treatment. All food processing surfaces should be covered during treatments or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

∂∕_j

NOTICE

. •

.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

