

VAPONA[†] DAIRY CATTLE SPRAY

Directions for Use

DAIRY AND BEEF CATTLE

FACE FLY - STABLE FLY - HORN FLY - HOUSE FLY - MOSQUITOES: Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers and belly. For Stable Flies, make sure to spray leg areas and particularly knees. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.


OVERHEAD MIST DISPENSERS For control of flies, gnats and mosquitoes in milkhouses and barns, use 1 pint for each 5,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 5 hours of direct applications.

HOUSE FLIES in Dog Kennels, Hog Pens, Stables, Compost areas, Manure, Poultry Houses, Barns, Porches and Patios, Drive In's, and Restaurants. Apply to thoroughly cover walls, flooring, etc., in kennels, barns and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies. For porches, patios and Drive In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 square feet.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty, nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressly or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Distributed By
FARMERS REGIONAL COOPERATIVE



FELCO

VAPONA[†] Dairy Cattle Spray

Ready-to-Use

FOR FACE FLIES, STABLE FLY, HOUSE FLY, HORN FLY

INGREDIENT STATEMENT

ACTIVE INGREDIENTS:	By Weight
2,2-Dichlorovinyl Dimethyl Phosphate	0.93%*
Related Compounds	0.07%*
Petroleum Distillate	99.00%
	100.00%

* Equivalent to 1.00% w of DDVP Insecticide

† Registered Trade-Mark Shell Chemical Company

WARNING: KEEP OUT OF THE REACH OF CHILDREN

May Be Fatal If Swallowed, Inhaled Or Absorbed Through The Skin! Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, milk or milk handling equipment. Do not contaminate food or food utensils. If swallowed, induce vomiting.

NET CONTENTS 1 GALLON

VAPONA[†] DAIRY CATTLE SPRAY

Directions for Use

DAIRY AND BEEF CATTLE

FACE FLY - STABLE FLY - HORN FLY - MOSQUITOES: Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers and belly. For Stable Flies, make sure to spray leg areas at ankles. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

OVERHEAD MIST DISPENSERS For control of flies, gnats and mosquitoes in milkhouses and barns, use 1 pint for each 5,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 5 hours of direct applications.

HOUSE FLIES in Dog Kennels, Hog Pens, Stables, Compost areas, Manure, Poultry Houses, Barns, Porches and Patios, Drive In's, and Restaurants. Apply to thoroughly cover walls, flooring, etc., in kennels, barns and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies. For porches, patios and Drive In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 square feet.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty, nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressly or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Distributed By
FARMERS REGIONAL COOPERATIVE

Fort Dodge, Iowa

DAIRY
SPRAY

For Use

OF CATTLE

HORN FLY - HOUSE FLY - MOSQUITOES: Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers and belly. For Stable Flies, make sure to spray leg areas and particularly ankles. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.


OVERHEAD MIST DISPENSERS: For control of flies, gnats and mosquitoes in milkhouses and barns, use 1 pint for each 8,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 8 hours of direct applications.

HOUSE FLIES: in Dog Kennels, Hog Pens, Stables, Compost areas, Manure, Poultry Houses, Barns, Porches and Patios, Drive In's, and Restaurants. Apply to the rough cover walls, flooring, etc., in kennels, barns and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies. For porches, patios and Drive In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 square feet.

Product conforms to its label description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

By
FARMERS REGIONAL COOPERATIVE

NET CONTENTS 1 GALLON



FELCO

ACCEPTED
May 23, 1968
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE ACT AND FEDERAL ACT
FOR ECONOMIC PEST CONTROL REGISTERED
UNDER NO. 1381-9

VAPONA Dairy Cattle Spray

Ready-to-Use

FOR FACE FLIES, STABLE FLY, HOUSE FLY, HORN FLY

INGREDIENT STATEMENT

ACTIVE INGREDIENTS:	By Weight
2,2-Dichlorovinyl Dimethyl Phosphate	0.93%*
Related Compounds	0.07%*
Petroleum Distillate	99.00%
	100.00%

* Equivalent to 1.00% w of DDVP Insecticide

† Registered Trade-Mark Shell Chemical Company

WARNING: KEEP OUT OF THE REACH OF CHILDREN

May Be Fatal If Swallowed, Inhaled Or Absorbed Through The Skin! Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, milk or milk handling equipment. Do not contaminate food or food utensils. If swallowed, induce vomit-

VAPONA DAIRY CATTLE SPRAY

Directions for Use

cattle
DAIRY AND BEEF CATTLE

FACE FLY - STABLE FLY - HORN FLY - HOUSE FLY - MOSQUITOES: Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers and belly. For Stable Flies, make sure to spray leg areas and particularly ankles. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

OVERHEAD MIST DISPENSERS: For control of flies, gnats and mosquitoes in milkhouses and barns, use 1 pint for each 8,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 8 hours of direct applications.

HOUSE FLIES: in Dog Kennels, Hog Pens, Stables, Compost areas, Manure, Poultry Houses, Barns, Porches and Patios, Drive In's, and Restaurants. Apply to the rough cover walls, flooring, etc., in kennels, barns and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies. For porches, patios and Drive In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 square feet.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

© 1968 FARMERS REGIONAL COOPERATIVE

Distributed By
FARMERS REGIONAL COOPERATIVE