

1-17-90

LAND O'LAKES

VAPONA  
Dairy Cattle Spray  
Ready-To-Use

For the control of horn flies, face flies, stable flies,  
house flies, gnats and mosquitoes.

ACTIVE INGREDIENTS:

2,2-dichlorovinyl dimethyl phosphate.....	0.93%
Related Compounds* .....	0.07%

INERT INGREDIENTS:

Deodorized Kerosene .....	49.50%
Mineral Seal Oil .....	49.50%
Total .....	100.00%

\*Equivalent to 1% Vapona Insecticide  
Vapona is a registered trademark of Shell Chemical Co.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

NET CONTENTS           

Distributed by  
LAND O'LAKES, INC.  
Fort Dodge, Iowa 50501

Not Reviewed in accordance with  
Herbicide Registration Process for Phase 2  
Response, Page 2.12, Active to Inert Change in Status.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

May be fatal if swallowed, inhaled or absorbed through the skin! Wash thoroughly with soap and water after handling and before eating or smoking. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

**ENVIRONMENTAL HAZARDS**

Do not contaminate feed, water, milk or milk handling equipment. Do not contaminate food or food utensils.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a secure area, in original container only, away from fertilizers, food or feed.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**DAIRY AND BEEF CATTLE**

**FACE FLY-STABLE FLY-HORN FLY-HOUSE FLY-MOSQUITOES:** Use a hand mist sprayer and treat entire animal to wet hair with 1 ounce in morning and evening. For Horn Flies, especially treat the back line, withers and belly. For Stable Flies, make sure to spray leg areas and particularly ankles. Repeat daily as needed for control. Do not apply more than 2 ounces per animal daily. For walk-thru mist sprayers, set to treat animals at rate not to exceed 2 ounces daily. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organophosphates.

**OVERHEAD MIST DISPENSERS--**For control of flies, gnats, and mosquitoes in milkhouses and barns, use 1 pint for each 8,000 cubic feet. Do not contaminate feed or milk utensils. If a program of spraying animals directly is employed daily, do not fog within 8 hours of direct applications.

**HOUSE FLIES** in Dog Kennels, Hog Pens, Stables, Compost areas, Manure, Poultry Houses, Barns, Porches and Patios, Drive In's, and Restaurants. Apply to thoroughly cover walls, flooring, etc., in kennels, barns and stables. Use no more than 1 pint per 1,000 square feet. This is not a residual spray, therefore treat weekly or as needed for control of flies. For porches, patios and Drive In's, spray refuse area and trash. Treat walks, steps and adjacent walls with a light spray mist. Repeat as needed. Do not contaminate food or food utensils. Do not use more than 1 pint per 1,000 square feet.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller; and buyer assumes the risk of any such use.

EPA Reg. No. 1381-9

EPA Est. 407-IA-1