


DIRECTIONS FOR USE

GENERAL INFORMATION — Thorough, full coverage applications should be made. Repeat applications as necessary. Mix 57% Malathion with water at rate indicated. The per acre rate is mixed with sufficient water for thorough application.

APPLICATION CHART

For Use On	To Control	Per Acre in 1 Gal.	Comments
Apples.	European Red Mite	1-2 pts.	2 tsp.
Pears.	Cloyer Mite		
	Two Spotted Mite		
	Wooly Apple Aphid	1 pt.	1 tsp.
	Forbes Scale		
	Green Apple Aphid	1 1/2 pts.	1 1/4 tsp.
	Rosy Apple Aphid		
Tomatoes	Spider Mites	1 1/2 pts.	1 1/4 tsp.
	Aphids	1 pt.	1 tsp.
Beans	Mexican Bean Beetle, Spider Mites, Leafhopper	2 pts.	2 tsp.
Peas	Pea Aphid	2 pts.	2 tsp.
Potatoes	Aphids	2 pts.	2 tsp.
Cabbage.	Aphids, Imported	2 pts.	2 tsp.
Broccoli.	Cabbage Worm		
Kale.			
Mustard.			
Turnip.			
Alfalfa	Aphids	2 pts. per acre in at least 10 gal. water	Do not apply within 7 days of harvest, or during full bloom.
Flowers.	Oyster Shell Scale	1 pt.	1 tsp.
Ornamentals.	Euonymus Scale		
Shrubs	Aphids, Spider Mites, White Fly, Thrips, Mealy Bugs, Japanese Beetle Adult, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Bagworms.	1 1/2 pts.	1 1/4 tsp.
	Juniper Scale	2 pts.	2 tsp.
	Soft Brown Scale	2 1/2 pts.	2 1/4 tsp.
	Pine Needle Scale	4 pts.	4 tsp.

CAUTION: Do not use on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula and Canaerti Juniper, because of possible injury to these ornamentals.



FELCO

NET CONTENTS
1 PINT

57% MALATHION

EMULSIFIABLE CONCENTRATE

Contains 5 lbs. of Malathion per gal.

ACTIVE INGREDIENTS:

Malathion*	57.0%
Xylene Range Aromatic Hydrocarbon	34.0%
INERT INGREDIENTS:	9.0%
	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See side panel for other cautions.
Keep container closed and away from heat, sparks, and open flame. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

DIRECTIONS FOR USE

LIQUID FLY BAIT: To control house and stable flies in Barns, Feed Lots, Poultry Houses, City Buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add 2 gallons of water. Paint or spray desired surfaces such as window frames, stanchions, support beams and doorways.

SWINE: For lice, mix 1 gallon per 100 gallons of water. Apply a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat applications only if needed.

BEEF CATTLE & NON-LACTATING DAIRY CATTLE: For lice, add 1 gallon 57% Malathion to 100 gallons of water. For ticks, mix 1 to 2 gallons per 100 gallons of water. For horn flies, mix 1 to 1 1/2 gallons per 100 gallons of water. Apply complete coverage spray. Repeat at 2 week intervals if needed. Do not spray cattle during lactation or within two weeks of freshening. Do not apply to calves under one month of age.

NOTE: When applying Malathion sprays do not contaminate food containers and watering troughs. Do not contaminate milk, milking equipment, feed or drinking water. Toxic to fish and aquatic life. Do not contaminate any body of water.

BACK RUBBING DEVICES: As an aid in the control of lice or horn flies on Beef Cattle, apply a mixture of 2% Malathion (1 gallon 57% Malathion to 28 1/2 gallons fuel oil or kerosene, or 1 quart 57% Malathion to 7 1/4 gallons oil) to the back rubbing device. These devices should be made continuously accessible, one to each 35 to 45 head of cattle. Retreat the oiler every 2 to 3 weeks. Do not apply to dairy animals.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION: Keep out of the reach of children. Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTE TO PHYSICIAN: Malathion, upon repeated, prolonged, or careless use, may cause cholinesterase inhibition. Atropine is antidotal.

USDA Reg. No. 1381-11

J-6821A-J

Distributed by
LAND O'LAKES, INC.,
FORT DODGE, IOWA

ACCEPTED WITH COMMENTS