

TOXAPHENE

TOXAPHENE

LAND O LAXES

DIRECTIONS FOR USE

READ APPROPRIATE CROP LIMITATIONS BEFORE USE

Apply mixed with water to give thorough coverage in accordance with proper sprayer calibration. When applied by airplane, use at least 3 to 10 gallons of water per acre, consistent with crop growth and thorough coverage.

GRASSHOPPERS (In Corn, Oats, Soybeans, Wheat): For young hoppers, mix 1 qt. 1 1/2 lbs. actual toxaphene in sufficient water to cover one acre. For mature hoppers, increase dosage to 1 quart per 10 to 20 lbs. actual toxaphene in sufficient water per acre. Do not apply to soybeans within 21 days of harvest. Do not treat forage or silage made from any of the above crops to dairy animals or animals being finished for slaughter. Do not use on grain and stock to live stock or poultry.

FIELD CORN (Grown For Grain Only): Cutworms, mix 1 quart in 20 to 25 lbs. actual toxaphene in sufficient water to cover one acre. Apply at a 12 inch band over 16 row corn less than 12 inches tall, or at the base of the plant in corn over 12 inches tall. To improve control of cutworms in tall corn, cultivate and hoe to remove weeds. Do not feed treated corn to dairy animals or animals being finished for slaughter.

WARNING

Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.


Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.

Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.

Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.

Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.

Do not use this product on any crop not listed on the label. Do not use on any crop not listed on the label. Do not use on any crop not listed on the label.



TOXAPHENE

TOXAPHENE

DIRECTIONS FOR USE

READ APPROPRIATE CROP LIMITATIONS BEFORE USE

Apply diluted with water to give thorough coverage in accordance with proper sprayer calibration. When applied by airplane use at least 3 to 10 gallons of water per acre, consistent with good crop growth and thorough coverage.

GRASSHOPPERS (In Corn, Oats, Soybeans, Wheat): For young hoppers mix 1 qt. (1½ lbs. actual toxaphene) in sufficient water to cover one acre. For mature hoppers, increase dosage to 1 quart plus 10 oz. (2 lbs. actual toxaphene) in sufficient water per acre. Do not apply to soybeans within 21 days of harvest. Do not feed treated forage, or ensilage made from any of the above plants to dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

FIELD CORN (Grown For Grain Only): Cutworms: mix 1 quart plus 10 oz. (2 lbs. actual toxaphene) in sufficient water to cover one acre. Apply in a 12 inch band over the row (corn less than 12 inches high) or at the base of the plants (corn over 12 inches high). To improve control of older cutworms, in tall corn, cultivate at once to bury the insecticide. Do not feed treated forage to dairy animals or animals being finished for slaughter.

WARNING

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after using. Do not store near feed or food products.

Not for use on household pets, dairy animals, animals being finished for slaughter or humans. Do not use in dairy barns or milkhouses.

This product is toxic to fish, birds, or other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse container. Destroy when empty.

A COMPANY

3-5-73

1381-52

FOR THE FEDERAL BUREAU OF INVESTIGATION
WASHINGTON, D. C.
RECEIVED
MAY 10 1973
ATTACHED COMMENTS.

LANDOLAXES



FELCO

