

Center Panel

NET CONTENTS 1 GALLON

ACCEPTED
AUG 16 1967
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1211-42 SUBJECT TO ATTACHED COMMENTS.

FELCO/STATEX

STOCK-TOX PLUS

(Livestock Spray Concentrate)

Use as a Dip or Spray to Control Horn Flies, Lice, Mites, and Ticks

Active Ingredients:

Toxaphene (Technical Chlorinated Camphene containing 67 to 69% Chlorine	45%
✓ Gamma Isomer of Benzene Hexachloride from Lindane	1%
Petroleum Hydrocarbon	44%
Inert Ingredients:	10%
Total	100%

WARNING

KEEP OUT OF THE REACH OF CHILDREN


May be fatal if swallowed! Do not use on humans or household pets. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed and food products. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. When this container is empty, destroy beyond all use. Do not reuse for any purpose.

ANTIDOTE: External: Wash with soap and water.
Internal: Give sodium chloride 15 g. (One tablespoon of table salt) in glass of warm water and repeat until vomit is clear. CALL A PHYSICIAN.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

LEFT PANEL

1381-68



DIRECTIONS FOR USE

Dilute with water and use either as a spray or a dip.

Stock Tox Plus	Amount of water
1 gallon	100 gallons
1 quart	25 gallons
1 pint	12 gallons
1 cup	6 gallons

BACKRUBBER USE

1 gallon	10 gal. oil
1 pint	1 1/4 gal. oil

NOTE: Do not apply to dairy animals, or calves under 6 months of age. Do not spray meat animals within 30 days of slaughter.

For controlling horn flies, on cattle; lice, mites (mange/scab mites) on cattle, hogs, sheep and goats; and ticks on cattle, sheep and goats.

SPRAYERS - High-pressure sprayers get better results when large numbers of animals are to be treated. Low-pressure sprayers and hand sprayers do the job for small groups of animals if the spray solution is mixed thoroughly before application.

IMPORTANT: Animals should be sprayed on a warm sunny day during the winter months. This makes it easier to complete the operation in time for the animals to dry before nightfall.

DIPPING VATS - Dipping vats are of great importance when large numbers of livestock are to be treated at one location. Dipping is the preferred method for mange or scabies where complete coverage of the animal is essential. Do not dip meat animals within 60 days of slaughter.

BACKRUBBER USE: For horn fly and lice control, dilute 1 gallon with 10 gallons fuel oil. Apply the diluted material to saturate the backrubber mechanism. Recharge the backrubber every 2 weeks or as necessary for control. Locate backrubber to give cattle or hogs free access to it. Best locations are usually watering places, salt licks, or feed racks.

Note: Do not allow calves under 6 months of age, or dairy animals to treat themselves. Do not permit meat animals access to this treatment within 30 days of slaughter.

Distributed by
FARMERS REGIONAL COOPERATIVE
Fort Dodge, Iowa

Right Panel

DIRECTIONS FOR USE:

FACE FLIES: For use against face flies one end of the backrubber must be low enough (not more than 15 to 20 inches above the ground to allow cattle to rub their faces and heads. The product is only effective for those cattle that rub the face and head. Dilute as for backrubber use.

HOGS: lice, sarcoptic mites -- A single application will usually give control. Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Pigs should not be sprayed when less than weaning age. Do not spray meat animals within 30 days of slaughter.

HORN FLIES, TICKS (such as cattle fever ticks, ear tick, Gulf coast tick, Lone Star tick and winter tick) and LICE: cattle, sheep and goats. Dip or spray. When dipping make sure the animal is held in the vat at least two minutes and submerge each animal twice during dipping. For sheep tick apply during warm weather soon after shearing or before the wool is 2 inches long. When spraying, use about 2 quarts per beef animal and proportionately less for smaller animals. If reinfestations occur repeat applications in 15 to 21 days or as needed for control. If calves are infested and must be treated, avoid dipping, apply spray sparingly. Do not add infested animals to the treated herd. Do not apply to dairy animals, or calves under 6 months of age. Do not apply to dairy goats. Do not spray meat animals within 30 days of slaughter.

SARCOPTIC MANGE: (cattle, sheep and goats) Dip or spray cattle, dip sheep and goats, for thorough animal coverage apply to herd when mange is found. If infestation is severe plan on a second treatment 15 to 21 days later. Do not apply to dairy animals or calves under 6 months of age. Do not spray meat animals within 30 days of slaughter.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Distributed by

FARMERS REGIONAL COOPERATIVE
Fort Dodge, Iowa

USDA Reg. No. 1381-68

