

Right Panel (Continued)

SMALL GRAINS - (Wheat, Barley, Rye, Oats): For use in winter wheat, apply, treat only in the fall. Use 3/8 to 1 1/4 pints in sufficient water to cover the seed. Apply before the tiller to the jointing stage. In spring, wheat, Oats, and barley, use 3/8 to 4/5 pint per acre. Use higher dose for hard to kill weeds. Treat when crop is fully tillered (4 to 6 inch high) up until early boot stage. Do not spray during seedling stage or boot to dough stage. Grain interplanted with legumes should not be sprayed unless injury to legumes can be tolerated.

PASTURES: For broadleaved weeds (see list) use 2 1/2 pints per acre in sufficient water to thoroughly cover. Apply early in summer when weeds are actively growing and near bud stage. Do not apply to recently seeded pastures until grass is well established. Do not graze pastures to dairy animals within seven days of application. Do not apply when grass is in boot to milk stage, or when legumes are in bloom. Most legumes, including white clover, are not killed, and, under some conditions, killed. Avoid drift of spray onto vegetables, flowers, ornamental trees and shrubs, and other desirable plants.

Notice

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of the product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Distributed by

FARMERS REGIONAL COOPERATIVE
Fort Dodge, Iowa

USDA Reg. No. 1381-66

0-24 10 1167
1381-66