

DIRECTIONS FOR USE

MEAT AND FOOD PROCESSING PLANTS: Ants, Centipedes, Clover Mites, Cockroaches, Crickets, Earwigs, Silverfish, Spiders and Black Carpet Beetles, Trogoderma species; Clean premises thoroughly before applying and maintain good sanitation at all times. Use spray equipment and nozzles that will produce a coarse spray. Apply as a residual spray to all sections of the plant and warehouses where insects are present or may hide, into cracks and crevices (such as openings in expansion joints, between different elements of construction and between machinery and floors). Apply as a water solution at the rate of 1 pint in 2 1/2 gallons of water (3 tablespoons in one quart). To make an oil solution, dilute 1 part spray concentrate in 19 parts of a mixture of 4 parts kerosene type solvent and 1 part aromatic hydrocarbon type solvent. Avoid contamination of food products, equipment, utensils, work surfaces, containers, water, and liners. Do not use in edible products areas of meat and food processing plants. Repeat application as necessary.

BOX CARS: Residual Spray; For control of Confused Flour Beetle, Flat Grain Beetle, Granary Weevil, Indian Meal Moth, Lesser Grain Borer, Mites, Red Flour Beetle, Rice Weevil, Rusty Grain Beetle and Saw-Toothed Grain Beetle. Clean car thoroughly, remove and burn debris. Spray walls and floor to runoff with a solution of 1 gallon in 25 gallons of water or 1 gallon in 19 gallons of deodorized kerosene (a mixture of 4 parts kerosene type solvent with 1 part aromatic hydrocarbon solvent). Let the empty sprayed car stand with doors open until car is completely dry. Line walls and floor with paper before loading.

FLY CONTROL: For use in and around which buildings house domestic animals, around yards, homes and meat and food processing plants. Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

ADULT FLIES: Mix 5 tablespoons per gallon water or 1 cup in 2 1/2 gallons or 1 qt. in 12 gallons. Apply as a spray at the rate of 1 gallon per 1,000 sq. ft. on painted surfaces and 2 gallons per 1,000 sq. ft. on unpainted surfaces where flies gather, such as walls, ceilings, windows on dairy barns, around garbage cans and waste storage areas.

ADULT FLIES AND FLY MAGGOTS: As a Bait Spray, mix 5 tablespoons plus 7 tablespoons unsulfurized molasses or sugar or corn syrup per gallon water or 1 cup plus 1 cup in 2 1/2 gallons or 1 quart plus 2 1/2 pounds per 12 gallons water. Apply as a bait spray over the breeding sites and resting places such as waste and manure piles, bedding in loafing sheds or holding pens within 18 inches of walls and around upright braces. Do not apply to freshly white washed surfaces. Wait 14 days before reapplying. Avoid contamination of milk, milking equipment and water fountains and feed troughs. Remove lactating dairy animals before treating building. Also remove animals under one month of age.

JUN 12, 1975
1321-81

LAND O LAKES®

RESIDUAL SPRAY CONCENTRATE

For Use in and Around Meat and Food Processing Plants and in Box Cars and Stored Product Holding Areas.

ACTIVE INGREDIENTS:	
Malathion*	57.0%
Xylene Range Petroleum Hydrocarbons	34.0%
INERT INGREDIENTS	9.0%
Total	100.0%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.
*Cythion registered trademark American Cyanamid

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

WARNING—FLAMMABLE! KEEP AWAY FROM HEAT AND OPEN FLAME

Distributed By
EPA Reg. No. 1381-81 **Land O'Lakes, Inc.** N-3439B-I
EPA Est. 407-1A-1 **FORT DODGE, IOWA 50501**

NET CONTENTS _____

DIRECTIONS FOR USE

GRAINS (Barley, Corn, Grain Sorghums, Oats, Rye, Wheat) and Field and Garden Seeds: For Confused Flour Beetle, Cereal Leaf Beetle, Flat Grain Beetle, Granary Weevil, Indian Grain Moth, Lesser Grain Borer, Maize Weevil, Red Flour Beetle, Rice Weevil, Rusty Grain Beetle, Saw-toothed Grain Beetle.

RESIDUAL SPRAY BEFORE STORING GRAINS: For a residual wall, floor and machinery spray in grain elevators, in treating truck beds, box cars and ships holds, before loading grain, clean area to be treated thoroughly. Remove and burn all sweepings and debris. Mix 1 gallon in 25 gallons of water and make a thorough application.

GRAINS GOING INTO STORAGE: Apply to grain as it is being loaded or turned into final storage at the rate of 1 pint in 2 to 5 gallons of water per 1,000 bushel.

AS A SURFACE TREATMENT AFTER GRAINS ARE STORED: Apply evenly over the grain surface after loading into storage at the rate of 1/2 pint in 1 to 2 gallons of water per 1,000 sq. ft. of grain surface area. Repeat as necessary.

CAUTION

Harmful if swallowed. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Do not contaminate food or feed products.

NOTE TO PHYSICIAN: Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse container. Dispose of by crushing and burying in non-cropland away from water supplies.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressed or implied, extends to the use of this product contrary to label directions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.