

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When used on cattle, FLYSTOP™ will carry the flies through the life cycle system in the animal's manure where the fly larvae do not shortly after fly egg hatch. By preventing larval development, the flylarvicide helps to control house flies and face flies on pastured and stable flies in and around cattle lots.

FLYSTOP MINERAL HAYLAGE FORMULA at a rate of one half ounce per 250 pounds of body weight per head per day, mixing it with ground grain or complete daily ration.

GENEX FLYSTOP MINERAL HAYLAGE FORMULA early in the season before flies begin to appear and continues feeding throughout the summer into the fall until cold weather restricts fly activity.

FLYSTOP MINERAL HAYLAGE FORMULA may be fed up to six months to lactating dairy cows.

GENEX FLYSTOP MINERAL HAYLAGE FORMULA prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population of existing adult flies.

In some cases, supplementary control measures may be needed to keep around cattle lots and buildings to control adult house flies and stable flies which cannot be controlled by cattle manure treatment. Other decay and cleanup matter or sludge on the premises.

In order to achieve optimum fly control, FLYSTOP MINERAL HAYLAGE FORMULA should be used in conjunction with other good management and sanitation practices.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSITION: This product is not to be disposed of in any way that would result in the release of the active ingredient into the environment.



FLYSTOP MINERAL HAYLAGE FORMULA

(Contains RABON® Oral Laxative)

To prevent the development of Horn Flies, Face Flies, House Flies, and Stable Flies in the manure of treated cattle.

ACTIVE INGREDIENT	BY WEIGHT
2-Chloro-1,2,4,5-tetrachloro-1,1-dimethyl phosphate	1.19%
INERT INGREDIENTS	98.81%
TOTAL	100.00%

* Each pound contains 5.4g RABON®
* Refers only to ingredients which are not larvicidal.

GUARANTEED ANALYSIS

Phosphorus (P)	Minimum	20.0%
Calcium (Ca)	Maximum	1.0%
Equivalent Crude Protein from Non-Protein Nitrogen	Maximum	57.0%
Iodine (I)	Minimum	60%
Vitamin A	Not less than 250,000 USP Units per lb.	
Vitamin D	Not less than 125,000 USP Units per lb.	

FEED INGREDIENTS

Monoammonium Phosphate, Potassium Sulfate, Magnesium Sulfate, Sodium Bentonite, Magnesium Oxide, Dried Molasses, Potassium Chloride, Zinc Oxide, Manganese Oxide, Iron Carbonate, Iron Sulfate, Copper Oxide, Cobalt Carbonate, Ethylene Diamine, Dithiothiosulfate, Vitamin A Acetate, 25% Activated Animal Steroids Source of Vitamin D, Natural and Artificial Flavors, Mineral Oil.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists see a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of waste.

MANUFACTURED BY
FARMERS UNION CENTRAL EXCHANGE
GENERAL OFFICES, ST. PAUL, MINNESOTA

NET WEIGHT 5 LBS.