

1381-153

02/18/2005

1/3



ACCEPTED
 FEB 18 2005
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1381-153

**6% MALATHION
GRAIN DUST**

ACTIVE INGREDIENT:

Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate6%

OTHER INGREDIENTS:94%

TOTAL.....100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**FIRST AID
ORGANOPHOSPHATE**

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably to mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For further emergency medical information call toll-free 1-877-424-7452.

NOTE TO PHYSICIAN: Malathion is an acetyl cholinesterase inhibitor. Atropine is antidotal.

See below for additional Precautionary Statements.

EPA Reg. No. 1381-153

EPA Est. No. 1015-MO-1

Distributed By
Agriliance, LLC
P.O. Box 64089, St. Paul, MN 55164-0089

0/H27/4

NET WEIGHT 50 LBS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May be harmful if swallowed. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Use only with adequate ventilation. After using this product, do not enter treated grain storage areas until area has been ventilated and dust has settled. Avoid contact with treated surfaces.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its label.

STORAGE AND DISPOSAL

STORAGE: Store in a secure area, in original container only, away from fertilizers, food, or feed.
PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste facility. Do not contaminate water, food, or feed by storage or disposal.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned, stayed out of smoke.

GENERAL INFORMATION

FOR USE ON STORED GRAINS: Wheat, corn, oats, rye, barley. Protects against damage from Confused Flour Beetle, Rice Weevil, Saw-toothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, and Indian Meal Moth.

IMPORTANT: USE OF THIS PRODUCT, AS DIRECTED, WILL BE WITHIN THE TOLERANCE OF 8 PARTS PER MILLION FOR MALATHION. TREATED GRAIN IS NOT SUBJECT TO DOWNGRADING AND CAN BE MILLED INTO FLOUR OR FEED USING NORMAL PRODUCTION METHODS.

STORED GRAIN: Apply this product to the grain at the rate of ten lbs. per 1,000 bushels as the grain is being loaded or turned into final storage. To protect stored grain from attack by Indian Meal Moth, apply this product on the surface of grain at the rate of 5 lbs. per 1,000 square feet of surface area. Apply the dust evenly over the surface immediately after the grain is loaded into storage. Repeat if necessary.

METHODS OF APPLICATION: One convenient way to apply this product to shelled corn, wheat, corn, oats, rye, and barley is to treat it when it is in the field wagon or truck prior to binning. Determine the amount in the wagon or truck by multiplying the length by width by depth (all in feet) and multiply this total by 0.8 to get the number of bushels. Measure out the correct amount of this product and spread evenly over the surface of the grain in the wagon or truck bed. Cut the dust into the grain with a shovel. The dust will become mixed with the grain as it is hauled from the field to the storage bin, and as it is dumped from the bed into the auger hopper. Further mixing will occur when the grain is augered or blown into the storage bin. Mixing is completed as the treated grain rolls down the cone shaped pile in the bin. Another method of applying this product to grain is to evenly meter the correct amount of dust onto the grain as it is being transferred into storage, using a suitable metering device. Mixing is completed as the grain is transferred into storage and rolls down the cone shaped pile in the bin.

NOTICE OF WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.