



DELTA-COAT AD

Seed Treatment Fungicide

ACTIVE INGREDIENTS:

Chloroneb (1,4-dichloro-2,5-dimethoxybenzene)30.0%
 Metalaxyl (N-(2,6-dimethylphenyl)-N-(methoxyacetyl) alanine methyl ester.....3.5%

INERT INGREDIENTS*:66.5%
TOTAL.....100.0%

Contains 2.9 lbs. a.i. chloroneb and 0.34 lbs. metalaxyl per gal.


* Contains petroleum distillates, xylene or xylene range aromatic solvent.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

- If in eyes**
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If swallowed**
 - Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If inhaled**
 - Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably to mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.
- If on skin or clothing**
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.



Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For further medical emergency information, call toll-free: 1-877-424-7452.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 1381-166

EPA Est. No.

Distributed By:
Agrilliance, LLC
P.O. Box 64089,
St. Paul, MN 55164-0089

Net Contents
60 fl. oz.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, or inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapors of spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants; chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton; and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS: When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contaminations.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Failure to follow directions on this label may result in crop injury or poor disease control.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted entry intervals. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If the product is used for seed treatment, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton; shoes plus socks.

CONDITIONS OF SALE: Delta-Coat AD is a systemic fungicide having a specific mode of action. Delta-Coat AD could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, Agrilience, LLC cannot assume liability for crop damage resulting from resistant strains of fungi. Consult with your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible Delta-Coat AD resistant strains of fungi which may occur.

→ Use the high rate when conditions for disease expression are favorable.

BEANS (String beans, black-eyed peas, cowpeas, field beans, green beans, kidney beans, lima beans, navy beans, pinto beans, pole beans, snap beans, wax beans, and lupine): For Pythium and Rhizoctonia damping-off, apply Delta-Coat AD as a seed treatment at the rate of 5.5 to 7.0 fl. ounces per 100 pounds of seed.

COTTON: For Pythium and Rhizoctonia seed rot and damping-off of acid or machine delinted cotton seed, apply Delta-Coat AD as a seed treatment at the rate of 11.75 fl. ounces per 100 pounds of seed.

SOYBEANS: For early season Phytophthora control and Pythium and Rhizoctonia damping-off, apply Delta Coat AD as a seed treatment at the rate of 5.5 to 7.0 fl. ounces per 100 pounds of seed.

Do not use excess seed for food, feed or oil purposes.

Do not graze beans or soybeans within 45 days of planting. California registration prohibits grazing plants grown from treated seed.

Planter Box Application:

1. Pour one bushel of seed into each planter box.
2. Pour enough Delta-Coat AD on top of the seed to treat one bushel.
3. Mix thoroughly with an axe handle or paddle (an axe handle or paddle used for seed treatment purposes only) until seed is uniformly coated.

Drill Box Application:

1. Fill drill one-half full, counting the number of bags (bushels) put into the drill.
2. Pour 4 fluid ounces of Delta-Coat AD on top of each bushel of seed just put into the drill.
3. Mix thoroughly with an axe handle or paddle (an axe handle or paddle used for seed treatment purposes only) until seeds are uniformly covered.
4. Fill the remainder of the drill and repeat the process in items 2 and 3.

Vacuum Planters:

Pour 50 pounds of seed into the planter box. Pour enough Delta-Coat AD on top of the seed to treat 50 pounds of seed. Mix thoroughly with an axe handle or paddle until seed is uniformly covered. Then add any plantability additive at the recommended rate to improve seed flowability (talc, etc.). Mix thoroughly once more with an axe handle or paddle to get uniform coverage. This process will provide best seed flow through the planter while providing all the benefits of a seed treatment product.

Commercial Seed Treaters:

Delta-Coat AD may be applied with other registered seed treatment insecticides and fungicides through mist-type or other commercial seed treatment equipment. When using with other pesticides, observe and follow all precautions, restrictions and directions on the other pesticide product labels.

NOTE: Federal law requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with Delta-Coat AD fungicide. Do not use excess treated seed for food, feed or oil purposes."

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring. Do not store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Notice of Warranty

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

Treatment of highly mechanically damaged seed or seed of known low vigor and poor quality except for the purposes of curative control of existing disease or insect pests may result in reduced germination and/or reduced vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Agrilience, LLC no claims are made to guarantee germination of carry-over seed.