

1381-181

9/22/2003

Tacked
1/6



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
1381-181

Date of Issuance:
SEP 22 2003

Term of Issuance:
Conditional

NOTICE OF PESTICIDE:

Registration
Reregistration

(under FIFRA, as amended)

Name of Pesticide Product:
PRO SOURCE #1
FERTILIZER with 1.00%
RONSTAR® Herbicide

Name and Address of Registrant (include ZIP Code):

Agriliance LLC
P.O. Box 64089
St Paul, MN 55164-0089

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Add the phrase "EPA Registration No. 1381-181" to the label before you release the product for shipment.

COMMENTS CONTINUED ON PAGE 2 OF THIS NOTICE OF REGISTRATION

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Enclosure

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Signature of Approving Official:

Joanne I. Miller

Date:

SEP 22 2003

2/6

2. Submit the following data required for the registration of this pesticide product within 1 year from the date of this Notice of Registration:

<u>EPA Guideline Data Number</u>	<u>Guideline Descriptor</u>
830.6317	Storage Stability Study
830.6320	Corrosion Characteristics Study

3. Resubmit the Confidential Statement of Formula (CSF) on EPA Form 8570-4 to reflect the present registrant of Ronstar Intermediate 50% the source of oxadiazon in this pesticide product.
4. Delete from top of page 2 of labeling the name: "Pro Source #1 Magic Carpet Fertilizer with 1.00% ronstar®. And, give the name of the present owner of the trade mark name "Ronstar" and move the CAS No. from active ingredient line to a line below the "Total" line of the ingredient statement.
5. Submit one (1) copy of the final printed labeling before you release this product for shipment.
6. Submit and/or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA, section 4.

A Stamped copy of the label is enclosed for your records.

PRO SOURCE #1 FERTILIZER with 1.00% RONSTAR® HERBICIDE

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF MANAGEMENT AND LANDSCAPE PERSONNEL ONLY. NOT FOR USE ON HOME LAWNS, FOR USE ON GOLF COURSES, NURSERIES, PARKS, SOD FARMS, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS
 A Pre-emergent Weed Control Product plus a Quality Fertilizer for use on varieties of Perennial Bluegrass, Bentgrass, Bermuda grass, Buffalograss, Perennial Ryegrass, St. Augustine grass, Seashore Paspalum, Tall Fescue and Zoysiagrass. Also For Use Around Ornamental Shrubs, Vines, Trees & Ground Covers.

ACTIVE INGREDIENT: Oxadiazon CAS [19666-30-9] - 1.0%
 OTHER INGREDIENTS: - - - - - 99.0%
 Total 100.0%

**KEEP OUT OF REACH OF CHILDREN
 WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF IN EYES** - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING**: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED**: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; chemical resistant gloves and footwear, plus socks; protective eyewear; chemical-resistant headgear for overhead applications for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigations system. Apply only as specified on this label.


AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical resistant gloves, and footwear, plus socks, and protective eyewear. 09-18-02

Agrilance LLC

P.O. BOX 64089 - St. Paul, MN 55164-0089



Controls Crabgrass, Goosegrass, Poa Annua, Annual Sedge and other weeds as listed, on Golf Courses, Commercial Lawns, and other Athletic, Recreational, Ornamental Turf.

WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS

This product may be used at full rate or as a split application in established perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass area. **Not for use on home lawns.** The turf should be free of leaves and thatch and in condition to allow granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply using a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" may temporarily occur from the "doubled rate". See section on this label entitled; **PRECAUTIONS IN USING ON TURFGRASS** for the maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. a.i. per acre which is equiv. to 6.9 lbs. of this product per 1,000 sq. ft. After applications if rain is not expected within 24 hours sprinkle irrigate the treated turf with up to a 1/4 inch of water to get maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination. It will not control growing weeds. See section on this label entitled; **FOR SPRING APPLICATION ON GOLF COURSES** for instructions to apply the 3 lb. or 4 lb. a.i. per acre rate as a Split Application.

Pre-emergence Weed Control

Weeds Controlled	lbs. a.i. / Acre	lbs of Product per 1,000 sq.ft.	Directions - Refer to Map of U.S.
Crabgrass (<i>Digitaria</i> , spp.)	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	Apply in late winter or early spring prior to weed seed germination. See map. In areas 4 & 5 applications made after April or May, may not be fully effective.
Goosegrass (<i>Elysiene indica</i>)	200 to 400 lbs. product / acre		
Field Sandbur (<i>Cynchus incertus</i>)	2 to 4 lbs. a.i./A	4.6 to 9.2 lbs.	Apply in late summer to early fall prior to weed seed germination.
Annual Sedge (<i>Cyperus compressus</i>)	200 to 400 lbs./a		
Annual Bluegrass (<i>Poa annua</i> spp.) (See Note Below)	4 lbs. a.i./A	9.2 lbs.	Apply in early spring prior to weed seed germination.
Virginia Buttonweed (<i>Diodia virginiana</i>)	400 lbs. product /acre		
Pigweed (<i>Amaranthus</i> spp.)			
Florida Pusley (<i>Richardia scabra</i>)			
Oxalis (<i>Oxalis</i> spp.)			
Stinging Nettle (<i>Urtica dioica</i>)			
Carpentergrass (<i>Axonopus officinis</i>)			

Fall Application for Spring Weed Control in Cool Season Turf

Crabgrass, Goosegrass and other annual weeds as listed.	3 to 4 lbs. a.i./A 300 to 400 lbs. product / acre	6.9 to 9.2 lbs. / 1,000 sq.ft.	For use in Bluegrass, Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup.
---	--	--------------------------------	---

Annual Bluegrass Control in Bermudagrass Fairways Overseeded with Perennial Ryegrass Only.


Annual Bluegrass (<i>Poa annua</i> spp.)	2 lbs. a.i./A 200 lbs./acre	4.6 lbs. / 1,000 sq.ft.	Apply in late summer to early fall prior to <i>Poa annua</i> spp., seed germination. Any <i>Poa annua</i> spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where <i>Poa annua</i> is present in the overseed mixture. Apply the full 2 lb. a.i. rate in one application.
---	--------------------------------	-------------------------	---

NOTE: In many areas annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype, "Poa Annua spp. established when this product is applied" which germinates from seed every year is controlled, while the biotype, "Poa Annua spp. raptans" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning your local biotype.

Annual Weed Control during establishment from "Sprigs" of Bermudagrass, Zoysiagrass, and Seashore Paspalum.

Crabgrass and Goosegrass, Annual Sedge and other annual weeds.	2 to 3 lbs. a.i./A 200 to 300 lbs. per acre	4.6 to 6.9 lbs. per 1,000 sq.ft.	Directions - Refer to Map of U.S. on this label. For Bermudagrass make application immediately prior to, or, immediately after sprigging. For Seashore Paspalum and Zoysia make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkle irrigate the "sprigged" area with up to 1/4 inch water after application. RONSTAR® is a Registered Trademark of Aventis Corp.
--	--	----------------------------------	---

PRO SOURCE #1 Magic Carpet FERTILIZER WITH 1.00% IONSTAR®
EPA Reg. No. 1381-RIR EPA EST. NO. 1381-FL-1



1. Mid-north & northwestern After May 30
2. Upper/mid-states & northeast After May 10
3. Lower mid-states After April 20
4. Upper southern states After Mar 20
5. Warm season states Jan 1 to Mar 20

WEEDS CONTROLLED

- Annual Bluegrass
- Annual Sedge
- *Barnyardgrass
- Bittercress
- Carpetweed
- Carpetgrass
- Common Groundsel
- Common Purslane
- Crabgrass
- *Evening Primrose
- Fall Panicum
- Field Sandbur
- Fireweed
- Foxtail; Green, Bristly
- Florida Pusley
- *Galinsoga
- Golden Ragwort
- *Goosegrass
- **Junglerice
- Lambsquarter
- **Liverwort
- Oxalis
- **Panama Paspalum
- Pennsylvania Smartweed
- Redroot Pigweed
- Sensitive plant
- Shepardspurse
- Sow thistle
- Speedwell-veronica
- Spotted Catsear
- Spurge; Garden, Petty,
- **Prostrate
- Stinging Nettle
- *Swinecress
- Virginia Buttonweed
- Yellow Woodsorrel.

NON-AGRICULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and chemical resistant gloves when handling. Launder exposed clothing before reuse. Do not re-enter treated areas until dust has settled.

*For the weeds with 1 asterisk * above, use 3 to 4 lbs. a.i. per acre (equal to 300 lbs. to 400 lbs. of this product per acre or 6.9 lbs. to 9.2 lbs. per 1,000 sq.ft.).
For the difficult-to-control weeds with 2 asterisks ** above, use the 4 lbs. a.i. per acre (equal to 400 lbs. of this product/acre or 9.2 lbs. per 1,000 sq.ft.) **NOTE: For newly seeded lawns do not apply this product until after the second cutting. For best turf management, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied

FOR SPRING APPLICATION ON GOLF COURSES

The 3 lb. and 4 lb. a.i. per acre rates may be applied as a split application. Weed control and better turf management can be programmed by splitting the a.i. per acre rate of active ingredient into "two" separate applications. For example to accomplish this in region #5 (See map of U.S. on this label); first apply 1.5 lbs. a.i. (for required 3 lbs. a.i./acre rate) or first apply 2 lbs. a.i. acre (for a 4 lb. a.i./acre rate) as the 1st application 1 to 2 weeks before estimated weed seed germination or in late December or early January; Then follow with the second "1/2 rate" of a.i. in 60 days or the first of March. The timing for these applications should be closely programmed to the geographic region; (see map of U.S. on this label) and adjusted to an "early" or "late" spring. Do not split the 2 lb. a.i./acre rate. The 3 lb. a.i. / acre rate should not be split if applied to new turf areas first under turf management where weed pressure is expected to be heavy.

SPREADER SETTINGS

Spreader settings are calculated to apply the recommended rates of active ingredient per acre based on this product bulk density of 50 to 60 lbs. per cubic foot. Settings will vary if your turfgrass requires this EPA registered herbicide/fertilizer combination product with a lighter bulk density. To check your spreader settings refer to the spreader manual or manufacturer. Test apply enough product to cover 500 sq. ft. based on the amount of product a.i. you wish to apply per acre. Select a suitable rate from in the Pre-emergence Weed Control chart, on this label. Adjust and record the setting if required. Apply as directed on this label.

Spreader Settings	for 2 lbs. a.i./Acre	for 3 lbs. a.i./Acre	for 4 lbs. a.i./Acre
Cyclone B3-Pro	# x.xx	# x.xx	# x.x
Scotts Pro R-8A	# x-x	# xxx	# xxx

Plan on an approximately 3 mph spreader speed to obtain best coverage. Know the exact square footage before application. Incorrect application will result in poor weed control.

BERMUDA AND PASPALUM DIRECTIONS

Refer to U.S. Map on this label. For Bermudagrass make application immediately prior to or after sprigging. For Seashore Paspalum apply 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable paspalum injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" area with up to 1/4 inch water after application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

PRECAUTIONS FOR USING ON TURFGRASS AND ALSO PRECAUTIONS WHEN SEEDING

For St. Augustinegrass and Bentgrass turf, DO NOT exceed the rate of 3 lbs. of active ingredient per acre which is equal to 6.9 lbs. of this product per 1,000 sq.ft. DO NOT apply to Bentgrass turf mowed less than 3/8 inch. This product may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass but this discoloration is temporary and normally outgrown within 2 to 3 weeks. Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. **DO NOT** apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. • **DO NOT** apply to Dichondra or Centipedegrass. • **DO NOT** use on golf course putting greens or tees because of the varying cultural practices used.

Read the entire Directions for Use,

Conditions, Disclaimer of Warranties, and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties and Limitations of Liability.
CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However it is impossible eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Agrilience LLC, option, the replacement of the product.
Disclaimer of Warranties: There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Agrilience LLC, option, the replacement of the product, is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Agrilience LLC, disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product. **LIMITATION OF WARRANTIES:** The exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Agrilience LLC, option, the replacement of the product.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	variable	%
Nitrate Nitrogen	variable	%
Ammoniacal Nitrogen	variable	%
Water Soluble Nitrogen and/or Urea Nitrogen	variable	%
Water Insoluble Nitrogen	variable	%
Available Phosphoric Acid (P ₂ O ₅)	variable	%
Soluble Potash (K ₂ O)	variable	%
Derived from Ammonium Sulfate, Ureaform, Muriate of Potash; Manganese as Mn %; Iron as Fe %; Chlorine <		%

Agrilience LLC

P.O. BOX 64089 - ST. PAUL MN 55164-0089

FOR THE FOLLOWING EMERGENCY PHONE 24 HOURS A DAY
TRANSPORTATION: CHEMTREC - - - - 1-800-424-9300

