

1381-182

03/21/2005

1/12

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

☐ Registration  
☐ Amendment  
☒ Other

OPP Identifier Number

## Application for Pesticide - Section I

1. Company/Product Number 1381-182	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Magic Carpet Fertilizer w Barricade 0.21%	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Agrilliance, LLC, c/o Alice Walker Consulting 481 Country Club Drive Senatobia, MS 38668 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

## Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b>  MAR 21 2005
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification per PR Notice 98-10 to add language recommending that product be kept off concrete to avoid staining. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

## Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container
		If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	
		5. Location of Label Directions <input type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Alice Walker, Ph.D.	Title Agent	Telephone No. (Include Area Code) 662-562-5995
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Data Application Received  (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Alice Walker, Ph.D.	5. Date February 28, 2005	

2/12  
NOTIFICATION  
MAR 21 2005

# ProSource One

## Magic Carpet™ FERTILIZER WITH BARRICADE® 0.21%

**ACTIVE INGREDIENT:**

Prodiamine<sup>1</sup>.....0.21%

**OTHER INGREDIENTS:**.....99.79%

**TOTAL**.....100.00%

<sup>1</sup> CAS No. 29091-21-2

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials & wildflower plantings

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### FIRST AID

<b>If Swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If in Eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information. You may also contact 1-877-424-7452 for emergency medical treatment information.

EPA Reg. No. 1381-182

EPA Est. No. \_\_\_\_\_

Distributed By:

NET WEIGHT \_\_\_\_\_

Agrilience, LLC

P.O. Box 64089, St. Paul, MN 55164-0089

0/B28/5

3/12<sub>2</sub>

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

**If material is spilled:** Contain and sweep up material of spill for disposal as per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE**

Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides.

**PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL**

**Paper and plastic bags:** Do not reuse empty bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or at an incineration facility, or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns, landscape ornamentals, and established perennials & wildflower plantings.

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

**Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.**

**APPLICATION DIRECTIONS**

This product must be applied prior to crabgrass seed germination and prior to germination of seed of the other weeds listed on this label. Apply uniformly through suitable, calibrated application equipment.


### ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in Table 1. Maximum Annual Rates on this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

### CRABGRASS GERMINATION DATES

Approximate Date



### USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

1. Do not graze or feed livestock forage cut from areas treated with this product.
2. Do not use on food producing plants.
3. Do not apply this product through any type of irrigation system.
4. Do not apply aerially.
5. Do not apply to golf course putting greens.
6. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
7. Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
8. Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.
9. Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
10. Disturbing the herbicide barrier with cultural practices, such as disking may result in reduced weed control.
11. Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bentgrass (*Poa annua*) are desirable species.
12. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

### RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Do not exceed the Maximum Annual Rate per calendar year (See Table 1.)

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

### MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Table 1. Maximum Application Rates <sup>1</sup>		
Turf Species:	Lbs. Product/A	Lbs. a.i./A
Bermudagrass <sup>2</sup> Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	715 <sup>3</sup>	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	475	1.0
Creeping Red Fescue	355	0.75
Creeping Bentgrass	310	0.65

<sup>1</sup> These are the maximum rates per calendar year by species limitations.

<sup>2</sup> May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 248 lbs./A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

<sup>3</sup> Do not apply more than 715 lbs. /A (1.5 lb. a.i./A) of this product per calendar year.

- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in Table 1. Maximum Application Rates.

### WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Carpetweed
Bluegrass, Annual ( <i>Poa annua</i> )	Chickweed, Common
Crabgrass (large, smooth)	Chickweed, Mouseear (from seed)
Crowfootgrass	Henbit
Cupgrass, Woolly	Knotweed
Foxtails, Annual	Kochia
Goosegrass <sup>1</sup>	Lambsquarter, Common
Itchgrass	Pigweed
Johnsongrass (from seed)	Purslane, Common
Junglerice	Pusley, Florida
Lovegrass	Shepherd's purse <sup>2</sup>
Panicum (Texas, Fall, Browntop)	Speedwell, Persian
Rescuegrass <sup>3</sup>	Spurge, Prostrate

Signalgrass, Broadleaf  
Sprangletop

Witchgrass  
Woodsorrel, Yellow (from seed)

<sup>1</sup>In many areas a single application of 310 to 715 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 310 to 475 lbs./A followed after 40 to 60 days by a second application at doses that would not exceed those given in Table 1. **Maximum Application Rates.** Do not exceed the maximum rate for turf grass species listed in Table 1. **Maximum Application Rates.**

<sup>2</sup>Applications for Shepherd's purse should be made in late summer, fall, or winter prior to germination.

<sup>3</sup>Suppression only on rescuegrass.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

### WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

**TABLE 2. Months After Application Before Overseeding.**

Rate Lbs. Product/A	North	Transition	South
240	4	4	4
310	5	4	4
355	6	5	5
380	---	6	6
475	---	7	7
540	---	---	9
620	---	---	10
715	---	---	12

### LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

### USE RATES

Apply this product at 310 to 715 lbs./A (7 lbs. to 16 lbs. per 1,000 sq. ft.) of landscape plants in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 715 lbs./A.

### APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. Calibrate granular applicator per manufacturer's instructions to assure appropriate dosage. To assure that the product is applied at the proper rates, apply at a rate of 4 lbs. per 1,000 sq. ft. or for flowerbeds, use 6.4 oz. per 100 sq. ft.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate post-emergence herbicide.

### ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

### NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

### TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

#### Scientific name

*Abies* spp.  
*Abelia grandiflora*  
*Acer palmatum*  
*A. platanoides*  
*Achillea* spp.  
*Actinidia chinensis*  
*Agapanthus africanus*  
*Agapanthus orientalis*  
*Akebia quintata*  
*Allium cernuum*  
*Anemone hybrida*  
*Aquilegia* spp.  
*Arctostaphylos densiflora*  
*Arctotheca calendula*  
*Artemisia* spp.  
*Aster* spp.  
*Aster X Frikartii*  
*Athrium Filiz-femina*  
*Aucuba japonica*  
*Begonia* spp.  
*Berberis gladwynensis*  
*B. julianae*  
*B. mentorensis*  
*B. thunbergii*  
*B. verruculosa*  
*Bergenia cordifolia*  
*Boltonia asteroides*  
*Bougainvillea* spp.  
*Buddleia davidii*  
*Buxus microphylla*  
*Callistemon viminalis*  
*Calluna vulgaris*  
*Campanula carpatica*  
*Campis X Tagliabuana*  
  
*Carpobrotus edulis*  
*Cassia artemisoides*  
*Ceanothus rigidus*

#### Common name

Fir species\*\* (Balsam, Fraser, Noble, etc.)  
 Abelia: Sherwood  
 Japanese Maple  
 Norway Maple  
 Yarrow: King Edward  
 Kiwi\*  
 Lily-of-the-Nile (African Lily)  
  
 Five-Leaf or Chocolate Vine  
 Lady's Leek, Nodding Onion  
 Japanese Anemone  
 Aquilegia: Red and Gold  
 Vine Hill Manzanita  
 Cape Weed  
 Wormwood: Silver Mound, Castle  
 Aster: Bonny Blue, Purple Dome  
  
 Lady Fern: Fern Lady  
 Japanese Aucuba  
 Fibrous Begonia: Hardy Grandis  
 Barberry  
 Wintergreen Barberry  
 Mentor Barberry  
 Japanese Barberry  
 Warty Barberry  
  
 Snowbank  
 Bougainvillea  
 Butterfly-Bush (Dwarf Blue): Royal Red  
 Japanese Boxwood  
 Weeping Bottlebrush  
 Scotch Heather  
 Tussock Bellflower, (White Clips)  
 Trumpet Creeper, Trumpet Flower,  
 Madame Galen  
 Hottentot Fig (Ice Plant)  
 Feathery Cassia  
 Wild Lilac

*Ceratostigma plumbagonoides*

*Chamaecyparis pisifera*

*Chrysanthemum nipponicum*

*Cleyera japonica*

*Citrus* spp.

*Coreopsis* spp.

*Cornus florida*

*C. stolonifera*

*Cortaderia selloana*

*Cotoneaster apiculatus*

*C. buxifolius*

*C. dammeri*

*C. microphyllus*

*Cretagus* spp.

*Crocasmia* spp.

*Cupressus sempervirens*

*Delosperma alba*

*Delosperma* spp.

*Delphinium* spp.

*Dianthus deltoides*

*D. gratianopolitanus*

*Dodonea viscosa*

*Echinacea purpurea*

*Elaeagnus pungens*

*Euonymus fortunei*

*E. japonica*

*E. kiautschovica*

*Fatsia japonica*

*Forsythia intermedia*

*F. suspensa*

*F. viridissima*

*Gaillardia* spp.

*Gardenia jasminoides*

*Gaura* spp.

*Gentiana dahurica*

*Geranium cinereum*

*Gladiolus* spp.

*Gypsophila repens*

*Hedera helix*

*Hellanthemum* spp.

*Hemerocallis* spp.

*Hibiscus*

*Hibiscus* spp.

*Hibiscus rosa-sinensis*

*Hosta plantaginea*

*H. Sieboldiana*

*Houttuynia cordata* var. *variegata*

*Hydrangea macrophylla*

*Ilex cornuta*

*I. crenata*

*I. opaca*

*I. pemyi*

*I. vomitoria*

*Inula ensifolia*

*Iris ensata*

*Iris siberica*

*Iris* spp.

*Jasminium nudiflorum*

False Cypress

Cleyera

Citrus species\*

Coreopsis (Calliopsis): Early Sunrise, Moonbeam

Flowering Dogwood

American Dogwood

Pampas Grass

Cranberry Cotoneaster

Cotoneaster

Bearberry Cotoneaster

Rockspray Cotoneaster

Hawthorn

Lucifer

Italian Cypress

White Trailing Ice Plant

Cooperi Pink

Larkspur: Blue Elf

Dianthus: Maiden Pinks "Zing"

Cheddar Pink

Hop Bush

Coneflower, Purple; Magnus

Silverberry

Wintercreeper

Japanese Spindle Tree (Evergreen Euonymus)

Spreading Euonymus

Japanese Aralia

Border Forsythia

Weeping Forsythia

Greenstem Forsythia

Gaillardia, Blanket Flower: "Goblin"

Gardenia, Cape-Jasmine, Radicans

Gentian

Cranesbill

Gladiolus species\*\*

Baby's Breath

English Ivy

Sunrose

Daylily: Aztec Gold, Stella de Oro, Tender Love

Rose of Sharon\*\*

Mallow: Disco Belle White

Chinese Hibiscus

Hosta, Plantain Lily (Fragrant)

Hosta, "Searsucker"

Bigleaf Hydrangea

Chinese Holly

Japanese Holly: Helleri

American Holly

Holly

Yaupon Holly: Schillings

Sword-Leaved Iris: Jodlesong

Siberian Iris: Cabernet

Iris species\*\*

Winter Jasmine

<i>Juniperus chinensis</i>	Chinese Juniper: Nick's Compact, Parsonnii
<i>J. conferta</i>	Shore Juniper: Blue Pacific
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> spp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica</i> X <i>Fauriei</i>	Crape Myrtle: Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender</i> spp.	Lavender: Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet: Variegata
<i>Ligustrum amurense</i>	Amur Privet
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> spp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily; Turf: Silver Mound: Evergreen Giant
<i>Liriope Muscari</i> var. <i>variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum</i> spp.	Loosestrife; Modern Pink
<i>Magnolia</i> spp.	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus</i> spp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	Avocado*
<i>Persea americana</i>	Fraser's Photinia (Redtip)
<i>Photinia fraseri</i>	Dragonhead, False: Vivid
<i>Physotegia Virginiana</i>	Spruce species**(Colorado Blue, Norway, etc.)
<i>Picea</i> spp.	Lily-of-the-Valley Bush
<i>Pieris japonica</i>	Calabrian Pine
<i>Pinus brutia</i>	Canary Island Pine
<i>P. canariensis</i>	Slash Pine
<i>P. elliotii</i>	Aleppo Pine
<i>P. halepensis</i>	Austrian Pine
<i>P. nigra</i>	Longleaf Pine
<i>P. palustris</i>	Monterey Pine
<i>P. radiata</i>	Japanese Black Pine
<i>P. thunbergiana</i>	Eastern White Pine
<i>P. strobus</i>	Scotch Pine
<i>P. sylvestris</i>	Loblolly Pine
<i>P. taeda</i>	Virginia Pine
<i>P. virginiana</i>	Pistachio*
<i>Pistachio</i> spp.	Queensland Pittosporum
<i>Pittosporum rhombifolium</i>	

10/12<sup>9</sup>

*P. tobira*  
*Podocarpus macrophyllus*  
*Prunus laurocerasus*  
*Prunus* spp.

*Pseudotsuga menziesii*  
*Pyracantha coccinea*  
*P. fortuneana*  
*P. koidzumii*  
*Pyrus* spp.  
*Quercus Shumardii*  
*Quercus* spp.  
*Raphiolepis indica*  
*Raphiolepis umbellata*  
*Rhododendron*  
 (including Azalea)

*Rosmarinus officinalis*  
*Rudbeckia* spp.  
*Santolina virens*  
*Saxifraga* spp.  
*Scabiosa* spp.  
*Sedum album*  
*Sedum cauticola*  
*Sedum dasyphyllum*  
*Sedum spurium*  
*Spiraea Bumalda*  
*Syzygium paniculatum*  
*Taxus cuspidata*  
*T. media*  
*Teucrium* spp.  
*Thalictrum dipterocarpum*  
*Thuja occidentalis*  
*Trachelospermum asiaticum*  
*Tsuga canadensis*  
*Tulipa* spp.  
*Veronica* spp.  
*Viburnum japonicum*  
*V. odoratissimum*  
*V. plicatum*  
*V. rigidum*  
*V. suspensum*  
*V. tinus*  
*V. trilobium*  
*V. wrightii*  
*Vinca major*  
*Vinca minor*  
*Vitis* spp.  
*Weigela florida*  
*Yucca aloifolia*  
*Y. filamentosa*

Japanese Pittosporum  
 Japanese Yew  
 English Laurel  
 Almond, Apricot, Nectarine,  
 Peach, Plum and Prune\*  
 Douglas Fir\*\*  
 Firethorn (Scarlet)  
 Firethorn  
 Firethorn  
 Bradford Pear sp.  
 Oak, Shumard's Red  
 Oak species  
 Indian Hawthorn  
 Yedda Hawthorn  
 Coral Bells, Delaware Valley White, Flame  
 Creeper, Formosa, George L. Tabor, Girard  
 Crimson, Hino-crimson, PJM, Roseum Elegans,  
 Wakeiebisu, White Gumpo  
 Rosemary\*  
 Black-Eyed Susan; Goldstrum  
 Santolina  
 Saxifrage: Purple Dome  
 Pincushion Flower  
 Stonecrop  
 Stonecrop: Lidakense  
 Stonecrop  
 Stonecrop: Dragon's Blood  
 Spirea: Anthony Waterer  
 Australian Brushcherry  
 Japanese Yew  
 Yew  
 Germander  
 Meadow Rue  
 American Arborvitae  
 Star Jasmine  
 Canada Hemlock  
 Tulip species  
 Veronica, Speedwell: Sunny Border  
 Japanese Viburnum  
 Sweet Viburnum  
 Japanese Snowball  
 Canary Island Viburnum  
 Arrowwood Viburnum  
 Laurustinus  
 Cranberry Bush  
 Leatherleaf Viburnum  
 Greater Periwinkle, Vinca  
 Common Periwinkle, Vinca  
 Grape\*  
 Old Fashioned Weigela  
 Spanish Bayonet, Yucca  
 Adam's Needle, Yucca

\*Do not use on food-producing trees, vines, or plants.

\*\*Not for use on container-grown plants.

**Notice of Warranty:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER

11/12<sub>10</sub>

EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

Barricade® is a registered trademark of Syngenta.  
Magic Carpet™ is a trademark of Agrilience, LLC.

**GUARANTEED FERTILIZER ANALYSIS: (To be attached to final printed labeling)**

<b>X-X-X *</b>	
Total Nitrogen (N) .....	X.XX%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....	X.XX%
Soluble Potash (K <sub>2</sub> O).....	X.XX %
Chlorine not more than X.X%	
Derived from xxxxxxxxxxxxxx.	

\* [Editor's Note: all information will vary depending on grade used]

**CERTIFIED MAIL**

February 28, 2005

Document Processing Desk (Notif)  
Office of Pesticide Programs (7504C)  
U.S. Environmental Protection Agency  
1200 Pennsylvania Ave., NW  
Washington, DC 20460

Re: Magic Carpet Fertilizer with Barricade 0.21%, EPA Reg. No. 1381-182  
Notification under PRN 98-10 of Addition of Language Regarding Possible Staining of  
Concrete

Attn: Sherada Hobgood

This correspondence will constitute notification from AGRILIANCE, LLC to add label language recommending that the product be kept off concrete to avoid staining. We have highlighted the new text on page 3 for your convenience. Following a telecon with Jim Stone on this date, we are submitting this label revision as a notification rather than an amendment.

I believe this notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Thank you for adding this notification to the registration records for this product. We would appreciate a stamped "Notification" copy for our files.

Sincerely,



Alice Walker, Ph.D.  
Regulatory Agent for  
AGRILIANCE, LLC

Enclosures

cc: Mr. Gary Halvorson

