

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 13, 2014

Danielle Erding
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164

Subject: Label Notification per PRN 2001-05 – Add Resistance Management Language
Product Name: BATTERY 2.5 EC
EPA Registration Number: 1381-188
Application Date: October 6, 2014
Decision Number: 496690

Dear Ms. Erding:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2001-05 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 2001-05 and finds that the action requested falls within the scope of PRN 2001-05.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

If you have any questions, you may contact Maggie Rudick at 703-347-0257 or via email at rudick.maggie@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", enclosed within a hand-drawn oval.

Kable Bo Davis, Product Manager 3
Invertebrate & Vertebrate Branch 1
Registration Division (7505P)
Office of Pesticide Programs

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Invertebrates

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification

GROUP

3

INSECTICIDE

BATTERY[®] 2.5EC Insecticide

Active Ingredient

*Cypermethrin** 30.6%

Inert Ingredients*** 69.4%

Total 100.0%

*(±) α-Cyano (3-phenoxyphenyl)methyl(±) *cis/trans* 3-(2,2-dichloroethenyl)-2,2 dimethylcyclopropanecarboxylate

**Cis/trans ratio: Max. 55% (±) *cis* and min. 45% (±) *trans*

*** Contains petroleum distillates.

Contains 2.5 pounds cypermethrin per gallon.

NOTIFICATION

1381-188

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

11/13/2014

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.

FIRST AID

| | |
|-------------------------------|---|
| IF SWALLOWED | <ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| IF IN EYES | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| IF INHALED | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300

EPA Reg. No. 1381-188

EPA Reg. Nos. 53883-TX-1, 279-FL-1, 70989-AR-1, 70989-MO-1

Distributed By:

NET CONTENTS Gals. (Liters)

Winfield Solutions, LLC

P.O. Box 64589

St. Paul, MN 55164-0589

1/0613/3 1/1006/4

WINFIELD

AgriSOLUTIONS
THE

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Warning: May be fatal if swallowed. Causes skin irritation. Harmful if absorbed through skin. Causes moderate eye irritation. Do not get on skin or clothing. Avoid contact with eyes.

Product should be opened and poured in a well-ventilated area.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are made of barrier laminate, butyl rubber, nitrile rubber, or viton. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Pilots must wear:

- Long-sleeved shirt and long pants, and
- Shoes and socks.

Mixers, loaders, other applicators and handlers must wear:

- Coveralls over short-sleeved shirt and short pants,
- Chemical-resistant footwear plus socks,
- Chemical-resistant gloves,
- Chemical-resistant headgear for overhead exposure, and
- Chemical-resistant apron when mixing, loading or cleaning equipment.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

See engineering controls for additional requirements.

User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

User Safety Recommendations

Users should:

Wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Physical/ Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~**Resistance.** Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.~~

~~If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for our area. [editors note: moved below the next paragraph]~~

RESISTANCE MANAGEMENT RECOMMENDATIONS

Battery 2.5EC contains a Group 3 insecticide (or acaricide). Insect/mite biotypes with acquired resistance to Group 3 may eventually dominate the insect/mite population if Group 3 insecticides/acaricides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Battery 2.5EC or other Group 3 insecticide (or acaricide).

~~If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for our area.~~

~~To delay insecticide (or acaricide) resistance consider:~~

- ~~• Avoiding the consecutive use of Battery 2.5EC or other group 3 insecticides/acaricides that have a similar target site of action, on the same insect/mite species.~~
- ~~• Using tank-mixtures or premixes with insecticides/acaricides from a different target site of action Group as long as the involved products are all registered for the same use and have different sites of action.~~
- ~~• Basing insecticides/acaricide use on a comprehensive IPM program.~~
- ~~• Monitoring treated insect/mite populations for loss of field efficacy.~~
- ~~• Contacting your local extension specialist, certified crop advisors, and/or Winfield Solutions, LLC representative for insecticides/acaricide resistance management and/or IPM recommendations for the specific site and resistant pest problems.~~

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Shoes plus socks, Chemical-resistant gloves made of any waterproof material.

CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform, distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Battery 2.5EC should be applied continuously for the duration of the water application. Battery 2.5EC should be diluted in sufficient volume to insure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation is not required when a suitable diluent is used.

PRODUCT INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

Rotational Crops

With the exception of the crops listed below, rotational crops may not be planted within 30 days after last application.

Tank Mixture

Battery 2.5EC may be applied in tank mixtures with other products approved for use on cotton, brassicas, bulb vegetables, head lettuce, and pecans. Observe all restrictions and precautions which appear on the labels of these products.

For improved control of aphids and mites on cotton, Battery 2.5EC may be tank-mixed with Curacron® 8E. The tank-mixture is most effective when pest populations are first observed. Subsequent applications may be needed to maintain control.

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing cypermethrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses*. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp. <http://www.in.csusda/v/technical/agronom/newconbuf.pdf>.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application

Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial application

Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

SPRAY DRIFT REQUIREMENTS

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size. Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Maximum Usage When Applying Both Zeta-Cypermethrin and Cypermethrin Products to the Same Crop within the Same Season or Year, as indicated.

Do not apply more than the maximum seasonal or yearly total for either product when used alone, and do not apply more than the combined maximum seasonal total for both products as outlined in the table below.

| Crop | Maximum Seasonal or Yearly Total for Either Product Used Alone (pounds active ingredient/acre) | | Maximum Seasonal Total When Applying Both Products to the Same Crop (pounds ai/acre) |
|------------------------|--|------------------------------|--|
| | Zeta-cypermethrin Maximum Seasonal Total | Cypermethrin (Battery 2.5EC) | Zeta-cypermethrin plus Cypermethrin (Battery 2.5EC) |
| Cotton | 0.3 | 0.4 lbs. per year | 0.6 |
| Head and Stem Brassica | 0.3 | 0.6 per season | 0.6 |
| Leafy Brassica Greens | 0.3 | 0.4 per season | 0.4 |
| Head Lettuce | 0.3 | 0.6 per season | 0.6 |
| Bulb Vegetables | 0.25 | 0.5 per season | 0.5 |
| Pecans | 0.3 | 0.5 lbs. per year | 0.6 |

Cotton (14 day phi)

| PEST | DOSAGE | | REMARKS |
|--|--------------|-----------|--|
| | LB/AI/A | FL. OZ./A | |
| Preemergent Use: Cutworms | 0.025 to 0.1 | 1.3 to 5 | Use Battery 2.5EC in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air, banded (including T-brand) or in-furrow spray using sufficient spray volume to achieve adequate coverage. Use a minimum of 1 gallon of water per acre by air. Reduced volumes of water may be used with specialized equipment. Use the higher rates of Battery 2.5EC when incorporating into the soil. |
| Foliar Use: Cutworms Tobacco Thrips Soybean (banded) Thrips | 0.025 to 0.1 | 1.3 to 5 | |
| Foliar use: Boll Weevil Cabbage Looper Cotton Bollworm Cotton Fleahopper Cotton Leaf Perforator European Corn Borer Fall Armyworm Lygus Bugs Other Plant Bugs Pink Bollworm Saltmarsh Caterpillar Tarnished Plant Bug Tobacco Budworm White Flies* Yellow Striped Armyworm | 0.04 to 0.1 | 2 to 5 | Battery 2.5EC may be applied in water or refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment. When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray. When using oil, use a minimum of one quart per acre in the finished spray. Battery 2.5EC may be injected into overhead sprinkler irrigation water provided 1) an anti-backflow check valve is present between the injection port and the water source, 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the irrigation injection system has interlocking on-off switches. |

| | | | |
|--------------------------------|-------------|--------|--|
| Foliar Use: Beet Armyworm** | 0.06 to 0.1 | 3 to 5 | <p>For boll weevil control, re-apply Battery 2.5EC until pest numbers are reduced to acceptable levels.</p> <p>All use patterns: Do not make applications less than 5 days apart.</p> <p>A maximum of 0.4 lb. Active ingredient (20 fl. oz. of Battery 2.5EC) may be applied per acre per year.</p> <p>Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Asana® XL, Baythroid®, Capture®, Danitol®, Karate®, Scout® X-Tra, and Warrior®.</p> <p>Do not graze or feed cotton for forage.</p> |
|--------------------------------|-------------|--------|--|

*Aids in control. **For control of beet armyworms only in the high plains of Texas, Arizona, and California.

Head and Stem Brassica Vegetables including: Broccoli; Chinese Broccoli (gai lan, white flowering broccoli); Brussels Sprouts; Cauliflower; Cavolo broccolo, Kohlrabi; Cabbage; Chinese Cabbage (napa) (tight-heading varieties only) (1 day phi)

| PEST | DOSAGE | | REMARKS |
|--|--------------|-----------|---|
| | LB/AI/A | FL. OZ./A | |
| Cutworms Corn Earworm Tobacco Budworm Lygus Bugs Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids* Whiteflies* | 0.05 to 0.1 | 2.5 to 5 | <p>Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.</p> <p>Lower rates of Battery 2.5EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. Do not make applications less than 7 days apart.</p> <p>A maximum of 0.6 lb. active ingredient (30 fl. oz. of Battery 2.5EC) may be applied per acre per season.</p> <p>In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.</p> |
| Armyworms Loopers Stinkbugs Crickets Ground Beetles Onion Thrips Wireworm (adults) | 0.075 to 0.1 | 3.75 to 5 | <p>Follow appropriate spray drift precautions listed for cotton, brassicas, lettuce, and onions.</p> <p>*aids in control</p> |

**Leafy Brassica Greens including: Broccoli Raab (rapini), Chinese cabbage (bok choy); Chinese Mustard Cabbage (gai choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens.
(1 day phi)**

| Pest | DOSAGE | | Remarks |
|---|--------------|-----------|--|
| | LB/AI/A | FL. OZ./A | |
| Cutworms Corn Earworm Tobacco Budworm Lygus Bugs Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids* Whiteflies* | 0.05 to 0.1 | 2.5 to 5 | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Battery 2.5EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. Do not make applications less than 7 days apart. A maximum of 0.4 lb. active ingredient (20 fl. oz. of Battery 2.5EC) may be applied per acre per season. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. |
| Armyworms Loopers Stinkbugs Crickets Ground Beetles Onion Thrips Wireworm (adults) | 0.075 to 0.1 | 3.75 to 5 | Follow appropriate spray drift precautions listed for cotton, brassicas, lettuce and onions. *aids in control |

Lettuce, Head. (5 day phi)

| PEST | DOSAGE | | REMARKS |
|---|--------------|-----------|---|
| | LB/AI/A | FL. OZ./A | |
| Corn Earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Lygus Bugs Saltmarsh Caterpillar Tobacco Budworm | 0.05 to 0.1 | 2.5 to 5 | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Battery 2.5EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. Do not make applications less than 7 days apart. A maximum of 0.6 lb. active ingredient (30 fl. oz. of Battery 2.5EC) may be applied per acre per season. |
| Armyworms Crickets Loopers Onion Thrips Stinkbugs | 0.075 to 0.1 | 3.75 to 5 | In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions listed for cotton, brassicas, lettuce, and onions. |

Bulb Vegetables (*Allium* spp.) including: Garlic; Garlic, great-headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions. (7 day phi)

| PEST | DOSAGE | | REMARKS |
|--|-------------|-----------|---|
| | LB/AI/A | FL. OZ./A | |
| Onion Thrips | 0.08 to 0.1 | 4 to 5 | Apply Battery 2.5EC in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear. Do not make applications less than 7 days apart. A maximum of 0.5 lb. active ingredient (25 fl. oz. of Battery 2.5EC) may be applied per acre per season. To control Onion Thrips: Use higher rates as population increases and avoid rescue situations. Use of a crop oil concentrate at 16 fl. Oz/A is recommended. Do not graze livestock in treated areas or cut treated crops for feed. Follow appropriate spray drift precautions listed for cotton, brassicas, lettuce and onions. |
| Aphids Armyworms Cutworms Leafminers Onion Maggot Adults Stink Bugs | 0.04 to 0.1 | 2 to 5 | |

Pecans (21 day phi)

| PEST | DOSAGE | | REMARKS |
|---|-----------------|-----------|---|
| | LB/AI/A | FL. OZ./A | |
| Black Pecan Aphid Hickory Shuckworm Pecan Nut Casebearer Pecan Weevil Yellow Pecan Aphid | 0.06 to 0.10 | 3 to 5 | Applications at the lower rate should be made when pest populations are low. Rates should be increased as the pest pressure increases. Do not make applications less than 7 days apart. Apply by ground equipment to the point of drip. Use 100 gallons of dilute spray per acre for smaller trees. For larger trees which require higher gallonage to achieve adequate coverage, apply in 200 to 300 gallons of water. In order to calculate the correct number of gallons of water needed to spray one acre of your trees to the point of drip, you may need to conduct a test. If you do not know how to conduct such a test with your equipment, you should request assistance from your equipment dealer. Up to 0.5 lb. active ingredient (25 fl. oz. of Battery 2.5 EC) per acre season may be applied prior to shuck split. Do not graze livestock in treated orchards or cut treated cover crops for feed. |

STORAGE AND DISPOSAL

Pesticide Storage

Do not store below 10°F, (-12°).

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Do not use or store near heat or open flame. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal: Use label language appropriate for container size and type.

Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call
CHEMTREC 1-800-424-9300**

Dealers Should Sell in Original Packages Only.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE

GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Ambush, Curacron, Karate and Warrior-Trademarks of Syngenta Crop Protection.

Asana-Trademark of E.I. Dupont de Nemours and Co., Inc.

Baythroid-Trademark of Bayer AG

Scout-Trademark of Hoechst Roussel Agri-Vet Company

Capture-Trademark of FMC Corporation

Danitol-Trademark of Valent U.S.A. Corporation

AgriSolutions™ is a trademark and Battery® is a registered trademark of Winfield Solutions, LLC.