

1381-194

11/23/2010

1/52



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Ms Diana Williams
Winfield Solutions, LLC
P O Box 64589
St Paul, MN 55164-0589

NOV 23 2010

Subject Notification per PR Notice 98-10
Charger Max
EPA Reg No 1381-194
Application dated October 13, 2010
Text corrections per Notice of Registration

Dear Ms Williams,

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 of the subject product The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10 The label submitted with the application has been date-stamped "Notification" and will be added to our records

If you have any questions, please contact Hope Johnson at 703-305-5410

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2/52

<h1 style="margin:0;">EPA</h1>	United States Environmental Protection Agency Washington DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--------------------------------	--	--	-----------------------

Application for Pesticide -Section I

1 Company/Product Number 1381 194	2 EPA Product Manager Kathryn V. Montague	3 Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) Charger Max	PM# 23	
5 Name and Address of Applicant (Include ZIP Code) Winfield Solutions LLC P O Box 64589 St Paul MN 55164 0589 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to EPA Reg No _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment Explain Below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other explain below

Explanation Use additional page(s) if necessary (For Section I and Section II)
 Notification to correct text per EPA Notice of Registration of Sept 19 2005 I believe this notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA

Section III

1 Material This Product Will Be Packaged In

Child Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No <i>Certification must be submitted</i>	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If Yes Unit Package Wt _____ No Per Container _____	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If Yes Package Wt _____ No Per Container _____	2 Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3 Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4 Size(s) of Retail Container	5 Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6 Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input type="checkbox"/> Paper Glued <input type="checkbox"/> Stenciled			

Section IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)

Name Diana Williams	Title Product Registration	Telephone No (Include Area Code) 651 494 5205
------------------------	-------------------------------	--

<p style="text-align: center;">Certification</p> I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6 Date Application Received (Stamped)
2 Signature 	3 Title Product Registration on Behalf of Winfield Solutions LLC	
4 Typed Name Diana Williams	5 Date October 13 2010	



October 13, 2010

Ms Kathryn V Montague
Product Manager (23)
Herbicide Branch
Registration Division (7505C)
Document Processing Desk (NOTIF)
U S EPA – OPP
1200 Pennsylvania Ave NW
Washington D C 20460

Subject Notification of Text Corrections
EPA Reg No 1381-194 Charger Max

Dear Ms Montague

We recently discovered that the label copies submitted with Notifications filed November 19 2007 and April 16, 2008 did not include all of the corrections noted in the Agency’s September 19 2005 Notice of Pesticide Registration We are herewith correcting this oversight by Notification of the inclusion of the following statements into the label on file with the Agency

On page 4 addition of This product may be stored at temperatures down to 30 degrees below 0 degrees F to the pesticide storage statement

On page 10 addition of “soil before “surface in the statement i e untreated soil should not be brought to soil surface ” under ‘Special Application Procedures California only corn, safflower pod crops Preplant incorporated ’

On page 48, addition of “Use only where soils have 0 5-5 0% organic matter, and do not apply to soils with pH greater than 6 8 Do not apply to sand ” Under the “Pursuit + paraquat or Cornerstone (glyphosate)” tank mix directions

Enclosed are a Notification application and five copies of corrected labeling The corrections have been highlighted on one copy of the labeling

If there are any questions concerning the enclosed, please do not hesitate to contact me

Sincerely

Diana Williams
Product Registration
On Behalf of Winfield Solutions LLC

Enclosures

Cc Alice C Walker PhD Winfield Solutions LLC

4/52



NOTIFICATION
NOV 23 2010

**HERBICIDE FOR WEED CONTROL IN CORN, COTTON, GRAIN OR FORAGE
SORGHUM, SOYBEANS, SAFFLOWER, PEANUTS, POD CROPS, AND POTATOES**

ACTIVE INGREDIENT

s-metolachlor ((s) 2 chloro n (2-ethyl 6-methylphenyl) n-(2 methoxy 1-methylethyl) acetamide) 82.4%

OTHER INGREDIENTS

17.6%

TOTAL

100.0%

Contains 7.64 lb s-metolachlor per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, mouth-to-mouth if possible. Call a Poison Control Center or doctor for further advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.</p>	

**SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND USE
DIRECTIONS**

EPA Reg No 1381-194

EPA Est No 11773-IA-01

Manufactured for
Winfield Solutions LLC
P O Box 64589
St Paul MN 55164-0589

NET CONTENTS 2.5 Gals
LOT NO _____
1/1013/0

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Causes eye irritation Do not get in eyes on skin or on clothing Harmful if swallowed or absorbed through skin This product may cause skin sensitization in some people

You may also contact (Poison Control Centers) 800-222-1222
(ASPCA – animal health) 800-345-4735

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants chemical resistant gloves such as barrier laminate or Viton and shoes plus socks Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

Engineering controls statement

Mixers and loaders supporting aerial applications are required to use closed systems The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4-6)] When using the closed system the mixers and loaders PPE requirements may be reduced or modified as specified in the WPS

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240 (d) (4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Recommendations

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash and change into clean clothing

ENVIRONMENTAL HAZARDS

Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water when cleaning equipment or disposing of equipment washwaters Use only as directed on this label

Ground Water Advisory

The active ingredient in Charger Max has the potential to leach through soil into ground water under certain conditions as a result of agricultural use Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in ground water contamination

Surface Water Advisory

The active ingredient in Charger Max has the potential to contaminate surface water through spray drift. Under some conditions, the active ingredient may have a high potential for runoff into surface water (primarily via dissolution into runoff water) for several months post-application. These include poorly-drained or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain into surface waters, areas not separated from adjacent surface waters by vegetated surface strips, and areas overlying tile drainage systems that drain into surface water.

Mixing/ Loading Instructions for Water Protection

To prevent point-source contamination, do not mix and load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams or rivers, or natural or impounded lakes or reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells, and does not apply to impervious pads or properly diked mixing/ loading areas as described below. Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% of that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment, or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/ loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment. Care must be taken with this product to prevent a) back-siphoning into wells, b) spills, or c) improper disposal of excess pesticide spray mixtures or rinsates. Check valves or anti-siphoning devices must be used on all mixing equipment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NOTE NOT FOR SALE, USE, OR DISTRIBUTION IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. EXCEPTION: If this product is soil-incorporated or soil-injected, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves such as barrier laminate or Viton, and shoes plus socks.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, AND/ OR ILLEGAL RESIDUES

To avoid spray drift, do not apply under windy conditions. Avoid spray overlaps, as crop injury may result.

STORAGE AND DISPOSAL

Prohibitions Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. Open dumping is prohibited. This product may not be mixed, loaded, or used within 50 feet of wells, including abandoned wells, drainage wells, and sinkholes.

Pesticide Storage Store in original container in a well-ventilated area separate from fertilizer, food, and feed. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent for disposal. The risk of groundwater contamination will be reduced by diking and flooring permanent liquid storage sites with impermeable material. This product may be stored at temperatures down to 30 degrees below 0 degrees F.

Pesticide Disposal Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

CONTAINER DISPOSAL [Use label language appropriate for container size and type]

Nonrefillable containers Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable container Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY Spill, leak, fire, exposure, or accident, call
CHEMTREC 1-800 424-9300

GENERAL INFORMATION

Charger Max is a herbicide recommended as a preplant surface-applied preplant incorporated or pre-emergence treatment in water or fluid fertilizer for control of most annual grasses and some broadleaf weeds in corn (all types) cotton peanuts pod crops potatoes safflower grain or forage sorghum and soybeans

Follow all Directions for Use found on the labeling of pesticide products used in tank mixture with Charger Max. Observe all precautions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those States where the tank mix partner is registered.

DO NOT USE IN NURSERIES TURF OR LANDSCAPE PLANTINGS

Do not apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.

To prevent off site movement due to runoff or wind erosion:

- Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions the soil surface should first be settled by rainfall or irrigation.
- Do not apply to impervious substrates such as paved or highly compacted surfaces.
- Do not use tailwater from first flood or furrow irrigation of treated fields to treat nontarget crops unless at least 1/2 inch of rainfall has occurred between application and first irrigation.

Notes, atrazine tank mixtures Follow rates, recommendations, and limitations of the atrazine label being used. Certain States may have established rate limitations for atrazine within specific geographic areas. Consult your State pesticide control agency for additional information. It is a violation of law to deviate from State use regulations. If Charger Max is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Dry weather following an application of Charger Max alone or in tank mixture may reduce effectiveness. Cultivate if weeds emerge.

Where this label refers to partial control of weeds, partial control means either erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial weed control.

Precaution Injury may occur to crops other than corn following the use of Charger Max under abnormally high soil moisture conditions during early development of the crop.

SOIL TEXTURES AND APPLICATION RATES

Soil textural classes are categorized as in the following table

Coarse	Medium	Fine
Sand	Loam	Sandy clay loam
Loamy sand	Silt loam	Silty clay loam
Sandy loam	Silt	Clay loam
		Sandy clay
		Silty clay
		Clay

Within rate ranges in the rate tables and elsewhere in this label use the lower rate on soils that are relatively coarse textured or low in organic matter Use the higher rate on soils that are relatively fine textured or high in organic matter

Charger Max may be applied pre-emergence alone or in tank mixtures as specified in this label following pre-plant incorporated herbicides used according to their label directions provided such use is not prohibited by those labels

Thoroughly clean sprayer or application device before using Dispose of cleaning solution in a responsible manner Do not use a sprayer or application device contaminated with any other materials to prevent crop damage or clogging of the application device

WEEDS CONTROLLED BY CHARGER MAX ALONE

Weeds Controlled

barnyard grass (water grass)	prairie cupgrass	yellow foxtail
bristly foxtail	red rice	yellow nutsedge
crabgrass	robust foxtails (purple white)	carpetweed
crowfoot grass	signalgrass (Brachiaria)	common waterhemp
fall panicum	southwestern cupgrass	eastern black nightshade
foxtail millet	wild proso millet*	Florida pusley
giant foxtail	witchgrass	galinsoga
goosegrass	wooly cupgrass*	pigweed
green foxtail		tall waterhemp

* For control of these weeds in corn only refer to the Corn - wooly cupgrass and wild proso millet control program section of this label

Weeds Partially Controlled***

common purslane	sandbur	volunteer sorghum
eclipta	seedling Johnsongrass	wild proso millet
Florida beggarweed**	shattercane	wooly cupgrass
hairy nightshade	Texas panicum*	

* For partial control of this weed use a minimum of 1.33 pints per acre and apply pre-emergence

** For partial control of this weed use a minimum of 1.33 pints per acre and apply through a center-pivot irrigation system

*** Control of these weeds may be erratic due partially to variable weather conditions

Control may be improved by the following procedures

- 1 In corn use 2 to 2.5 pints per acre or the pre-plant surface-applied rates for Charger Max alone or in allowable tank mixtures when making pre-plant incorporated or pre-emergence applications
- 2 Thoroughly till moist soil to destroy germinating and emerged weeds. If Charger Max is to be applied pre-plant incorporated, this tillage may be used to incorporate Charger Max if uniform 2 inch incorporation is achieved as recommended under Application Procedures
- 3 Plant crop into moist soil immediately after tillage. If Charger Max is used pre-emergence, apply at planting or immediately after planting
- 4 If available, sprinkler irrigate within 2 days after application. Apply 1/2 to 1 inch of water. Use the lower water volume on coarse textured soils and the higher water volume on fine-textured soils. Also refer to the Center Pivot Irrigation Application section for this method of application
- 5 If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform shallow cultivation as soon as weeds emerge is recommended

ROTATIONAL CROPS

Do not rotate to food or feed crops other than those listed below

Charger Max applied alone

1 If crop treated with Charger Max alone is lost, any crop on this label may be planted immediately. Do not make a second broadcast application of Charger Max. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied.

2 Barley, oats, rye, or wheat may be planted 4 1/2 months following treatment. Alfalfa may be planted 4 months following application. Tomatoes may be planted 6 months after application.

3 Any crop on this label, in addition to root crops, tobacco, barley, buckwheat, milo, oats, rice, rye, wheat, cabbage, or peppers may be planted in the spring following treatment. Clover may be seeded 9 months following application. Do not graze or feed forage or fodder from cotton to livestock.

4 Following lay-by treatment or multiple treatments applied the previous season, any crop on this label, in addition to tobacco, cabbage, peppers, stone fruits, or tree nuts may be planted in the spring.

Charger Max in tank mixtures

For rotational crop restrictions, refer to the statements above for Charger Max applied alone and to the respective labels of the tank mix partner products for any additional restrictions.

Important Notes

To avoid injury to rotational alfalfa or clover, do not apply more than 1.9 lb active ingredient (2.0 pints Charger Max) pre-emergence, including preplant surface, preplant incorporated, postplant incorporated, etc. and do not make lay-by or other postemergence applications of Charger Max.

APPLICATION PROCEDURES

Application Timing

Charger Max alone or in some tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to the appropriate crop section of the label to determine if application timings listed below are recommended.

Preplant surface-applied

For minimum-tillage or no-tillage systems only. Charger Max alone or in some tank mixtures may be applied up to 45 days before planting certain crops. Use only split applications for treatments made 30 to 45 days before planting with 2/3 the recommended rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments made less than 30 days before planting may be made either as a split or single application. Refer to the directions for the individual crop to determine if early preplant surface application is recommended. If weeds are present at the time of treatment, apply in tank mixture with a contact herbicide such as Gramoxone MAX (paraquat) or Cornerstone (glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting or weed control will be decreased.

Preplant incorporated

Apply Charger Max to the soil and incorporate into the top 2 inches of soil within 14 days before planting using a finishing disk, harrows, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use preplant incorporation if furrow irrigation is used or when a period of dry weather after application is expected. If the crop will be planted in beds, apply and incorporate Charger Max after bed formation unless otherwise specified.

Pre-emergence

Apply Charger Max during planting (behind the planter) or after planting but before weeds emerge.

Post-emergence

Charger Max may be applied post-emergence to corn up to 40 inches tall. Charger Max will not provide post-emergence weed control, so it must be applied to weed-free soil or in combination with products that provide post-emergence control of weed species present at the time of application. Use only where allowed in crop-specific directions for use.

Special Application Procedures

California only, corn, safflower, pod crops

Preplant incorporated

Broadcast apply Charger Max to soil alone or in tank mixture with partner herbicides listed on this label and thoroughly incorporate with a disk or similar implement set to till 4 to 6 inches deep. For more thorough incorporation, cross-till the soil in 2 different directions. Crops may be planted in beds or on a flat surface. Caution should be used when forming beds such that only soil taken from the Charger Max treated area is used -

the untreated soil should not be brought to soil surface. If the application is made to pre-formed beds, incorporate Charger Max with a tillage implement set to till 2 to 4 inches deep. Care should be taken during tillage to keep the tilled Charger Max - treated soil on the beds.

Pre-emergence

Apply Charger Max after planting. Water with sprinkler or flood irrigation within 7 to 10 days.

Fall application only in IA, MN, ND, SD, WI, and portions of NE and IL

See specific instructions in the corn, soybeans, and pod crops sections of this label for timing of application and other information.

Do not apply to frozen ground. Use only on medium- and fine-textured soils with more than 2.5% organic matter that will be planted to corn or soybeans the following spring. Ground may be tilled either before or after application. Do not exceed 2 to 3 inch depth of incorporation if tilled after treatment. If a spring application is made, the total rate of spring plus fall applications must not exceed the maximum total rate for the specific crop or illegal residues might result.

Ground application

Apply Charger Max alone or in tank mixtures by ground application equipment in a minimum of 10 gallons per acre unless otherwise specified. Use a sprayer that provides accurate and uniform application. For tank mixtures with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after application is completed. The amount of herbicide required is calculated as follows:

<u>Band width in inches</u>		broadcast rate	=	amount needed
Row width in inches	X	per acre		per acre of field

For information on application in a lower volume, see "Low Volume Application". For application by air or by center-pivot systems, see "Aerial Drift Management" and "Aerial Drift Reduction Advisory Information" sections. For information on impregnating dry fertilizer, see "Dry Bulk Granular Fertilizer" section. For information on application via variable-rate equipment, see "Variable rate application" section.

SPRAY EQUIPMENT

Low Volume Application

Broadcast ground application

Use a sprayer such as AgChem RoGator®, Hagie, John Deere Hi-Cycle®, Melroe Spracoupe®, Tyler Patriot™, or Wilmar Air Ride® that provides accurate and uniform application. Use only water as a carrier. Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens be as fine as 100 mesh for some nozzles. Use a pump with capacity to maintain 35 to 40 PSI at the nozzles and provide sufficient agitation in the tank to keep the mixture in suspension. Use a minimum of 5 gallons of spray dilution per acre. Maximum speed recommended is 15 mph.

Rinse sprayer thoroughly after each use.

NOTE Low pressure nozzles are recommended to reduce drift and increase application accuracy Care should be taken when using automatic rate-controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected Nozzle screens should be used as recommended by the manufacturer All nozzles should be placed on 20-inch centers except flood nozzles which should be placed on 40-inch centers When flat-fan nozzles are used angles of 80° to 110° are recommended Always read and follow the manufacturer s directions for optimum set-up and performance of nozzles or tips

Aerial Application

Apply Charger Max in water alone or in tank mixtures with Atrazine linuron (e g Lorox®) or Sencor® in a minimum total volume of 2 gallons per acre by aircraft Charger Max may also be applied by air in mixture with Balan® pendimethalin (e g Prowl®) or trifluralin (e g Trust®) See specific tank mixture sections for use directions Avoid applications under conditions where uniform coverage cannot be obtained or where excessive drift might occur In order to assure that spray will be controllable within the target area when used according to label directions make applications at a maximum height of 10 ft above the crop using low-drift nozzles at a maximum pressure of 40 PSI and restrict application to periods when wind speed does not exceed 10 mph Do not apply in dead calm conditions which favor temperature inversion To ensure spray drift does not adversely affect adjacent sensitive non-target plants apply Charger Max alone or with Atrazine by aircraft at a minimum upwind distance of 400 ft from sensitive crops

Aerial Spray Drift Management

DRIFT MAY CAUSE DAMAGE TO ANY OTHER VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION APPROPRIATE BUFFER ZONES MUST BE MAINTAINED

Avoiding spray drift at the application site is the responsibility of the applicator The interaction of many equipment and weather-related factors determine the potential for spray drift The applicator and the grower are responsible for considering all these factors when making decisions

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops These requirements do not apply to forestry applications public health uses or to applications using dry formulations

- 1 The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor
- 2 Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees Where states have more stringent regulations they should be observed

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory

Aerial Drift Reduction Advisory

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion).

Controlling Droplet Size

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- **Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is released backwards, parallel to the airstreams, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length – For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application height – Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversion

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid application to humans or animals. Flaggers and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Center pivot Irrigation Application

Charger Max alone or in tank mixture with other herbicides specified on this label and which are registered for center pivot application may be applied in irrigation water pre-emergence (after planting but before emergence of crop or weeds) at rates recommended on this label. Charger Max may also be applied post-emergence to the crop but pre-emergence to weeds in crops where post-emergence application is allowed on this label. Follow all restrictions regarding height, timing, rate, etc. to avoid illegal residues. Do not apply through any type of irrigation system except center pivot. Crop injury, lack of effectiveness, or illegal residues in the crop may result from non-uniform distribution of treated water. Advice on calibration may be obtained from State Extension Specialists, equipment manufacturers, or other experts. Do not connect an irrigation system, including greenhouse systems, to a public water system unless the safety devices described on the label for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, must shut the system down and make adjustments should the need arise.

Operating instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.

- 2 The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump
- 3 The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is automatically or manually shut down
- 4 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- 5 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 6 Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump or piston pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- 7 Do not apply when wind speed favors drift beyond the area intended for treatment
- 8 Prepare a mixture with a minimum of 1 part herbicide to 1 part water and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension
- 9 Meter into irrigation water during the entire period of water application
- 10 Apply in 1/2 to 1 inch of water. Use the lower volume on coarse-textured soils and the higher volume on fine textured soils. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil

Precaution for center pivot application

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler patterns overlap excessively, crop injury may result.

Dry Bulk Granular Fertilizer

Many dry bulk granular fertilizers may be impregnated or coated with Charger Max alone or in tank mixtures which are registered for preplant incorporated or preplant surface applications to control weeds in crops on the Charger Max label and are not prohibited from use on dry bulk granular fertilizers.

When applying Charger Max alone or in tank mixture with dry granular fertilizers, follow all directions for use and precautions on the respective labels regarding target crops, rates, soil texture, application methods, timing, and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide / fertilizer mixture.

Prepare the herbicide/ fertilizer mixture by using any closed drum belt ribbon or other commonly used dry bulk fertilizer blender. Nozzles used to spray Charger Max or tank mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer and to avoid spraying the walls of the blender.

If the herbicide/ fertilizer mixture is too wet, add a highly absorptive material such as Agsorb® or Celatom MP-79® or similar clay or diatomaceous earth material to obtain a dry free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% by weight of absorptive material.

Calculate amounts of Charger Max, Atrazine, Atrazine plus Simazine, Simazine, Balance®, Sencor, or ethalfluralin (e.g. Sonalan®) by the following formulae:

2000 lb fertilizer/ ac X Pints/ acre liquid or flowable = pints liquid or flowable per ton fertilizer
or
2000 lb fertilizer/ ac X lb/ acre dry product = lb dry product per ton fertilizer

Pneumatic (compressed air) application (Charger Max alone)

High humidity, high urea concentration, low fertilizer use rates, or dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, pre-mix Charger Max with Aromatic® 200 at a rate of 1 to 4 pints per gallon of Charger Max. Aromatic 200 is a non-combustible/ non-flammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

Notes: Mixtures of Charger Max and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water-based or liquid fertilizer solutions for spraying applications. When impregnating Charger Max in a blender before application, a drier mixture can be obtained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size is recommended. Drying agents are not recommended for use with on-the-go impregnation equipment.

Precautions: To avoid potential for explosion, do not impregnate Charger Max alone or in mixture on ammonium nitrate, potassium nitrate, or sodium nitrate alone or in blends with other fertilizers. Do not use Charger Max alone or in mixture on straight limestone since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated. To avoid crop injury, do not apply on crops where bedding occurs.

Application: Apply 200 to 700 lb of the herbicide/ fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application is essential to prevent crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On fine- or medium-textured soils, in areas where soil incorporation is not planned, i.e. reduced tillage situations or some conventional tillage.

situations make applications approximately 30 days before planting to allow moisture to move the herbicide/ fertilizer mix into the soil On coarse-textured soils make applications approximately 14 days prior to planting

Variable-Rate Application

Variable-rate broadcast ground application in corn and soybeans Variable-rate technology may be used for more precise applications Two factors apply to variable rate applications including A) soil texture itself and B) % organic matter (%OM) and soil texture together Use the formulae below to determine a more precise rate for any given area within a field Do not use on soils with greater than 8% OM

Soil texture	Pints per acre			
	Preplant surface applied Preplant incorporated Pre-emergence		Early preplant	
	Factors		Factors	
	A	B	A	B
coarse	1 0	+ (0 1 x %OM)	1 2	+ (0 1 x %OM)
medium	1 17	+ (0 1 x %OM)	1 4	+ (0 08 x %OM)
fine	1 13	+ (0 1 x %OM)	1 67	+ (0 067 x %OM)

Formulae for fall applications Charger Max may be used only where corn or soybeans will be grown the following year in IA MN SD ND and WI and in portions of NE and IL Refer to specific instructions and restrictions regarding fall application in the Corn and Soybeans sections of this label Do not use on soils with more than 8% OM

Soil texture	Fall Applications	
	Pints per acre	
	Factors	
	A	B
coarse	Do not use	
medium	1 4	+ (0 113 x %OM)
fine	1 8	+ (0 067 x %OM)

Mixing Procedures

Charger Max alone Fill the spray tank 1/2 to 3/4 full with water or fluid fertilizer add the proper amount of Charger Max then add the rest of the water or fluid fertilizer Provide sufficient agitation during mixing and application to maintain a uniform emulsion

Tank Mixtures Fill the spray tank 1/4 full with water (or fluid fertilizer where allowed in the table below) and start agitation Add tank mix partner as listed below and allow it to become dispersed Products listed in the table below as Group 1 should be added before those listed as Group 2' Finally fill the rest of the tank with water or fluid fertilizer A compatibility test as described below is recommended before tank mixing

Group 1 (add before adding Group 2 products)	Group 2
2 4-D **Atrazine *Balance Pro benfluralin (e g Balan) bentazon (e g Basagran®) *Canopy® *Command® *Eptam® *ethalfuralin (e g Sonalan) *Fluometuron Liberty® Liberty ATZ *linuron Sencor *pendimethalin *Prometryn *Pursuit® *Scepter® *Simazine *Trifluralin	Cornerstone (glyphosate) glyphosate + 2 4-D paraquat (e g Gramoxone® MAX)
NOTES See further mixing instructions for tank mixtures with Atrazine and Fluometuron, or pendimethalin and Simazine * may be dispersed in fluid fertilizer ** may be dispersed in fluid fertilizer except for postemergence application	

COMPATIBILITY TEST

A jar test is recommended before tank mixing to ensure compatibility of Charger Max with other pesticides and fertilizers. The following directions assume a spray volume of 25 gallons per acre. If applying in other volumes, increase or reduce the amount of each product added proportionally.

Nitrogen solutions or complete fluid fertilizers may be used to replace some or all of the water in the spray mixture. Not all fertilizers are the same regardless of the N-P-K analyses. A compatibility test is necessary before dilution Charger Max or tank mixing in fluid fertilizers.

TEST PROCEDURE

- 1 Use 2 clear glass containers that hold 1 quart each and have tight-sealing lids. Add 1 pint of carrier (water or fluid fertilizer) to each jar. Use the same water source to be used for spraying and at the same temperature as expected during actual tank mixing and spraying.
- 2 Add 1/4 teaspoon (1.2 ml) of a compatibility agent such as COMPLETE COMPATIBILITY® to one of the jars and shake or stir to mix well.
- 3 Add the proportionate amount of each pesticide to be mixed according to label rates to each jar. For example, if a pesticide product is used at a rate of 1 quart

per acre in 25 gallons per acre add one teaspoon of that product to each jar
Add dry pesticides first then flowable concentrates then emulsifiable
concentrates and aqueous solutions (e.g. paraquat) last

- 4 After all ingredients have been added close and tighten the lids and mix well by inverting the jars 10 times each Let the jars stand for 20 to 30 minutes then examine for signs of incompatibility such as lumps flakes gelling or oily film on the jar If the mixture containing the compatibility agent shows improved mixing compared to jar without this should be added to the spray tank at the rate indicated on the label of the compatibility agent If the mixture separates but be remixed by several more inversions of the jar the mixture can be used in the spray tank so long as adequate agitation is maintained If the mixture is still incompatible with the compatibility agent repeat the test using the following methods to improve mixing a) slurry the dry pesticide in water before adding to the jar or b) add half the compatibility agent to the fertilizer or water and the other half directly to the emulsifiable concentrate or flowable pesticide before adding to the jar If the mixture is still incompatible do not use that combination of products

NOTE TESTED MATERIAL NOT USED IN THE ACTUAL APPLICATION MUST BE DISPOSED OF IN ACCORDANCE WITH THE STORAGE AND DISPOSAL INSTRUCTIONS ON THIS LABEL

CROP USE DIRECTIONS

CORN (ALL TYPES)

NOTE For all applications to corn do not graze or feed forage from treated areas for 30 days following application and do not harvest sweet corn ears from treated acres for 30 days following application or illegal residues might result

Charger Max alone, preplant surface, fall application

Restrictions Apply after Sept 30 in ND SD MN WI and north of Route 30 in IA Apply after Oct 15 north of Route 91 in NE and south of Route 30 in IA Apply after Oct 31 north of Route 136 in IL

In all locations apply to crop stubble after harvest when the sustained soil temperature is less than 55° F and falling In minimum tillage and no-till systems on soils having more than 2.5% organic matter use 1.67 to 2 pints per acre on medium-textured soils and 2 pints per acre on fine-textured soils Do not apply on frozen ground Tillage may precede the application A fall and spring tillage may follow the application but do not exceed an incorporation depth of more than 2 to 3 inches Minimize ridge and furrow formation in the tillage operations

NOTE If a spring application is made the total rate of the fall + spring application must not exceed the maximum total rate for corn or illegal residues might result

On medium- and fine-textured soils with minimum tillage or no-till systems in CO IA IL IN KS KY MN MO MT ND SD TN WI and WY apply 2/3 the recommended rate as a split treatment 30 to 45 days before planting and the remainder at planting

Applications made less than 30 days before planting may be either split or single application. On coarse soils, apply 1.33 pints per acre no more than 2 weeks before planting.

On medium- and fine-textured soils with minimum-tillage or no-till systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be made as above. If the amount of rainfall results in an unsatisfactory duration of weed control following the earlier treatment, a postemergent application of a herbicide registered for that use may be used, e.g., Atrazine, Beacon®, Bicep®, MAGNUM, Bicep II, MAGNUM, Exceed®, Accent®, bentazon, Moxy® (bromoxynil), or 2,4-D. If the post-emergence treatment includes the herbicide used preplant surface applied, do not exceed the total label rate for corn on that soil texture. Observe all directions for use, precautions, and limitations on the label of the post-emergence herbicide.

Preplant incorporated or pre-emergence

Follow instructions for use of Charger Max alone under **APPLICATION PROCEDURES**. On coarse soils, apply 1 to 1.33 pints per acre if organic matter content is less than 3% or 1.33 pints per acre if organic matter content is 3% or more. On medium soils, 1.33 to 1.67 pints per acre. On fine soils, apply 1.33 to 1.67 pints per acre if organic matter content is less than 3% or 1.67 to 2 pints per acre if organic matter is 3% or more.

Postemergence or lay by

To extend the duration of weed control in corn, a maximum rate of 2 pints per acre of Charger Max may be applied after corn emergence until the corn plants reach 40 inches tall, following any preplant surface applied, preplant incorporated, or pre-emergence herbicide application including Charger Max. For best results, apply to soil free of emerged weeds and directed toward the base of any corn plants more than 5 inches tall. The total rate of Charger Max applied on corn in any one crop year must not exceed 3.9 pints per acre, depending on the soil texture.

Control of problem weeds

For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 2 to 2.5 pints per acre as a single application or 1 to 1.33 pints per acre preplant incorporated followed by 1 to 1.33 pints per acre pre-emergence. However, do not apply more than a total of 2.5 pints per acre including both applications. Make the pre-emergence application during or after planting but before weeds or corn emerge. Apply the 1.33 pints per acre rate when a heavy infestation is expected. A shallow cultivation may be needed to control late-emerging weeds.

For control of woolly cupgrass and wild proso millet, use the following 3-step program: 1) Apply Charger Max early preplant, preplant incorporated, or pre-emergence at a rate of 1.67 pints per acre on medium soils or 2 pints per acre on fine soils, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 to 7 days. 2) Apply a postemergence tank mixture of Beacon at a rate of 0.38 oz per acre or Exceed at a rate of 1 packet per 4 acres, plus Accent at a rate of 0.33 oz per acre, plus 1 quart of crop oil concentrate, plus 1 gallon per acre of 28% nitrogen or the equivalent amount of ammonium sulfate when grasses are 2 to 3 inches tall and corn is at least 4 inches tall. 3) Cultivate 14 to 21 days after the postemergence application.

NOTES

- Do not apply more than the labeled rate for a given soil texture per year as either a single or split application or illegal residues might result
- In corn Charger Max may be used up to 2.5 pints per acre as preplant surface preplant incorporated or pre-emergence treatment on soils having an organic matter content from 6% to 20% or up to 2 pints per acre on any soil for extended residual control and where severe stands of problem weeds are expected
- In the event of escape of annual weeds following preplant surface preplant incorporated or pre-emergence treatment with Charger Max follow with application of an appropriately registered postemergence herbicide such as Atrazine Beacon Bicep MAGNUM Bicep II MAGNUM Exceed Accent bentazon (e.g. Basagran) Moxy (bromoxynil) or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment do not exceed the total annual labeled rate for corn on that soil texture
- Moxy (bromoxynil) may be applied postemergence alone or in combination with Atrazine. Do not exceed 1.2 lb ai of Atrazine in tank mix combination with Moxy. Refer to the labels of the tank mix partners for specific rates and precautions
- Do not use Charger Max on peat or muck soils

Tank mixtures for corn

Charger Max may be applied in tank mixtures with other herbicides diluted in water or fluid fertilizer before corn emerges. After corn emerges use only water as a carrier.

FOR TANK MIXTURES WITH ATRAZINE ALL RESTRICTIONS AND RATE LIMITATIONS ON THE ATRAZINE LABEL MUST BE FOLLOWED IF MORE RESTRICTIVE THAN ON THIS LABEL. IN ADDITION IF ATRAZINE MUST BE APPLIED AT A RATE LOWER THAN THOSE RECOMMENDED ON THIS LABEL BROADLEAF WEED CONTROL MAY BE REDUCED. REFER TO THE ATRAZINE LABEL FOR WEEDS CONTROLLED AT THE REDUCED RATES.

**TABLE 1
Additional weeds controlled in tank mixtures and special instructions**

Mixture (Charger Max plus)	Atrazine or Simazine (preplant surface PPI pre emergence)	Atrazine (post)	Atrazine + linuron	Atrazine or Simazine + pendimethalin (e.g Prowl)	Balance	Atrazine + Balance
Comments	2 3 4 5 7 8	2 3 4 5	2 3 4 5 6	1 2 3 4 5	2 3 7	2 3 5 7
browntop	X		X	X		X
panicum						
cocklebur	X	O	X	X		O X
common purslane	X		X	X	X	X
hairy nightshade	X		X	X		X
jimsonweed		X			X	X
kochia		X			X	X
lambquarters	X		X	X	X	X
morningglory	X	O	X	X	X	X
mustard		X			X	X
pigweed			X	X	X	X
prickly sida		X				
ragweed	X	X	X	X	X	X
smartweed	X	X	X	X	X	X
velvetleaf	X	X	X	X	O X	O X

X = control O = partial control

Comments

- 1 Fill the spray tank 1/4 full with water or fluid fertilizer start agitation add a compatibility agent such as Complete Compatibility according to its label directions then add Atrazine or Simazine and allow it to disperse Then add Charger Max and Prowl 4E Finally add the remaining water or fluid fertilizer
- 2 Atrazine or Simazine 4L or 90DF may be used The active ingredient in 1 lb of 90DF is equivalent to that in 1 8 pints of 4L
- 3 If using other brands or formulations of atrazine follow directions on that label
- 4 Observe all additional mixing directions on the Atrazine label
- 5 Do not exceed 2 5 lb ai of atrazine per acre per year Certain states may have rate limitations for atrazine within specific geographic areas Consult your state pesticide control agency for additional information It is a violation of this label to deviate from state use restrictions
- 6 Observe the rate recommendations on the brand of linuron being used
- 7 In minimum tillage and no till situations mix with paraquat (e.g Gramoxone) for control of most emerged annual weeds and suppression of perennial weeds Mix with Landmaster BW for suppression of emerged field bindweed and control or suppression of annual weeds Mix with Cornerstone (glyphosate) for control of most emerged annual or perennial weeds

Tank mixtures with Atrazine, Simazine, or mixtures of atrazine and simazine

In addition to the weeds controlled by Charger Max alone tank mixtures with Atrazine and/ or Simazine also control the following weeds

browntop panicum	cocklebur	common purslane
hairy nightshade	lambsquarters	morningglory
ragweed	smartweed	velvetleaf

Preplant surface applied - for use in minimum tillage or no till systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY

Follow instructions for use of Charger Max alone under Application Procedures and under application instructions for Charger Max in corn On medium soils use 1.67 pints/ acre Charger Max + 3.2 to 4 pints per acre Atrazine 4L or Simazine 4L or combination of the two On fine soils use 1.67 to 2 pints per acre Charger Max + 4 pints per acre Atrazine 4L or Simazine 4L or combination On coarse soils apply 1.33 pints per acre Charger Max + 3.2 pints Atrazine 4L or Simazine 4L or combination of the 2 Apply as a split or single treatment

Preplant incorporated or pre-emergence

Follow instructions for use of Charger Max alone under Application Procedures Apply mixtures with Atrazine 4L or Simazine 4L as indicated in TABLE 2

DO NOT apply more than the label rate for a given soil texture per year or illegal residues might result

Partial control of shattercane, wild proso millet, wooly cupgrass, and eclipta

For more consistent partial control of shattercane wild proso millet wooly cupgrass and eclipta where Charger Max is used alone or in tank mixtures or sequentially with other registered herbicides for use in corn apply at a rate of 2 to 2.33 pints per acre as a single application or as follows

- 1 Apply 1 to 1.33 pints per acre Charger Max + 2 lb ai per acre atrazine or simazine preplant incorporated followed by 1 to 1.33 pints per acre of Charger Max pre-emergence Make the pre-emergence application during or after planting but before weeds and corn emerge
- 2 Apply 1.33 pints per acre Charger Max alone or in tank mixture with up to 2 lb ai/ acre of atrazine or simazine preplant incorporated Do not exceed the total rate of triazine herbicide recommended for corn on the specific soil texture Follow with a post-directed application of ametryn (e.g. Evik®) at 2 lb ai per acre (2.5 lb 80DF or 1 gal 4L) Refer to the specific ametryn product label for further directions
- 3 Apply Eradicane® or Sutan® or equivalent (EPTC or butylate products) preplant incorporated at labeled rates followed by a pre-emergence application of Charger Max at 1 to 1.33 pints per acre Make the pre-emergence application during or after planting and before weeds and corn emerge Do not use in soils where rapid degradation of butylate or EPTC has been shown to occur

Precautions When following the application regimes above a shallow cultivation may be needed after the pre emergence or postemergence application to help control any late emerging shattercane or wild proso millet

TABLE 2 Charger max plus Atrazine, Simazine or Atrazine + Simazine Broadcast Rates per Acre			
Soil Texture	less than 3% organic matter		more than 3% organic matter
	with Atrazine 90DF ¹ or with Simazine 90DF ¹	with Atrazine 90DF ² and Simazine 90DF ²	with Atrazine 90DF ¹ or with Simazine 90DF ¹
coarse	0 8 to 1 pint Charger Max + + 1 2 to 2 2 lb 90DF	0 8 to 1 pint Charger Max + 0 6 to 1 1 lb Atrazine 90DF + 0 6 to 1 1 lb Simazine 90DF	1 pint Charger Max + 0 7 to 1 1 lb Atrazine 90DF + 0 7 to 1 1 lb Simazine 90DF
medium	1 to 1 33 pints Charger Max + 1 3 to 2 2 lb 90DF	1 to 1 33 pints Charger Max + 0 7 to 1 1 lb Atrazine 90DF + 0 7 to 1 1 lb Simazine 90DF	1 33 pint Charger Max + 0 9 to 1 1 lb Atrazine 90DF + 0 9 to 1 1 lb Simazine 90DF
fine	1 33 pints Charger Max + 1 8 to 2 2 lb 90DF	1 33 pints Charger Max + 0 9 to 1 1 lb Atrazine 90DF + 0 9 to 1 1 lb Simazine 90DF	1 33 to 1 67 pint Charger Max + 0 9 to 1 1 lb Atrazine 90DF ³ + 0 9 to 1 1 lb Simazine 90DF ³
muck/ peat (> 20% organic matter)	DO NOT USE		

NOTES 1) Use Simazine in preference to Atrazine when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter Charger Max may be used up to 2 33 pints per acre in tank mix combination with 2 2 lb/ acre of Atrazine 90DF or equivalent rate of Atrazine 4L. Refer to the Atrazine 4L label for weeds controlled at this reduced rate.

2) When tank mixing with Atrazine + Simazine use equal rates of each as shown in TABLE 2 when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected a 1 2 ratio of Atrazine to Simazine instead (e.g. for 1 2 lb total Atrazine + Simazine use 0 4 lb Atrazine + 0 8 lb Simazine). If using Atrazine 4L and Simazine 4L see the conversion formula in TABLE 1 above.

3) For cocklebur yellow nutsedge and velvetleaf control on fine-textured soils with more than 3% organic matter apply 2 25 lb per acre Atrazine 90DF or equivalent rate of Atrazine 4L or the same amount of Atrazine + Simazine with 1 33 to 1 67 pints Charger Max per acre.

Tank mixture with Atrazine postemergence

DO NOT APPLY THIS TANK MIXTURE IN FLUID FERTILIZER OR SEVERE CROP INJURY MAY OCCUR

Weeds controlled

barnyardgrass (watergrass)	yellow foxtail	prickly sida
crabgrass	jimsonweed	purslane
crowfootgrass	kochia	ragweed
fall panicum	lambsquarters	smartweed
giant foxtail	mustard	velvetleaf
green foxtail	pigweed	

Weeds partially controlled

cocklebur	morningglory	yellow nutsedge
-----------	--------------	-----------------

Apply 1 pint per acre of Charger Max plus 1 3 lb/ acre Atrazine 90DF on coarse soils
 1 33 pints per acre Charger Max plus 1 8 lb per acre Atrazine 90DF on medium soils or
 1 33 to 1 67 pints per acre Charger Max plus 1 88 to 2 2 lb per acre Atrazine 90DF on
 fine soils For better control of cocklebur morningglory velvetleaf and yellow nutsedge
 in fine soils with more than 3% organic matter use the higher rate (2 2 lb/ acre) rate of
 Atrazine If using Atrazine 4L 1 8 pints is equivalent in active ingredient to 1 lb of 90DF
 Over-the top application may be made before grasses and broadleaf weeds pass the 2-
 leaf stage and before corn is more than 5 inches tall Application to weeds greater than
 2 inches in height may result in unsatisfactory control

Lay-by application may be made to corn plants not more than 12 inches tall Application
 to corn more than 5 inches tall should be directed toward the base of the corn plants
 Occasionally some leaf burn may result but this will not affect later growth or yield

Tank mixtures of Charger Max plus Atrazine may be applied following use of any
 registered pre-plant surface-applied preplant incorporated or pre-emergence corn
 herbicide (including Charger Max plus Atrazine) The total rate of metolachlor from
 Charger Max or other brands should not exceed 3 9 pints (3 72 lb ai) per acre and the
 total rate of atrazine from any brand may not exceed 2 5 lb ai per acre during one crop
 year or illegal residues can result Refer to the Atrazine label for further information on
 restrictions related to soil textures geographic areas and rotational crops

Tank mixture with Atrazine plus linuron for control of pigweed and lambsquarters

For prolonged control of lambsquarters and pigweed in DE MD NJ NY PA VA and
 WV Charger Max may be applied pre-emergence in a mixture with Atrazine and linuron
 (e g Lorox) Apply Charger Max and Atrazine according the rates in TABLE 2 plus
 linuron as follows

Soil Texture	Broadcast Rate per Acre*
sandy loam (1% to 3% organic matter)	0 67 lb linuron 50% DF
sandy loam (3% to 6% organic matter)	1 lb linuron 50% DF
medium and fine textured soils (1% to 6% organic matter)	1 lb linuron 50% DF
* If using linuron 4L one pint of 4L is equivalent to 1 lb 50% DF	

Observe all directions for use precautions and limitations on all the labels when applying these products in tank mixture

Tank mixture with Atrazine or Simazine plus pendimethalin for prolonged control of lambsquarters and pigweed in field corn only

For prolonged control of lambsquarters and pigweed in addition to a broad spectrum of annual broadleaf and grass weeds in northeastern U S including MI IN KY and states east of these Charger Max may be applied in a tank mixture with Atrazine or Simazine plus pendimethalin Apply after planting but before corn or weeds emerge by ground equipment in a minimum of 10 gallons of water or 20 gallons of liquid fertilizer per acre If applied by air use a minimum of 5 gallons of water per acre Do not use wettable powder formulations of atrazine See TABLE 2 for rates of Charger Max Atrazine and Simazine Add pendimethalin as follows

Broadcast Rate per Acre (4 lb per gallon formulation of pendimethalin)			
soil texture	% organic matter in soil		
	less than 1.5%	1.5 to 3%	over 3%
coarse	1.5 to 2 pints	2 pints	3 pints
medium	2 pints	3 pints	3 pints
fine	2 pints	3 pints	3 pints

Observe all directions for use precautions and limitations on all the labels when applying these products in tank mixture Refer to the pendimethalin label for replanting instructions in the event of crop loss

Tank mixture with Balance

Use in field corn only A tank mixture of Charger Max plus Balance controls all the weeds listed on this label plus improves control of problem weeds including velvetleaf tall or common waterhemp jimsonweed kochia common lambsquarters common ragweed and others Also this tank mixture will improve control of biotypes of these weeds which are resistant to ALS-inhibitor and triazine herbicides It will also contribute toward control of some problem grass weeds Application may be made preplanting (surface applied up to 14 days prior to planting) preplant incorporated or pre-emergence in conventional tillage conservation tillage or no-till systems Use the rate of Charger Max applied alone listed on this label for the type of application plus Balance as follows For early preplant applications 8 to 14 days before planting use 1.5 ounces per acre Balance on coarse soils or 1.5 to 2 ounces per acre Balance in medium or fine soils For application 0 to 7 days before planting or for preplant incorporated or pre-emergence application use 1 ounce per acre Balance in coarse soils or 1 to 1.5 ounces per acre Balance in medium or fine soils Where difficult to control weed species are present use the maximum allowable rate where ranges are listed Observe all directions and restrictions on the Balance label

Tank mixture with Balance plus Atrazine

The mixture of Charger Max plus Balance plus Atrazine controls all weeds listed on this label for Charger Max plus Atrazine and adds improved control of velvetleaf kochia

pigweeds tall or common waterhemp jimsonweed common lambsquarters common ragweed and biotypes of these weeds which are resistant to ALS-inhibitor and triazine herbicides Balance will also contribute toward the control of problem grass and other broadleaf weed species Application may be made preplanting (surface applied up to 14 days prior to planting) preplant incorporated or pre-emergence in conventional tillage conservation tillage or no-till systems Use the rate of Charger Max plus Atrazine listed on this label for the type of application plus Balance as follows For early preplant applications 8 to 14 days before planting use 1.5 ounces per acre Balance on coarse soils or 1.5 to 2 ounces per acre Balance in medium or fine soils For application 0 to 7 days before planting or for preplant incorporated or pre-emergence application use 1 ounce per acre Balance in coarse soils or 1 to 1.5 ounces per acre Balance in medium or fine soils Where difficult to control weed species are present use the maximum allowable rate where ranges are listed Observe all directions and restrictions on the Balance label

Tank Mixtures with Charger Max plus Atrazine, Simazine, or Balance, or Charger Max plus Atrazine plus Balance, or Charger Max plus Simazine plus Balance For Minimum Tillage Or No-Till Systems

Paraquat, Cornerstone (glyphosate), or Landmaster® BW

In minimum tillage or no-till systems where corn is planted directly into a cover crop stale seedbed established sod or previous crop residues paraquat (e.g. Gramoxone Max) glyphosate (e.g. Cornerstone) or Landmaster BW may be added to a mixture of Charger Max plus Atrazine Simazine or Balance or Charger Max plus Atrazine plus Balance or Charger Max plus Simazine plus Balance Mix with paraquat for control of most emerged annual weeds and suppression of perennial weeds Mix with Landmaster BW for suppression of emerged field bindweed and control or suppression of annual weeds Mix with Cornerstone (glyphosate) for control of most emerged annual or perennial weeds Apply before during or after planting but before corn emerges at the rates indicated below

Paraquat

<u>Weed height</u>		
1 to 3 inches	3 to 5 inches	5 to 6 inches
<u>Rate per acre, paraquat*(as paraquat ion)</u>		
56 to 75 lb ai	75 to 94 lb ai	94 to 112.5 lb ai

Add 1 to 2 pints of high-active-ingredient non-ionic surfactant such as Activate Plus per 100 gallons of spray mix Do not apply in suspension-type fluid fertilizers This mixture will not control weeds taller than 6 inches

* Check the label of the paraquat product being used to determine the amount of active ingredient (as paraquat ion) per gallon of solution

Landmaster BW

Apply at a rate of 1.7 to 3.4 pints per acre depending on weed species and size See the Landmaster BW label for specific instructions

Cornerstone (glyphosate)

See the Cornerstone label for specific directions. Apply in 20 to 60 gallons of water of fluid fertilizer per acre by ground equipment. Mix Cornerstone with Charger Max plus Atrazine, Simazine, or Atrazine plus Simazine as follows. Use Simazine in preference to Atrazine when heavy infestations of crabgrass or fall panicum are expected.

On coarse soils, apply 1.0 pints per acre of Charger Max plus 1.3 lb per acre Atrazine 90DF or Simazine 90DF, or with 0.7 lb per acre Atrazine 90DF plus 0.7 lb per acre Simazine 90DF.

On Medium soils, apply 1.33 pints per acre Charger Max plus 1.8 lb per acre Atrazine 90DF or Simazine 90DF, or with 0.9 lb per acre Atrazine 90DF plus 0.9 lb per acre Simazine 90DF.

On fine soils, apply 1.33 to 1.67 pints per acre Charger Max plus 1.8 to 2.2 lb per acre Atrazine 90DF or Simazine 90DF, or with 0.9 to 1.1 lb per acre of Atrazine 90DF plus 0.9 to 1.1 lb per acre Simazine 90DF.

Atrazine, Atrazine plus 2,4-D, Atrazine plus Balance

In minimum tillage or no-till systems where corn is planted directly into a cover crop stale seedbed, established sod, or previous crop residues, Charger Max in combination with Atrazine or Atrazine plus Balance will kill most emerged small annual weeds. Use Balance combinations only in field corn. Apply as described in TABLE 2. Where crop residues are heavy, add 0.8 to 1.6 pints per acre of 2,4-D Amine 4 (3.8 lb ai per gallon) to the spray tank last, and apply in a minimum of 25 gallons of spray mixture per acre.

Nitrogen solutions or complete liquid fertilizers applied before corn emergence enhance burndown of existing weeds, and are therefore recommended instead of water as carrier for this mixture. Add an organosilicone surfactant such as Silkin as its recommended rate, or Prime Oil plus 28% nitrogen liquid. Apply before weeds exceed 3 inches in height.

In fields with existing sod grasses such as bromegrass, orchardgrass, rye, or timothy when weeds exceed 3 inches in height or under very dry conditions, add paraquat at a rate of 94 lb ai per acre (equivalent to 2.5 pints Gramoxone MAX) in place of or in addition to 2,4-D as indicated above. Do not apply paraquat with suspension-type liquid fertilizers.

Marksman

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, Charger Max plus Marksman applied at rates on the Marksman label will kill most small emerged weeds. Apply Charger Max plus Marksman before, during, or after planting but before corn emerges in medium and fine soils with more than 2.5% organic matter. This mixture may also be used postemergence on corn that is less than 3 inches tall and before grasses exceed the 2-leaf stage. Nitrogen solutions and complete liquid fertilizers applied before corn emergence enhance burndown of existing weeds, and are recommended as carriers. In fields with existing vegetation

more than 3 inches tall or under very dry conditions add paraquat at labeled rates Do not apply paraquat in liquid fertilizers

Tank Mixtures for Postemergence Salvage Weed Control (Field Corn Only)

For postemergence control of weeds in specific types of field corn Charger Max may be tank mixed with the herbicides listed below Full season weed control from early preplant preplant incorporated or pre-emergence treatments provide maximum yield potential However if control of emerged weeds becomes required a postemergence program as described below may be used to provide residual control for the remainder of the season

NOTES Follow all label directions restrictions and precautions for each product used Apply only to varieties of field corn specified on the label of the tank mix product Do not use fluid fertilizers with these mixtures or corn injury may result In-row weed control may be reduced because of lack of coverage when applied to corn more than 4 inches tall

Liberty (in Liberty Link Corn or corn warranted by Bayer CropScience as tolerant to Liberty Herbicide)

Charger Max may be mixed with Liberty herbicide for postemergence application in corn grown from Liberty Link seed or corn warranted by Bayer CropScience as being tolerant to Liberty herbicide Liberty provides postemergence control of a broad spectrum of grass and broadleaf weeds while Charger Max provides residual control of grasses and some broadleaf weeds For tank mixing use the rate listed on the Liberty label for the species of weeds to control plus the minimum rate listed on this label for Charger Max applied alone preplant incorporated or pre-emergence If multiple weed species are present use the maximum rate on the Liberty label for the species and stages present

Liberty ATZ (in Liberty Link Corn or corn warranted by Bayer CropScience as tolerant to Liberty ATZ Herbicide)

Charger Max may be mixed with Liberty ATZ herbicide for postemergence application in corn grown from Liberty Link seed or corn warranted by Bayer CropScience as being tolerant to Liberty ATZ herbicide Liberty ATZ provides postemergence control of a broad spectrum of grass and broadleaf weeds while Charger Max provides residual control of grasses and some broadleaf weeds For tank mixing use the rate listed on the Liberty ATZ label for the species of weeds to control plus the minimum rate listed on this label for Charger Max applied alone preplant incorporated or pre-emergence If multiple weed species are present use the maximum rate on the Liberty ATZ label for the species and stages present Do not apply to corn more than 12 inches tall

Cornerstone/ Cornerstone Plus (glyphosate) in Roundup Ready Corn

Charger Max may be tank mixed with Cornerstone Cornerstone Plus or other glyphosate liquid herbicides for postemergence application in corn designated as Roundup Ready Application may be made from emergence of corn to when the corn reaches 30 inches tall or the V8 stage (8 leaves with collars) which ever comes first This mixture will provide control of all weeds listed on the glyphosate label plus the residual control of weeds listed on this label for Charger Max Use the minimum

Charger Max rate postemergence with Cornerstone or Cornerstone Plus as listed in the Corn - Charger Max alone instructions for preplant incorporated or pre-emergence application Refer to the Cornerstone/ Cornerstone Plus labeling for use directions precautions limitations requirement for additional surfactants etc for those products Apply 24 to 34 fl oz per acre of Cornerstone or Cornerstone Plus for control of labeled broadleaf and grass weeds Refer to additional directions on the Cornerstone/ Cornerstone Plus labeling regarding control of problem species

Cornerstone/ Cornerstone Plus (glyphosate) plus Atrazine in Roundup Ready Corn

Charger Max plus Atrazine may be tank mixed with Cornerstone Cornerstone Plus or other glyphosate liquid herbicides for postemergence application in corn designated as Roundup Ready Application may be made from emergence of corn to when the corn reaches 12 inches in height This mixture will provide control of all weeds listed on the glyphosate label plus the residual control of weeds listed on this label for Charger Max plus Atrazine Use the minimum Charger Max plus Atrazine rate postemergence with Cornerstone or Cornerstone Plus as listed in the Corn - Charger Max plus Atrazine or Simazine instructions above for preplant incorporated or pre-emergence application Refer to the Cornerstone/ Cornerstone Plus labeling for use directions precautions limitations requirement for additional surfactants etc for those products Apply 24 to 34 fl oz per acre of Cornerstone or Cornerstone Plus for control of labeled broadleaf and grass weeds Refer to additional directions on the Cornerstone/ Cornerstone Plus labeling regarding control of problem species

COTTON

Charger Max alone

Precautions To avoid crop injury do not apply Charger Max in sand or loamy sand soils nor in areas where water is likely to pond over the bed To avoid concentration in seed furrows do not make broadcast applications to cotton planted in furrows more than 2 inches deep Band applications may be made to cotton planted in furrows deeper than 2 inches but band width should not exceed the width of the bottom of the furrow DO NOT APPLY IN GAINES COUNTY TX DO NOT APPLY ON TALOKA SILT LOAM DO NOT GRAZE OR FEED FORAGE OR FODDER FROM COTTON TO LIVESTOCK

In AR KS LA MS TN and bootheel region of MO apply Charger Max pre-emergence at a rate of 0.5 to 1 pint per acre on sandy loams 0.66 to 1.33 pints per acre on medium soils or 1 to 1.33 pints per acre on fine soils

In NM OK and TX apply preplant incorporated or pre-emergence at a rate of 1 pint per acre on sandy loams 1 to 1.33 pints per acre on medium soils or 1.33 pints per acre on fine soils For preplant incorporated application apply to the soil and incorporate into the top 1 inch of soil immediately before planting At planting or after planting but before crop or weeds emerge Use a rolling cultivator or similar implement to uniformly incorporate not more than one inch deep Use preplant incorporated application when furrow irrigation will be used or when a period of dry weather after application is expected If furrow irrigation is used wet the top of the bed for best results If the crop is to be planted in beds apply and incorporate after beds have been formed Plant cotton below the zone of incorporation i.e. at least 1 inch deep in fine soils and 1.5

inches deep in medium and coarse soils. If incorporated before planting, use a planter that causes a minimum of soil disturbance.

For best control of yellow nutsedge and suppression of seedling Johnsongrass, apply preplant incorporated at the maximum rate for the soil texture. See also tank mixtures with Prometryne below.

Tank Mixtures for Cotton

Prometryne 4L

Charger Max may be tank mixed with Prometryne 4L using either water or fluid fertilizer as the carrier. If using fluid fertilizer, ensure only enough is mixed to be sprayed in one operation. Do not let mixtures with fluid fertilizers stand without agitation.

In addition to weeds controlled by Charger Max alone applied preplant incorporated or pre-emergence, the tank mixture will also control junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of cocklebur and coffeeweed.

Apply this mixture preplant incorporated or pre-emergence at the rates indicated below. Plant cotton below the zone of incorporation, i.e., at least 1 inch deep in fine soils and 1 1/2 inches deep in medium and coarse soils. If incorporated before planting, use a planter that minimizes soil disturbance. Refer to the Prometryne 4L label for all appropriate use directions, precautions, and restrictions.

Tank Mixtures with Prometryne, NM, OK, TX			
Use Area	Soil Texture	Broadcast Rate per Acre	
		Charger Max	Prometryne 4L
all	sand, loamy sand	DO NOT USE	
OK blacklands and gulf coast TX	loams	0.8 to 1.33 pints	2.4 pints
	clays	1.33 pints	4.8 pints
Rio Grande Valley TX	loams	0.8 to 1.33 pints	3.2 pints
	clays	1.33 pints	4.8 pints
NM high plains rolling plains Edwards Plateau southwest TX	sandy loams	0.8 to 1 pint	1.6 pints
	loams	0.8 to 1.33 pints	2.4 pints
	sandy clay loams	1.33 pints	2.4 pints
	other clay soil	1.33 pints	3.2 pints

Precautions: To avoid crop injury, do not apply Charger Max in sand or loamy sand soils nor in areas where water is likely to pond over the bed. To avoid concentration in seed furrows, do not make broadcast applications to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow. Do not apply in cut areas of newly leveled fields, nor in areas of excess salt. Do not use on glandless varieties of cotton.

DO NOT APPLY IN GAINES COUNTY TX. DO NOT APPLY ON TALOKA SILT LOAM. DO NOT GRAZE OR FEED FORAGE OR FODDER FROM COTTON TO LIVESTOCK.

Fluometuron

Charger Max may be tank mixed with appropriately registered herbicides containing fluometuron such as Fluometuron 4L or 80DF and applied pre-emergence for control of weeds on this label plus those on the fluometuron label including spotted spurge hyssop spurge nodding spurge and prostrate spurge Apply to the soil surface at planting or after planting but before weeds or crop emerge at the rates listed below To avoid any problems with incompatibility see the instructions above under Incompatibility Test and use a compatibility agent as required

Use Area	Soil Texture	Broadcast Rate per Acre	
		Charger Max	Fluometuron 4L*
all	sand loamy sand	DO NOT USE	
AR LA MS TN bootheel of MO	sandy loam	1/2 to 1 pint	2 pints
	loam silt loam silt	66 to 1 33 pints	2 to 3 25 pints
	fine soil	1 to 1 33 pints	3 25 to 4 pints
Rio Grande Valley eastern gulf coast TX eastern OK	sandy loam	0 8 to 1 pint	2 pints
	loam silt loam silt	1 to 1 33 pints	2 to 3 25 pints
	fine soil	1 33 pints	3 25 to 4 pints

* If using an 80 DF formulation of fluometuron active ingredient in 1 lb 80 DF = 1 7 pint 4L

Precautions To avoid crop injury do not apply Charger Max in sand or loamy sand soils nor in areas where water is likely to pond over the bed To avoid concentration in seed furrows do not make broadcast applications to cotton planted in furrows more than 2 inches deep Band applications may be made to cotton planted in furrows deeper than 2 inches but band width should not exceed the width of the bottom of the furrow The use of Fluometuron following use of a systemic insecticide at planting may result in crop injury
DO NOT APPLY IN GAINES COUNTY TX DO NOT APPLY ON TALOKA SILT LOAM DO NOT GRAZE OR FEED FORAGE OR FODDER FROM COTTON TO LIVESTOCK

Paraquat or Cornerstone (glyphosate) and/ or Fluometuron

In minimum tillage or no till systems where cotton is planted directly into a cover crop stale seedbed or previous crop residues Charger Max alone or in mixture with fluometuron may be tank mixed with paraquat (such as Gramoxone) or Cornerstone (glyphosate) herbicides Additional emerged weeds controlled will be those described on the glyphosate or paraquat label Observe all directions precautions and restrictions on the label of the tank mix products

Apply after emergence of weeds before during or after planting but before emergence of cotton Apply Charger Max at 0 8 to 1 pint per acre on sandy loam or medium to fine soils For fluometuron mixtures see the table above Add paraquat or Cornerstone as indicated below

Paraquat

Rate per acre paraquat (as ionized active ingredient*) * e.g Gramoxone MAX contains 3 lb ai per gallon	Weed Height		
	1 to 3 inches	3 to 5 inches	5 to 6 inches
	0 56 to 0 75 lb ai	0 75 to 0 94 lb ai	0 94 to 1 13 lb ai
	Add 1 to 2 pints of high active-ingredient non-ionic surfactant such as Activate Plus per 100 gallons of spray mix Do not apply in suspension type fluid fertilizers This mixture will not control weeds taller than 6 inches		

Cornerstone (glyphosate)

Refer to the Cornerstone (or other glyphosate) label for rates and weeds controlled as well as all other directions precautions and restrictions Apply with ground equipment in 20 to 60 gallons per acre of spray mixture diluted with water or liquid fertilizer

Precautions If a heavy rain occurs soon after application crop injury may result especially in poorly drained areas where water stands for several days or where the seeding slit has not fully closed
DO NOT USE IN GAINES COUNTY TX

PEANUTS

Charger Max applied alone

Do not graze peanut forage or fodder to livestock for 30 days after application Do not apply within 90 days of harvest or illegal residues might result

Application may be made after any of the following preplant incorporated herbicides when used according to their label directions trifluralin (e.g. Trust) benfluralin (e.g. Balan) ethalfluralin (e.g. Sonalan) or pendimethalin (e.g. Prowl)

Apply Charger Max preplant incorporated postplant incorporated or pre-emergence at a rate of 1 to 1.33 pints per acre in the southeast for most weeds or 0.8 to 1.33 pints per acre in NM OK and TX For partial control of beggarweed in the southeast apply at a rate of 1.33 to 2 pints per acre For preplant incorporation see the instructions above under Application Procedures For postplant incorporation apply to the soil and shallowly incorporate after planting but before peanut germination Keep incorporation above depth of seeds or seed injury might result

Where weed pressure is heavy or where difficult-to-control species are expected Charger Max is most effective when used as follows

IN THE STATES OF AL FL GA NC SC VA Apply Charger Max preplant incorporated alone as directed above Apply Charger Max preplant incorporated alone as directed above or with benfluralin as described below or pre-emergence before ground cracking alone at a rate of 1 to 2 pints per acre **Do not use Charger Max, Dual® II Magnum, or equivalent products after peanut emergence** If peanuts have emerged use Dual Magnum (NOT Dual II Magnum or Charger Max) according to its directions Follow with a lay-by application of Dual Magnum (NOT Dual II Magnum or Charger Max) according to its label directions

Do not apply more than a total of 2.67 lb of active ingredient (s-metolachlor) in this product plus Dual Magnum used sequentially per acre in any one crop year or illegal residues might occur

IN THE STATES OF NM OK and TX Apply Charger Max preplant incorporated alone as directed above or with benfluralin as described below **Do not use Charger Max, Dual II Magnum, or equivalent products after peanut emergence** If peanuts have emerged use Dual Magnum (NOT Dual II Magnum or Charger Max) according to its directions Follow with a lay-by application of Dual Magnum (NOT Dual II Magnum or Charger Max) according to its label directions

Do not apply more than a total of 2.67 lb of active ingredient (s-metolachlor) in this product plus Dual Magnum used sequentially per acre in any one crop year or illegal residues might occur

Tank mixture with benfluralin

A tank mixture with benfluralin (e.g. Balan) will control all weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those listed on the benfluralin label. Apply 1 to 1.33 pints per acre Charger Max plus 1 to 1.25 lb ai of benfluralin in a minimum of 10 gallons of spray mixture per acre by ground or 5 gallons per acre by air. Follow the recommended procedures for soil preparation and incorporation on the benfluralin label. Apply and incorporate up to 14 days before planting.

Tank mixture with Pursuit

A tank mixture with Pursuit will control all weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those listed on the Pursuit label. Refer to both labels for rates, timing, precautions, and restrictions. Charger Max will not control emerged weeds.

Tank mixture with ethalfluralin

A tank mixture with ethalfluralin (e.g. Sonalan) will control all weeds in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those listed on the ethalfluralin label. Refer to both labels for rates, timing, precautions, and restrictions.

Apply the mixture preplant incorporated at the rate specified in the following table. Follow directions for incorporation on the ethalfluralin label.

Soil Texture	Broadcast rates per acre			
	Southeast		NM, OK, TX	
	Charger Max	ethalfluralin*	Charger Max	ethalfluralin*
coarse	1 to 1.33 pints	0.5 to 7.5 lb ai	0.8 to 1.33 pints	0.5 to 7.5 lb ai
medium	1 to 1.33 pints	0.66 to 1 lb ai	0.8 to 1.33 pints	0.66 to 1 lb ai
fine	1 to 1.33 pints	0.84 to 1.13 lb ai	0.8 to 1.33 pints	0.84 to 1.13 lb ai

* See the specific ethalfluralin label for active ingredient content per pint of liquid or per pound of dry flowable.

Tank mixture with pendimethalin

A tank mixture with pendimethalin applied preplant incorporated will control all weeds controlled by Charger Max alone plus Texas panicum, field sandbur, Johnsongrass, from seed, lambsquarters, kochia, annual spurge, and other species on the pendimethalin product label. Apply at the rates indicated in the table below by ground or aerial equipment within 14 days before planting. Incorporate into the top 1 to 2 inches of soil before planting and within 7 days of application using a finishing disk or other similar implement capable of providing uniform incorporation. If peanuts will be planted in beds, apply and incorporate after bed formation. Follow all directions, limitations, precautions, and application information for peanuts on the pendimethalin label.

Soil texture	Broadcast rate per acre*	
	NM, OK, TX	Other states
	Charger Max + pendimethalin	Charger Max + pendimethalin
sand loamy sand	0.8 pint + 1 to 1.5 pints	1 to 1.33 pints + 1.5 to 2 pints
sandy loam	0.8 to 1 pint + 1 to 1.5 pints	1 to 1.33 pints + 1.5 to 2 pints
fine soil	1.33 pints + 1 to 1.5 pints	1.33 pints + 1.5 to 2 pints

* Assumes a 3.3 lb/gallon liquid formulation. If using other formulations, use equivalent amount of active ingredient.

Sequential application of Storm®

Apply Charger Max alone as directed on this label and follow with a postemergence treatment of Storm as directed on the Storm label

Multiple Applications

Where weed pressure is heavy or where species difficult to control are expected Charger Max is most effective when used as follows

SOUTHEAST ONLY ((AL FL GA NC SC VA)

Preplant Incorporated

Apply preplant incorporated as directed above for peanuts alone or in mixture with benfluralin

OR

Apply any time from pre-emergence to before ground cracking at a rate of 1 to 2 pints per acre for extended control of weeds not yet emerged **Do not use Charger Max after peanut emergence** If peanuts have emerged use Dual Magnum according to directions on that label Follow the PPI or pre-emergence application with a lay-by application of Dual Magnum (NOT DUAL II MAGNUM) according to instructions on that label

NOTE Do not apply more than the equivalent of 2.67 lb per acre of active ingredient of Charger Max (i.e. s-metolachlor) during any one year or illegal residues might result If Dual Magnum is used as a sequential treatment the lb active ingredient of Charger Max plus the lb active ingredient of Dual Magnum may not exceed 2.67 lb/ acre (each contains 0.95 lb ai per pint)

Do not graze or feed peanut forage or fodder to livestock for 30 days following application and do not apply within 90 days before harvest or illegal residues might result

SOUTHWEST ONLY (NM OK TX)

First Application Apply Charger Max preplant incorporated or pre-emergence to before ground cracking as directed above for peanuts either alone or with benfluralin **Do not use Charger Max after peanut emergence** If peanuts have emerged use Dual Magnum according to its label

Second application Apply Dual Magnum (not Dual II Magnum) at lay-by as directed for peanuts alone on the Dual Magnum label Use only when late-germinating weeds are expected to be a problem

NOTE Do not apply more than the equivalent of 2.67 lb per acre of active ingredient of Charger Max (i.e. s-metolachlor) during any one year or illegal residues might result If Dual Magnum is used as a sequential treatment the lb active ingredient of Charger Max plus the lb active ingredient of Dual Magnum may not exceed 2.67 lb/ acre (each contains 0.95 lb ai per pint)

Do not graze or feed peanut forage or fodder to livestock for 30 days following application and do not apply within 90 days before harvest or illegal residues might result

POD CROPS

Charger Max may be applied to pod crops including garbanzo beans great northern beans kidney beans lima beans mung beans navy beans English peas* southern

peas (such as blackeye pinkey Crowder and others) pinto beans snap (green wax string) beans lupines (sweet white white sweet and grain)

* Use only pre-emergence on English peas Do not use on English peas in north-eastern states or crop injury may occur

Charger Max alone

FALL APPLICATION In ND SD MN WI and north of Route 30 in IA apply after Sept 30 In NE and south of Route 30 in IA apply after Oct 15 In IL north of Route 136 apply after Oct 31

Apply to crop stubble after harvest when the sustained soil temperature at a 4 inch depth is less than 55 degrees F and falling In minimum or no-till systems in soil having more than 2.5% organic matter use 1.67 to 2 pints per acre on medium textured soils and 2 pints per acre on fine textured soils Do not apply to frozen ground A tillage operation may precede the application A fall or spring tillage operation may follow the application but do not exceed a depth of incorporation of 2 to 3 inches Minimize ridge and furrow formation during tillage

NOTE If a spring application is also made the total rate applied in fall plus spring must not exceed a maximum total rate of 2 pints per acre

SPRING APPLICATION Apply preplant incorporated or pre-emergence at a rate as follows for the appropriate soil type On coarse soils with less than 3% organic matter apply 1 to 1.3 pints per acre or 1.33 pints per acre if organic matter is 3% or more On medium soils apply 1.33 to 1.67 pints per acre On fine soils apply 1.33 to 1.67 pints per acre if the organic matter content is less than 3% or 1.67 to 2 pints per acre if the organic matter content is 3% or more

Tank mixtures for pod crops

When using tank mixtures with Charger Max on pod crops do not cut for hay within 120 days after application or illegal residues might result Do not use forage within 60 days after an application of Charger Max Do not apply more than 2 pints per acre of Charger Max within any one year

Tank mix or sequential application with Eptam® - green or dry beans

This mixture controls all weeds controlled by Charger Max alone plus those controlled by Eptam alone For sequential application with Eptam apply Eptam alone preplant incorporated as specified on the Eptam label Observe all limitations on the Eptam label for rates geographic areas and species or varietal restrictions Do not exceed 3.5 pints per acre of Eptam 7E on small white beans or green beans grown in coarse textured soils Follow with application of Charger Max alone at or after planting but before weeds or crop emerge On coarse soils apply 0.8 pints per acre Charger Max if organic matter content is less than 3% or 1 pint per acre if organic matter is 3% or more On medium soils apply 1 pint per acre of Charger Max if the organic matter is less than 3% or 1.33 pints per acre if organic matter is 3% or more On fine soils apply 1.33 pints per acre of Charger Max if organic matter is less than 3% or 1.33 to 1.67 pints per acre if organic matter is 3% or more

Tank mixture with Trust (trifluralin)

Charger Max may be tank mixed with Trust (trifluralin) for preplant incorporation application to dry beans (kidney navy pinto etc) Lima beans and snap beans Weeds controlled include those on the Charger Max label as well as those on the trifluralin label Apply using either ground or aerial equipment and incorporate up to 14 days before planting Use incorporation equipment that will provide uniform incorporation to a depth of 2 inches Follow rate directions for each product alone for the appropriate soil texture Follow all restrictions and precautions on both labels

POTATOES

Charger Max alone

Apply Charger Max pre-emergence or preplant incorporated for control of weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label Within the rate ranges below use the higher rates on fine textured soils or soils high in organic matter and the lower rates on coarse textured soils or low in organic matter Effectiveness will be reduced if later cultural practices expose untreated soil For application via center pivot see the **Center pivot irrigation application** section of the label

For preplant or postplant incorporated application apply Charger Max at a rate of 1 to 2 pints per acre and incorporate into the top 3 inches using a finishing disk harrows rolling cultivator or similar implement Planting and later cultural practices should not bring untreated soil to the surface Postplant incorporated application may be made any time after planting but before emergence of potatoes Use an implement that evenly incorporates into the top 2 inches of soil Avoid damaging potato seed pieces or sprouts with incorporation equipment

For pre-emergence application apply Charger Max at a rate of 1 to 2 pints per acre after planting as a pre-emergence delayed pre emergence after drag-off or hilling treatment but before weeds emerge If soil organic matter is between 6% and 20% up to 2.5 pints per acre of Charger Max may be used

Do not use on peat or mulch soils if cool wet soil conditions occur after application Charger Max may delay maturity and/ or reduce yields of Superior and other early-maturing potato varieties To avoid crop injury do not use on sweet potatoes or yams do not apply as both pre-emergence and incorporated treatments Do not apply in Kern County CA

Tank mixtures for potatoes

Sencor

Charger Max may be tank mixed with Sencor to control those weeds controlled by Charger Max alone plus hairy nightshade hemp sesbania jimsonweed lambsquarters prickly sida ragweed smartweed velvetleaf Venice mallow and wild mustard Cocklebur will be partially controlled

Apply pre-emergence on coarse soils at a rate of 1 to 1.33 pints per acre Charger Max or on other soil textures a rate of 1.33 to 2 pints per acre Charger Max plus the rate indicated on the Sencor label for this use. Within this rate range use the higher rate on fine textured soils or those high in organic matter and the lower rates on soils that are coarse-textured or low in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. Charger Max does not control emerged weeds. Read and follow all use directions, precautions, and restrictions on the Sencor label.

Do not use in Kern County, CA Do not apply to sweet potatoes or yams. Do not use in peat or muck soils. Potatoes treated with Sencor may not be harvested within 60 days of application or illegal residues may result.

Linuron

East of the Rocky Mountains only. Charger Max may be mixed with linuron products for pre-emergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off at the rates listed in the table below.

Soil texture	Broadcast rate per acre			
	1% to less than 3% organic matter		3% to 5% organic matter	
	Charger Max	linuron*	Charger Max	linuron*
coarse sandy loam	1 pint	1 to 1.5 lb	1.33 pints	1.5 to 2 lb
medium loam silt loam silt	1.33 pints	1.5 to 2 lb	1.67 to 2 pints	2 to 2.5 lb
* Assumes 50% DF formulation. If using other formulations use the equivalent amount of active ingredient. Do not use on sands or loamy sands. Do not incorporate or spray over emerged potatoes.				

Pendimethalin (e.g. Prowl)

A tank mixture with pendimethalin will control the weeds listed on this label for Charger Max alone plus problem species such as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the pendimethalin label. Apply Charger Max plus pendimethalin pre-emergence or pre-emergence incorporated using the rates in the table below. Read and follow all use directions, restrictions, and precautions on both labels and follow the most restrictive label.

Soil texture	Broadcast rate per acre	
	less than 3% organic matter	3% organic matter or more
	Charger Max + pendimethalin*	Charger Max + pendimethalin*
coarse	1 to 1.33 pints + 1 to 1.5 pints	1 to 1.33 pints + 1 to 1.5 pints
medium	1.33 pints + 1.5 to 2 pints	1.33 to 1.67 pints + 2 to 3 pints
fine	1.33 to 1.67 pints + 2 to 3 pints	1.67 to 2 pints + 3 pints
* Pendimethalin rate based a 4E (4 lb/ gallon) formulation. If using other formulations use equivalent amount of active ingredient.		

Pendimethalin + Eptam

A tank mixture with pendimethalin + Eptam will control the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those on the pendimethalin and Eptam labels Follow the directions above for Charger Max plus pendimethalin and add Eptam 7E at a rate of 3.5 to 7 pints per acre Refer to the Eptam label for specific directions precautions and restrictions

SAFFLOWER

Charger Max alone

Follow the directions for preplant incorporated or pre-emergence application of Charger Max alone in the **Application Procedures** section of this label On coarse soils apply 1 to 1.33 pints per acre if organic matter is less than 3% or 1.33 pints per acre if the organic matter is 3% or more On medium soils apply 1.33 to 1.67 pints per acre On fine soils apply 1.33 to 1.67 pints per acre if the organic matter is less than 3% or 1.67 to 2 pints per acre if the organic matter is 3% or more

SORGHUM - GRAIN OR FORAGE

APPLY CHARGER MAX ALONE OR IN TANK MIXTURE ONLY TO SORGHUM SEED THAT HAS BEEN TREATED BY THE SEED COMPANY WITH CONCEP®. OTHERWISE SEVERE CROP INJURY WILL RESULT Under high soil moisture conditions prior to sorghum emergence, crop injury may occur following the use of Charger Max The crop will normally outgrow this effect Do not use Charger Max on sorghum grown under dry mulch tillage, or crop injury may occur Do not apply more than once per crop year (except as a split application described below) or illegal residues may occur

Charger Max alone

Apply preplant surface preplant incorporated or pre-emergence using the rates specified below

Preplant surface applied

For minimum tillage or no-till systems only Charger Max may be applied up to 45 days before planting in IA IL eastern KS MO NE and SD Use only split applications for treatments made 30 to 45 days prior to planting with 2/3 of the rate applied initially and the remaining applied at planting Apply 1.5 pints per acre Charger Max on medium soils or 1.67 pints per acre on fine soils Treatments made less than 30 days before planting may be made either as a split or single application Apply 1.33 pints per acre of Charger Max on coarse soils not more than 2 weeks prior to planting Under dry conditions irrigation after application is recommended to move Charger Max into the soil

Preplant incorporated or pre-emergence

Broadcast 1 to 1.33 pints per acre of Charger Max on coarse soils 1.33 to 1.5 pints per acre on medium soils or 1.33 to 1.67 pints per acre on fine soils

Tank Mixtures for Sorghum

Atrazine

If applying Charger Max in combination with Atrazine follow all restrictions and limitations on the Atrazine label if more restrictive than those on this label. If Atrazine is applied at rates lower than those recommended on this label broadleaf weed control may be adversely affected. Refer to the Atrazine label for weeds controlled at reduced rates. Applications on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.

A tank mixture of Charger Max + Atrazine applied preplant surface preplant incorporated or pre-emergence will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus cocklebur common purslane hairy nightshade lambsquarters morningglory ragweed smartweed and velvetleaf.

For preplant surface applications in minimum-tillage or no-till systems only Charger Max plus Atrazine may be applied up to 45 days before planting in IA IL eastern KS MO NE and SD. Use only split applications for treatments made 30 to 45 days prior to planting with 2/3 of the rate applied initially then the remaining 1/3 applied at planting. Treatments made less than 30 days prior to planting may be made as either split or single applications. Do not use on coarse soils. Apply 1.5 pints per acre of Charger Max + 1.7 to 2 lb per acre Atrazine 90DF* on medium soils with 1.5% organic matter or more. Do not use on medium soils with less than 1.5% organic matter. Apply 1.5 pints per acre of Charger Max + 1.7 to 2 lb per acre Atrazine 90DF on fine soils with less than 1.5% organic matter or apply 1.67 pints per acre Charger Max + 2 to 2.2 lb per acre Atrazine 90DF on fine soils with 1.5% organic matter or more. Under dry conditions irrigation after application is recommended to move the herbicides into the soil.

** If using other formulations of Atrazine use equivalent amount of active ingredient. One lb of Atrazine 90DF is equivalent to 1.8 pints of Atrazine 4L.*

For preplant incorporated or pre-emergence applications apply at the following rates. On medium soils with 1.5% organic matter or more apply 1 pint per acre Charger Max with 1.3 lb Atrazine 90DF*. On fine soils with less than 1.5% organic matter, apply 1 pint per acre Charger Max with 1.3 lb per acre Atrazine 90DF. On fine soils with 1.5% or more organic matter apply 1.2 to 1.33 pints per acre Charger Max with 1.6 to 1.8 lb per acre of Atrazine 90DF.

Precautions To avoid crop injury do not use on medium soils with less than 1.5% organic matter. Do not use on coarse soils. Do not use in NM OK or TX with the following exceptions. In TX use only in the gulf coast and blacklands areas. In OK use only in north-eastern OK. Do not apply preplant incorporated in AZ or the Imperial Valley in CA.

** If using other formulations of Atrazine use equivalent amount of active ingredient. One lb of Atrazine 90DF is equivalent to 1.8 pints of Atrazine 4L.*

Tank Mixtures For Minimum Tillage Or No-Till Systems

Paraquat, Cornerstone (glyphosate), or Landmaster BW

In minimum tillage or no-till systems where sorghum seed treated with Concep is planted directly into a cover crop stale seedbed established sod or previous crop residues paraquat (e.g. Gramoxone Max) glyphosate (e.g. Cornerstone) or Landmaster BW may be added to a mixture of Charger Max plus Atrazine. Mix with paraquat for control of most emerged annual weeds and suppression of perennial weeds. Mix with Landmaster BW for suppression of emerged field bindweed and control or suppression of annual weeds. Mix with Cornerstone (glyphosate) for control of most emerged annual or perennial weeds. Apply before, during, or after planting but before crop emerges at the rates indicated below.

Paraquat

	Weed Height		
	1 to 3 inches	3 to 5 inches	5 to 6 inches
Rate per acre paraquat (as ionized active ingredient*)	0.56 to 0.75 lb ai	0.75 to 0.94 lb ai	0.94 to 1.13 lb ai
* e.g. Gramoxone MAX contains 3 lb ai per gallon	Add 1 to 2 pints of high-active-ingredient non-ionic surfactant such as Activate Plus per 100 gallons of spray mix. Do not apply in suspension-type fluid fertilizers. This moisture will not control weeds taller than 6 inches.		

Landmaster

Apply at a rate of 1.7 to 3.4 pints per acre depending on weed species and size. See the Landmaster BW label for specific instructions.

Cornerstone (glyphosate)

See the Cornerstone label for specific directions. Apply in a minimum of 20 gal of water per acre with conventional spray equipment.

SOYBEANS

Charger Max alone

Apply Charger Max preplant surface-applied, preplant incorporated, pre-emergence, or postemergence using the appropriate rate specified below.

Preplant surface applied, fall application

Apply after Sept. 30 in ND, SD, MN, WI, and north of Route 30 in IA. Apply after Oct. 15 north of Route 91 in NE and south of Route 30 in IA. Apply after Oct. 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a depth of 4 inches is 55° F and falling. In minimum-tillage or no-till systems in soils

having greater than 2.5% organic matter use 1.67 to 2 pints per acre on medium textured soils and 2 pints per acre on fine soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/ or spring tillage operation may follow the application but do not exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in tillage.

NOTE: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate on soybeans or illegal residues might result.

Use on medium and fine soils with minimum tillage or no-till systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply 2/3 the recommended rate (2/3 of 1.67 to 2 pints = 1.11 to 1.33 pints) as a split treatment 30 to 45 days before planting and then apply the remainder at planting. Applications made less than 30 days before planting may be made as either a split or single treatment.

On coarse soils, apply 1.33 pints per acre not more than 2 weeks before planting.

Preplant incorporated or pre-emergence

On coarse soils, apply 1 to 1.33 pints per acre of Charger Max if organic matter is less than 3% or 1.33 pints per acre if organic matter 3% or more. On medium soils, apply 1.33 to 1.67 pints per acre of Charger Max. On fine soils, apply 1.33 to 1.67 pints per acre of Charger Max if organic content is less than 3% or 1.67 to 2 pints per acre if organic matter content is 3% or more.

NOTE: Charger Max may be used as a preplant surface applied, preplant incorporated or re-emergence treatment on soybeans at a rate up to 2.5 pints per acre on soils having an organic matter content between 6% and 20%. The total Charger Max rate applied to soybeans during any one crop season should not exceed 2.5 pints per acre.

Post emergence

Apply 1 to 1.33 pints per acre as a post emergence application to soybeans. This treatment will control only weeds that have not yet emerged. Apply from emergence of soybeans through the third trifoliolate leaf stage. A post emergence treatment with Charger Max may not be made if a preplant or pre-emergence application of Charger Max or other metolachlor product was already made.

NOTE: To avoid possible illegal residues following postemergence application to soybeans, do not apply more than 1.33 pints per acre, make postemergence applications at least 90 days before harvest, and do not graze or feed treated forage or hay from soybean crops to livestock following postemergence application.

Tank mixtures/ sequential treatments for soybeans

Charger Max may be diluted for application with water or fluid fertilizer when tank-mixed or used sequentially with the following herbicides: Sencor, linuron, Canopy, Pursuit, Scepter, ethalfluralin (e.g. Sonalan) or Command 3ME. For these mixtures, Charger Max may be applied at up to 2.33 pints per acre on soils having organic matter between 6% and 20%. The total rate from Charger Max or any other brand of s-metolachlor

applied to soybeans in any one crop year may not exceed the equivalent of 2.5 pints per acre

Refer to the Sencor label for planting details and soybean variety restrictions

Sencor

A tank mixture of Charger Max + Sencor applied preplant incorporated or pre-emergence at the rates specified in the following table will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus cocklebur hairy nightshade hemp sesbania jimsonweed (partial control only) lambsquarters prickly sida ragweed smartweed Venice mallow and wild mustard

Broadcast rate per acre		
Organic matter	0.5% up to 3%	3% or more
Soil texture	Charger Max + Sencor*	Charger Max + Sencor*
COARSE** - loamy sand (over 2% organic matter) sandy loam	0.8 to 1 pint + 0.33 lb	1 pint + 0.5 lb
MEDIUM	1 to 1.33 pints + 0.5 lb	1.33 pints + 0.67*** lb
FINE	1.33 pints + 0.67 lb	1.33 to 1.67 pints + 0.67 lb
Mississippi delta only silty clay clay	1.33 pints + 1 lb	1.33 to 1.67 pints + 1 lb
Muck or peat soils (more than 20% organic matter)	DO NOT USE	
NOTES		
* Assumes a 75% DF formulation. If using a 4L, multiply lb by 1.5 to get pints per acre		
** Do not use on any sand or loamy sand with less than 2% organic matter. Do not use this tank mixture on coarse soils preplant incorporated or crop injury might occur		
*** Use 0.5 lb Sencor if preplant incorporated		

For sequential application with Sencor, apply Charger Max alone preplant incorporated at rates specified in the table above. Follow with a pre-emergence application of Sencor during planting (behind the planter) or after planting but before weeds or soybeans emerge.

Linuron

A tank mixture of Charger Max + linuron applied pre-emergence at the rates specified in the following table will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus cocklebur jimsonweed (partial control only) lambsquarters morningglory (partial control only) prickly sida ragweed smartweed velvetleaf Venice mallow and wild mustard

Apply during planting (behind planter) or after planting but before weeds or soybeans have emerged. Refer to the linuron label for planting details. Do not use on soils with less than 0.5% organic matter or crop injury might occur. Do not use on sand, gravelly soils or exposed subsoils.

Broadcast rate per acre			
Organic matter	0.5% up to 3%		3% or more
Soil texture	Charger Max +	linuron*	Charger Max + linuron*
COARSE**	0.8 pint	+ 1 lb	1 pint + 1 to 1.5 lb
MEDIUM	1 pint	+ 1 to 1.5 lb	1.33 pints + 1.5 to 2 lb
FINE	1.33 pints	+ 2 lb	1.33 to 1.67 pints + 2.5 to 3 lb
Muck or peat soils (more than 20% organic matter)	DO NOT USE		
NOTES			
* Assumes a 50% DF formulation. If using a different formulation, use the equivalent amount of active ingredient.			
** Do not use on loamy sand except in the north eastern US on loamy sand with more than 1% organic matter.			

Trust (trifluralin)

A tank mixture of Charger Max + Trust (trifluralin) applied preplant incorporated will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those listed on the trifluralin label for trifluralin alone. This mixture may be applied by ground or aerial equipment and incorporated up to 14 days before planting. Follow directions, restrictions and precautions on both labels and use equipment that provides a uniform incorporation 2 inches deep. Apply at the rates specified on each label for Charger Max and trifluralin applied alone for the specific soils, texture, organic matter, and weed species expected.

To control DNA-resistant goosegrass and other species on the respective labels where the soil organic matter is 3% or less, apply at the rates in the following table. Where a range of rates is given, use the minimum rate where DNA-resistant goosegrass is the predominant species.

Soil Texture	Broadcast Rate per Acre		
	Charger Max	Trust 4EC*	
	Organic Matter	Organic Matter	
	less than 3%	less than 2%	2 to 3%
COARSE	0.8 to 1 pint	1 pint	1.5 pints
MEDIUM	1 pint	1.5 pints	1.5 pints
FINE	1.33 pints	2 pints	2 pints
* If using an alternate formulation of trifluralin, apply the equivalent amount of active ingredient.			

Scepter

A tank mixture of Charger Max + Scepter 70DG applied preplant incorporated or pre-emergence at the rates specified in the following table will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those listed on the Scepter label for Scepter alone. For preplant incorporated application, incorporate within 30 days before planting. Follow directions, restrictions and precautions on both labels and use equipment that provides a uniform incorporation.

2 inches deep Follow use directions under Application Instructions on the Scepter label and observe geographical restrictions on the Scepter label

Soil Texture	Broadcast Rate per Acre			
	Less than 3% organic matter		3% or more organic matter	
	Charger Max	Scepter 70DG	Charger Max	Scepter 70DG
COARSE	0 8 pint	0 18 lb	1 pint	0 18 lb
MEDIUM	1 pint	0 18 lb	1 33 pints	0 18 lb
FINE	1 33 pints	0 18 lb	1 33 to 1 67 pints*	0 18 lb
Muck or peat soils with more than 20% organic matter	DO NOT USE			
* Use the higher rate of Charger Max if heavy weed infestations are expected NOTES Do not apply within 90 days of harvest Do not graze or feed treated soybean forage hay or straw to livestock before 30 days following treatment or illegal residues might result				

Canopy SP

This tank mixture applied preplant incorporated or pre-emergence at the rates listed below will control all weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus all those for Canopy SP alone on the Canopy SP label For preplant incorporated application apply within 2 weeks of planting Uniformly incorporate into the top 1 to 2 inches of soil before planting soybeans For pre-emergence application apply after planting but before soybeans emerge NOTE Follow all directions precautions limitations varietal restrictions and varietal restrictions on the labels of both tank mix partners

Soil Texture	Broadcast Rate per Acre			
	0 5 to less than 3% organic matter		3 to 5 % organic matter	
	Charger Max	Canopy SP	Charger Max	Canopy SP
COARSE	0 8 pint	7 7 to 12 9 oz	1 pint	7 7 to 12 9 oz
MEDIUM	1 pint	10 3 to 15 4 oz	1 33 pints	10 3 to 15 4 oz
FINE	1 33 pint	12 9 to 18 oz	1 33 to 1 67 pints	12 9 to 18 oz
Precautions Do not apply to sand nor to any soil with less than 0 5 % organic matter nor to any soil with pH greater than 7 0 (alkaline soil) except as noted on the Canopy SP label				

Command 3ME

This tank mixture applied preplant incorporated at the rates listed in the table below will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label plus those on the Command 3ME label for Command 3ME alone Follow all use directions limitations precautions and rotational restrictions on labels of both products Follow all Command 3ME instructions regarding incorporation interval geographic restrictions equipment operation, soil moisture etc

Soil Texture	Broadcast Rate per Acre			
	Charger Max		Command 3ME	
	0 5 to 3% organic matter	More than 3% organic matter	Northern area	Southern area
COARSE	0 8 pint	1 pint	2 - 2 67 pints	2 67 to 3 33 pints
MEDIUM	1 pint	1 33 pints	2 to 2 67 pints	2 67 to 3 33 pints
FINE	1 33 pint	1 33 to 1 67 pints	2 to 2 67 pints	2 67 to 3 33 pints

Ethalfuralin (e.g. Sonalan)

Charger Max may be used in tank mixture with ethalfuralin as a preplant incorporated application at the rates listed in the table below. Charger Max may alternatively be applied pre-emergence as a sequential treatment following a preplant incorporated application of ethalfuralin. In either case, follow the soil preparation procedures specified on the ethalfuralin label. For sequential treatment, follow with a pre-emergence application of Charger Max using rates specified for Charger Max alone during planting (behind the planter) or after planting but before weeds or soybeans emerge. For preplant incorporated application of a tank mixture, use the rates specified in the table below.

Soil Texture	Broadcast Rate per Acre			
	Less than 3% organic matter		3% or more organic matter	
	Charger Max	ethalfuralin*	Charger Max	ethalfuralin*
COARSE	1 to 1.33 pints	1.25 to 2 pints	1.33 pints	1.25 to 2 pints
MEDIUM**	1.33 to 1.67 pints	1.75 to 2.5 pints	1.33 to 1.67 pints	1.75 to 2.5 pints
FINE**	1.33 to 1.67 pints	2.25 to 3 pints	1.67 to 2 pints	2.25 to 3 pints
Muck or Peat (soils with more than 20% organic matter)	DO NOT USE			
* Assumes a 3 lb per gallon formulation. If using an alternate formulation, use the equivalent amount of active ingredient per acre.				
** For eastern black nightshade on medium and fine soils, apply ethalfuralin at 3 pints per acre on medium soils or 3.5 pints per acre on fine soils, and follow with 2 incorporation passes.				
NOTE: Follow all use directions, limitations, precautions, and information regarding application to soybeans on both tank-mix partner labels.				

Pursuit (tank mixture)

This tank mixture applied early preplant, preplant incorporated, or pre-emergence after planting will control all the weeds listed in the WEEDS CONTROLLED BY CHARGER MAX ALONE section of this label, plus those on the Pursuit label for Pursuit alone. Refer to the Pursuit label for geographical restrictions regarding where this tank mixture may be applied, as well as rotational crop restrictions. Apply at rates indicated in the table below, diluting in water or liquid fertilizer. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Soil Texture	Broadcast rate per acre		
	less than 3% organic matter	3% organic matter or more	Any % soil organic matter
	Charger Max	Charger Max	Pursuit
COARSE	0.8 pint	1 pint	0.25 pint
MEDIUM	1 pint	1.33 pints	0.25 pint
FINE	1.33 pints	1.33 to 1.67 pints	0.25 pint

Pursuit (sequential)

Apply Charger Max early preplant preplant incorporated or pre-emergence after planting at a rate of 0.8 pint per acre on coarse soils or 1 pint per acre on medium or fine soils. Follow with a sequential postemergence application of Pursuit to control emerged weeds according to the Pursuit label. Charger Max will improve the consistency and level of control by Pursuit on most grass weed species. Refer to the Pursuit label for weeds controlled, rates and growth stage limitations.

Cornerstone (glyphosate) or paraquat (e.g. Gramoxone MAX) plus Sencor, Scepter, linuron, or Pursuit For minimum tillage or no-till systems

In minimum tillage or no-till systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat (e.g. Gramoxone MAX) or Cornerstone (glyphosate) may be added to a tank mixture of either Charger Max + Sencor or + Scepter or + linuron or + Pursuit. The paraquat portion of the mixture will control most emerged annual weeds and suppress many perennial weeds. Cornerstone combinations will control emerged annual and perennial weeds as directed on the Cornerstone label. For additional weeds controlled by Sencor, Scepter, Pursuit or linuron, refer to the respective labels of those products. Read and observe all use directions, planting details, variety restrictions, geographical restrictions, and all other precautions and limitations on the labels of all the products being used. Apply before, during, or after planting but before soybeans emerge at the rates specified below.

With paraquat. Apply at rates specified for Charger Max alone plus the following

	Weed Height		
	1 to 3 inches	3 to 5 inches	5 to 6 inches
Rate per acre paraquat (as ionized active ingredient*)	0.56 to 0.75 lb ai	0.75 to 0.94 lb ai	0.94 to 1.13 lb ai
* e.g. Gramoxone MAX contains 3 lb ai per gallon	Add 1 to 2 pints of high-active-ingredient non-ionic surfactant such as Activate Plus per 100 gallons of spray mix. Do not apply in suspension-type fluid fertilizers. This moisture will not control weeds taller than 6 inches.		

With Cornerstone (glyphosate)

Apply at rates specified for Charger Max alone plus Cornerstone or other glyphosate formulation. Follow the glyphosate label for weeds controlled, rates, and other directions. Apply with ground equipment in 20 to 60 gallons of water or fluid fertilizer per acre.

With Sencor + paraquat or Cornerstone (glyphosate)

Use rates of glyphosate or paraquat described above. On loamy sand with over 2% organic matter, apply 1 pint per acre of Charger Max plus 0.33 to 0.5 lb per acre of Sencor DF. On medium soils, apply 1.33 pints per acre of Charger Max plus 0.5 to 0.67 lb per acre of Sencor DF. On fine soils, apply 1.33 to 1.67 pints per acre of Charger

Max plus 0.67 lb per acre of Sencor DF. If using an alternate formulation of Sencor apply the equivalent amount of active ingredient.

PRECAUTIONS To avoid crop injury do not use this tank mixture on soils with less than 0.5% organic matter on alkaline soil with a pH more than 7.4 or on sand or loamy sand with less than 2% organic matter. If heavy rain occurs soon after application crop injury might result especially in poorly drained areas where water stands for several days or where the seeding slit has not been properly closed.

With Scepter + paraquat or Cornerstone (glyphosate)

Use rates of glyphosate or paraquat described above. On coarse soils apply 1 pint per acre of Charger Max plus 0.67 pints per acre of Scepter. On medium soils apply 1.33 pints per acre of Charger Max plus 0.67 pints per acre of Scepter. On fine soils apply 1.33 to 1.67 pints per acre of Charger Max plus 0.67 pints per acre of Scepter.

NOTES Do not apply within 90 days of harvest and do not graze or feed treated soybean forage, hay or straw to livestock or illegal residues might occur.

With linuron + paraquat or Cornerstone (glyphosate)

On coarse soils apply 1 pint per acre of Charger Max + 1 to 1.5 lb per acre of linuron 50% DF. On medium soils 1.33 pint per acre of Charger Max plus 1 to 2 lb per acre of linuron 50% DF. On fine soils apply 1.33 to 1.67 pints per acre of Charger Max plus 2 to 3 lb per acre of linuron 50% DF. If using a 4L formulation one pint of 4L is equivalent to 1 pound of 50% DF.

PRECAUTIONS To prevent crop injury do not use on loamy sand except in the northeastern US on loamy sand with over 1% organic matter and do not use on sand, gravelly soils or exposed subsoil. Do not use on any soil with less than 0.5% organic matter.

With Canopy + paraquat or Cornerstone (glyphosate)

Use rates of glyphosate or paraquat described above. Use only where soil has 0.5% to 5% organic matter. On coarse soils (except sand) apply 1 pint per acre of Charger Max on medium soils apply 1.33 pints per acre Charger Max and on fine soils apply 1.33 to 1.67 pints per acre of Charger Max. Apply in mixture with Canopy as specified on the Canopy label for the geographic location, soil type, pH and other limitations.

PRECAUTIONS Do not apply to sand nor to any soil with less than 0.5% organic matter nor to any soil with a pH greater than 7.0 except as specified on the Canopy label.

With Pursuit + paraquat or Cornerstone (glyphosate)

On coarse soils apply 1 pint per acre of Charger Max plus 0.25 pint per acre of Pursuit. On medium soils apply 1.33 pints per acre of Charger Max plus 0.25 pint per acre of pursuit. On fine soils apply 1.67 pints per acre of Charger Max plus 0.25 pint per acre.

of Pursuit Use only where soils have 0.5-5.0% organic matter and do not apply to soils with pH greater than 6.8 Do not apply to sand

Postemergence Tank Mixtures for Soybeans

Cornerstone (glyphosate)

USE THIS MIXTURE ONLY ON GLYPHOSATE TOLERANT OR ROUNDUP READY SOYBEANS

Charger Max may be applied in tank mixture with Cornerstone or other glyphosate products registered for this use from emergence through the first trifoliolate leaf stage of soybeans Charger Max alone will not control emerged weeds DO NOT USE THIS MIXTURE ON SOYBEANS NOT SPECIFIED AS BEING TOLERANT TO GLYPHOSATE APPLIED POSTEMERGENCE Apply at rates specified for the appropriate weeds soils geographic regions etc on this and the glyphosate label

Pursuit

Charger Max may be applied in tank mixture with Pursuit from emergence through the first trifoliolate leaf stage of the soybeans Apply Charger Max at a rate of 1 to 1.33 pints per acre plus Pursuit at rates specified for the appropriate weeds soils geographic regions etc on the Pursuit label Charger Max alone will not control emerged weeds

Liberty (glufosinate)

USE THIS MIXTURE ONLY ON GLUFOSINATE TOLERANT OR LIBERTY LINK SOYBEANS

Charger Max may be applied in tank mixture with Liberty (glufosinate) from emergence through the third trifoliolate leaf stage of the soybeans Apply Charger Max at a rate of 1 to 1.33 pints per acre plus Liberty at rates specified on the Liberty label Charger Max alone does not control emerged weeds DO NOT USE THIS MIXTURE ON SOYBEANS NOT SPECIFIED AS BEING TOLERANT TO GLUFOSINATE APPLIED POSTEMERGENCE

NOTES Follow the glufosinate product label for adjuvant recommendations The use of COC or UAN with Charger Max may result in temporary crop injury To avoid possible illegal residues when Charger Max is applied post-emergence to soybeans do not apply more than 1.33 pints per acre make postemergence applications at least 90 days before harvest and do not graze or feed treated forage or hay to livestock following postemergence application of Charger Max

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT

FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations on liability do not use the product and return it unopened to the Seller and the purchase price will be refunded

Trademark information

Activate Plus and AgriSolutions are trademarks and Charger Max Complete Compatibility Cornerstone Moxy Prime Oil Silkin and Trust are registered trademarks of Winfield Solutions LLC

RoGator and Spra-Coupe are trademarks of AGCO Corporation

HiCycle is a trademark of John Deere and Company

Tyler Patriot is a trademark of Case Corporation

Accent Lorox and Canopy are trademarks of E I DuPont de Nemours and Co

Sencor Balance Liberty Link and Liberty are trademarks of Bayer Crop Sciences

Balan and Sonalan are trademarks of Dow AgroSciences

Prowl Marksman Basagran Scepter and Pursuit are trademarks of BASF Ag Products

Agsorb is a trademark of Oil-Dri Corporation

Celatom MP 79 is a trademark of Eagle-Picher Industries

Aromatic 200 is a trademark of ExxonMobil Corp

Command is a trademark of FMC Corp

Storm is a trademark of United Phosphorus Inc

Concep Beacon Exceed Gramoxone Evik and Dual are trademarks of Syngenta Crop Protection Inc

Eptam and Eradicane are trademarks of Gowan Company LLC

Sutan is a trademark of Helm Agro US Inc

Landmaster is a trademark of The Monsanto Company