

1381-202

7/25/2014

1/28



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Danielle Erding
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164

JUL 25 2014

Subject: Notification; Per PR-Notice 98-10 & 2001-5
Brash
EPA Reg. No. 1381-202
Date Submitted: July 16, 2014

Dear Ms. Erding:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notices (PRN) 98-10 and 2001-5 dated July 16, 2014 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and 2001-5 and finds that the action requested falls within the scope of PRN 98-10 and 2001-5. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Kable Bo Davis at (703) 306-0415 or davis.kable@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "K. Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

for

EPA	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
------------	---	---	-----------------------

Application for Pesticide -Section I

1. Company/Product Number 1381-202	2. EPA Product Manager Kathryn Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Brush	PM# 23	

5. Name and Address of Applicant (Include ZIP Code) Winfield Solutions, LLC P.O. Box 64589 St. Paul, MN 55164-0589 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition to the product identified below. EPA Reg. No.: Product Name:
--	---

NOTIFICATION
JUL 25 2014

Section II

<input type="checkbox"/> Amendment - Explain Below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - explain below

Explanation: Use additional page(s) if necessary. (For Section I and Section II). Notification of Addition of Resistance Management Recommendations and MOA code to label, consistent with PR Notice 2001-5, and per PR Notice 98-10. I believe this notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR-Notice 98-10 and 40 CFR-152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Section III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No <small>*Certification must be submitted.</small>	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Unit Package Wt. No. Per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes," Package Wt. No. Per Container	2. Type of Container <input checked="" type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container 1, 2.5, 30, 55 gallons	
5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product			
6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other (self-adhesive booklet) <input type="checkbox"/> Paper Glued <input type="checkbox"/> Stenciled			

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Danielle Erding	Title Registration Specialist	Telephone No. (Include Area Code) 651-375-7228
-------------------------	----------------------------------	---

Certification		6. Date Application Received (Stamped)
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		
2. Signature <i>Danielle Erding</i>	3. Title Registration Specialist	
4. Typed Name Danielle Erding	5. Date July 16, 2014	

NOTIFICATION

JUL 25 2014

4/
28

GROUP 4 HERBICIDE

BRASH®

For Use on Conservation Reserve Program Land, Fallow Systems (Between Crop Applications), General Farmstead, Sorghum, Grass (Hay or Silage), Pastures, Rangeland, Rights-of-Way, Sugarcane, and Wheat

ACTIVE INGREDIENTS:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)*	12.4%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid**	35.7%
OTHER INGREDIENTS:	51.9%
TOTAL	100.0%

*This product contains 10.3% 3,6-dichloro-o-anisic acid (dicamba) or 1 pound per gallon (120 g/L)

**This product contains 29.6% 2,4-dichlorophenoxyacetic acid (2,4-D) or 2.87 pounds per gallon (344 g/L). Isomer specific by AOAC method 978.05, 15th Edition

**SHAKE WELL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

See inside booklet for additional Precautionary Statements

EPA Reg. No. 1381-202

EPA Est. No.

Distributed By:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0089

NET CONTENTS _____
LOT NO. _____
1/1212/3 107164

WINFIELD™

AGROSOLUTIONS™

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER**

Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Harmful if swallowed or absorbed through skin. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are butyl rubber \geq 14 mils, or natural rubber \geq 14 mils, or neoprene rubber \geq 14 mils or nitrile rubber \geq 14 mils. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (except for applicators using groundboom equipment, pilots and flaggers)
- Shoes plus socks
- Protective eyewear
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements and exceptions.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering Controls Statements:

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow,

may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et.al. v. EP, C01-0132C, (W.D. WA). For further information, please refer to: <http://www.epa.gov/espp/litstatus/wtc/index.htm>.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over short-sleeve shirt and short pants, chemical-resistant gloves made of any waterproof material, chemical-resistant headgear for overhead exposure, chemical-resistant footwear plus socks, and protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Pesticide wastes are toxic. Triple rinse pesticide from containers and use rinsates in the pesticide application. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Use label language appropriate for container size and type.

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call
CHEMTREC 1-800-424-9300.**

PRODUCT INFORMATION

BRASH® is a selective postemergence herbicide for controlling a wide spectrum of annual, biennial, and perennial broadleaf weeds and brush in grass forages and selected row crops. BRASH® may be used in/on Conservation Reserve Program Land*, Fallow Systems (Between Crop Applications)*, General Farmstead*, Grain Sorghum, Grass (Hay or Silage), Pastures, Rangeland, Sugarcane, and Wheat.

*These crops are considered Food/Feed crops only when harvested, grazed or foraged. Otherwise, they are considered as non-Food/Feed uses.

MODE OF ACTION: BRASH® contains dicamba and 2,4-D as the active ingredients. BRASH® is readily absorbed by plants through shoot and root uptake, translocates throughout the plant's system, and accumulates in areas of active growth. BRASH® interferes with the plant's growth hormones (auxins) resulting in death of many broadleaf weeds.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

Brash is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Brash and other Group 4 herbicides. Weed species with acquired resistance to Group 4 may eventually dominate the weed population if Group 4 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Brash or other Group 4 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of Brash or other target site of action Group 4 herbicides that have a similar target site of action, on the same weed species.

- Using tank-mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or Winfield Solutions, LLC representative for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

SPRAY EQUIPMENT CLEANING: Spray equipment may be cleaned by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and then triple rinsing the equipment before and after applying this product.

APPLICATION INSTRUCTIONS

Apply BRASH® at the specified rates and growth stages in the **Annual Weeds** and the **Biennial and Perennial Weeds** rate tables unless instructed differently in the **Food/Feed Crop Specific Information** or **Non-Food/Feed Use-Specific Information** sections of this label. Make applications of BRASH® to actively growing weeds using aerial, broadcast, band, or spot spray applications. BRASH® may be applied using water or sprayable fluid fertilizer as a carrier. For preplant or pre-emergence uses, sprayable fluid fertilizer may be used as the carrier for all crops listed on this label. Postemergence applications with sprayable fluid fertilizer may be made on pasture, hayland, or wheat crops only.

The most effective application rate and timing varies based on the target weed species. In mixed weed populations, the correct rate is determined by the weed species requiring the highest rate. Inadequate control may be observed if application is delayed since weeds may exceed the maximum size stated on this label.

IRRIGATION: In irrigated areas, it may be necessary to irrigate before application of BRASH® to ensure active weed growth.

SPRAY COVERAGE: Ensure weeds are thoroughly covered with spray. Dense leaf canopies may shield smaller weeds and prevent adequate coverage.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if:

- a) conditions of temperature inversion exist, or
- b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. These plants are most sensitive to BRASH® during their development or growing stage. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants. Agriculturally approved drift-reducing additives may be used. Do not use aerial equipment to apply BRASH® when sensitive crops and plants are growing in the vicinity of area to be treated. Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of BRASH® with the roots of desirable plants such as trees and shrubs.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

AERIAL APPLICATION METHODS AND EQUIPMENT

Water Volume: Use 3-10 gallons of water per acre. Use the higher spray volume when treating dense or tall vegetation.

GROUND APPLICATION (BANDING)

When applying BRASH® by banding, determine the amount of herbicide and water volume needed using the following formula:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Banding herbicide rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre} = \text{Banding water volume per acre}$$

GROUND APPLICATION (BROADCAST)

For optimal performance, use 5-40 gallons of spray solution per broadcast acre. Use the higher spray volume when treating dense or tall vegetation.

SPOT OR SMALL AREA APPLICATION

BRASH® may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to runoff) of foliage and stems. For knapsack or other small capacity sprayers, utilize the table below to calculate material needed. (The table below is based on the assumption that the spot treatment rate equates to 60 gallons per acre on the broadcast basis.)

Knapsack Sprayer Dilution Instructions

Sprayer Capacity (Gallons of water)	Fluid Ounces* of BRASH® to add per filled tank
1.0 gallon	1.0 oz
2.5 gallons	2.5 oz
3.0 gallons	3.0 oz
5.0 gallons	5.0 oz

The addition of a surfactant can help improve control. Add ½% (0.005) by volume. For example, 5 gallons (40 pt/640 fl oz) of herbicide solution would require 0.2 pt (3.2 fl oz) of surfactant.

*1 fluid ounce = 2 tablespoons and 1 cup (liquid) = 16 tablespoons

Weeds Controlled	BRASH® Rate Per Acre (according to weed growth stage)					
	0.5 pint	1 pint	1.5 pints	2 pints	3 pints	4 to 5.6 pints
¹ May require repeat applications. ² Listed rate provides top growth suppression only. ³ For improved root kill of woody species such as mesquite and eastern persimmon, spray 4 pints of BRASH® per acre each year for 3 consecutive years. For increased control of weeds such as blackberry and dewberry, BRASH® may be tank mixed with Ally® herbicide (0.1-0.2 ounces per acre), if labeled for the use site. ⁴ A second application may be required the following growing season under dense populations. ⁵ Not for use in California. ⁶ Treat with 4 pints of BRASH® per acre after full leaf but before leaves develop a heavy cuticle (waxy covering) in periods of extreme heat or drought stress.						

ADDITIVES

For improved burndown of emerged weeds, surfactants and/or low use rate of liquid fertilizers (28-0-0, 32-0-0), or crop oil concentrate may be used with BRASH® or BRASH® tank mixes applied after weeds have emerged. Crop oil concentrate is for non-food/feed crop uses only. Do not apply to tank mixes that include ammonium sulfate or crop oil concentrate to any food/feed crop use listed on this label. For food/feed crop uses, do not use liquid fertilizers that contain ammonium sulfate (AMS) as a source of nitrogen as tolerances in commodities derived from the crop may contain residues that exceed established tolerances. Consult your local WINFIELD SOLUTIONS, LLC representative for recommendations for your area. For additional information, refer to the **Compatibility Test for Mix Components** section of this label.

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria: be nonphytotoxic, contain only EPA-exempt ingredients, provide good mixing quality in the jar test and be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, refer to the **Compatibility Test for Mix Components** section of this label.

Mix Components

Adjuvants containing crop oil concentrates may be used for preplant, pre-emergence and between cropping applications. Do not use crop oil concentrate for postemergence applications in food/feed crops (i.e., sorghum, grass (hay or silage), pastures, rangeland, sugarcane and wheat).

Nitrogen Source

Sprayable Liquid fertilizers: Use one quart of sprayable liquid fertilizers (28-0-0, 32-0-0) per acre. Do not use brass or aluminum nozzles when spraying fertilizers.

Nonionic Surfactant

The standard label recommendation is 2-4 pints of an 80% active nonionic spray surfactant per 100 gallons of water. For certain weeds, use a higher spray surfactant rate. When an adjuvant is to be used with this product, Winfield Solutions, LLC recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Additive Rate Per Acre

Additive	Rate Per Acre
Nonionic Surfactant	2-4 pints per 100 gallons
Sprayable liquid fertilizers (28-0-0, 32-0-0)	2-4 quarts
Crop Oil Concentrate	1 quart*

*Refer to the manufacturer's label for specific rates.

PRODUCT TANK MIXING INFORMATION

The following products may be tank mixed with BRASH[®] according to the specific tank mixing instructions in this label and the respective product labels.

Aim [™] (carfentrazone-ethyl)	Gramoxone [®] Extra (paraquat)
Ally [®] (metsulfuron-methyl)	Harmony [®] Extra (thifensulfuron + tribenuron-methyl)
Amber [®] (triasulfuron)	Karmex [®] (diuron)
Asulox [®] (asulam)	Kerb [™] (pronamide)
Atrazine	Laddok [®] S-12 (bentazon + atrazine)
Albaugh Dicamba DMA Salt or Banvel [®] (dicamba)	Landmaster [®] BW (glyphosate + 2,4-D)
Basagran [®] (bentazon)	MCPA
BROX [™] -M Herbicide or Bronate [®] (bromoxynil + MCPA)	Paramount [®] (quinclorac)
BROX [™] 2EC Herbicide or Buctril [®] (bromoxynil)	Peak [®] (prosulfuron)
Canvas [®] (thifensulfuron + tribenuron + metsulfuron)	Permit [®] (halosulfuron-methyl)
Clarity [®] (dicamba)	Rave [™] (dicamba + triasulfuron)
Curtail [™] (clorpyralid + 2,4-D)	Gly Star [™] Plus or Roundup [®] Ultra (glyphosate)
Cyclone [®] (paraquat)	Sencor [®] (metribuzin)
Dakota [®] (fenoxaprop-p-ethyl + MCPA)	Sinbar [®] (terbacil)
Distinct [®] (diflufenzopyr)	Stinger [™] (clopyralid)
Evik [®] (ametryn)	Tiller [®] (fenoxaprop-p-ethyl + 2,4-D + MCPA)
Express [®] (thifensulfuron + tribenuron-methyl)	Tordon [™] (picloram)
Fallow Star [™] or Fallowmaster [®] (glyphosate + dicamba)	Touchdown [®] (sulfosate)
Finesse [®] (chlorsulfuron + metsulfuron-methyl)	2,4-D
Glean [®] (chlorsulfuron)	

Refer to the Food/Feed Crop-Specific Information section for additional information. Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Crop injury, reduced weed control, or physical incompatibility may result when mixing BRASH[®] with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. Winfield Solutions, LLC does not recommend using tank mixes other than those listed on this labeling. Local agricultural authorities may be a source of information when using tank mix components other than those recommended on this label.

Compatibility Test for Mixing Components

Always perform a compatibility test before mixing components. For 20 gallons per acre spray volume, use 3.3 cups (800 mL) of water. For other spray volumes, adjust accordingly. Only use water from the intended source at the source temperature. Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order

If an inductor is used, rinse it thoroughly after each component has been added. Maintain constant agitation during application.

1. Water* - Begin by agitating a thoroughly clean sprayer tank half full of clean water.
2. Agitation – Maintain constant agitation throughout mixing and application.
3. Products in PVA bags – Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. Water-dispersible products (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
5. Water-soluble products (such as BRASH®).
6. Emulsifiable concentrates (such as oil concentrate when applicable).
7. Water-soluble additives (such as liquid fertilizers (28-0-0, 32-0-0) when applicable).
8. Remaining quantity of water.

*If sprayable fluid fertilizer is used as the carrier, BRASH® must be diluted with a minimum of 5 parts water to 1 part BRASH®. Then add 0.25-0.05% volume/volume of a nonionic surfactant to the dilution before adding it to the sprayable fluid fertilizer to reduce the concern for compatibility problems with this mix. Always perform the **Compatibility Test** before mixing into the spray tank. Also, when using a sprayable fluid fertilizer as the carrier, any product contained in PVA bags must first be completely dissolved in water before the contents can be added to the fertilizer mix.

RESTRICTIONS AND LIMITATIONS

Crop	Livestock Grazing or Feeding ¹	Aircraft Application
Between Crop Applications	Yes	Yes
Pasture, Hay, Silage	Yes	Yes
Sugarcane	Yes	Yes
Sorghum	Yes	Yes
Wheat	Yes	Yes
¹ Refer to the Food/Feed Crop-Specific Information section of this label for grazing and feeding restrictions.		

- **Maximum single application rate:** 1 gallon per acre with no more than 2 applications per year.
- **Preharvest Interval (PHI):** Refer to the **Food/Feed Crop-Specific Information** section of this label.
- **Crop Rotational Restrictions:**
The interval between application and planting rotational crop is provided below. Always exclude counting days when the ground is frozen. Planting at intervals less than specified below may result in crop injury. Moisture is essential for the degradation of this herbicide in soil. If dry weather prevails, use cultivation to allow herbicide contact with moist soil.
 - **Planting/replanting restrictions for BRASH® applications of 6 pints per acre or less:** No rotational cropping restrictions apply at 120 days or more following application. Additionally, for annual crop uses in this label including sorghum, follow the preplant use directions in the **Food/Feed Crop-Specific Information** section of this label. For barley, oat, wheat, and other grass seedlings, the interval between application and planting is 10 days per pint per acre.

- **Planting/replanting restrictions for application of more than 6 pints and up to 8 pints of BRASH[®] per acre:** Corn, sorghum, cotton (east of the Rocky Mountains) and all other crops grown in areas with 30" or more of annual rainfall may be planted 120 days or more after application. Barley, oat, wheat, and other grass seedlings, may be planted if the interval from application to planting is 10 days per pint per acre east of the Mississippi River and 15 days per pint per acre west of the Mississippi River. For all other crops in areas with less than 30" of annual rainfall, the interval between application and planting is 180 days or more.
- **Rainfast period:** The effectiveness of BRASH[®] may be reduced if rainfall or irrigation occurs within 4 hours after postemergence applications.
- **Stress:** Unsatisfactory control may result if BRASH[®] is applied to crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures.
- Do not apply to crops that exhibit injury (leaf phytotoxicity or plant stunting) produced by any other, prior herbicide applications, because this injury may be enhanced or prolonged by applying BRASH[®].
- Do not apply through any type of irrigation equipment. Do not contaminate irrigation ditches or water used for domestic purposes.
- This product cannot be used to formulate or reformulate any other pesticide product.

FOOD/FEED CROP-SPECIFIC INFORMATION

Pastures, Rangeland and Grass (Hay, Silage)

BRASH[®] is recommended for use for pasture (including pasture grown for hay), rangeland and grass grown for hay or silage.

Refer to the **ANNUAL WEEDS** and **BIENNIAL and PERENNIAL WEEDS** Application Rate and Timing tables for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control. Rates above 4 pints of BRASH[®] per acre are for spot treatments only. Retreatments may be made as needed; however, do not exceed a total of 8 pints of BRASH[®] per treated acre during a growing season.

BRASH[®] uses described in this situation also pertain to small grains (such as barley, corn, forage sorghum, oats, rye, sudangrass or wheat) grown for pasture, hay, and silage only. Newly seeded areas, including small grains grown for pasture or hay, may be injured if rates of BRASH[®] greater than 2 pints per acre are applied.

In newly established hybrid Bermudagrass, Pangolagrass, and stargrasses (*Cynodon spp.*), use 1-2 quarts of BRASH[®] per acre to control or suppress weeds after planting vegetative propogules (stolens) of hybrid bermudagrasses. In addition to the weeds listed in the ANNUAL and BIENNIAL and PERENNIAL WEEDS tables, this rate of BRASH[®] will control or suppress annual sedges, broadleaf signalgrass, crabgrass, and goosegrass. Best results will be obtained if BRASH[®] is applied at the germinating stage of weeds. Under favorable conditions, this is usually 7-10 days after planting these grasses. Reduced control can be expected if weeds are allowed to reach 1" in height before application or if germination of weeds occurs 10 days after application.

Do not use on bentgrass, susceptible grass pastures (such as carpetgrass, buffalograss or St. Augustine grass), lespedeza, wild winter peas, vetch, clover and alfalfa pastures as injury will occur.

When perennial weeds are reaching maturity, mowing and allowing some regrowth will enhance control. Difficult to control weeds and brush may require repeat application.

For pasture renovations, wait 3 weeks per quart per acre of BRASH[®] used before interseeding or injury may occur.

If grasses are grown for seed or for seed-down purposes, do not apply after grass reaches the joint stage.

Grazing and feeding restrictions - Non-lactating animals

Remove meat animals from treated areas 30 days prior to slaughter. There is no waiting period between treatment and grazing for non-lactating animals.

Grazing and feeding restrictions - Lactating animals

Do not graze lactating dairy animals within 7 days of treatment.

Dry hay and silage - Treated grasses may be harvested for dry hay or silage but do not harvest within 37 days of treatment.

Pasture and Rangeland Tank Mixes: BRASH[®] may be applied in tank mixes with one or more of the following herbicides: Ally[®], Amber[®], Albaugh Dicamba DMA Salt or Banvel[®], Clarity[®], Rave[™].

Pasture and Rangeland Restrictions:

Do not cut forage for hay within 7 days of application.

Pasture and Rangeland Restrictions, Postemergence:

For susceptible annual and biennial broadleaf weeds: Use 2.78 pints product/acre per application.

~~For moderately susceptible biennial and perennial broadleaf weeds: Use 2.78 – 5.6 pints product/acre per application.~~

For difficult to control weeds and woody plants: Use 5.6 pints product/acre per application.

Spot Treatment: Use 5.6 pints product/acre.

Maximum of two applications per year.

Maximum of 11 pints product/acre per year.

Minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for Worker Protection Standard are applicable.

SORGHUM

Rates and Timings

Apply 1 pint of BRASH[®] per acre to sorghum in the 3-5 leaf stage (4-8" tall). Apply BRASH[®] when weeds are small (less than 3" tall) for best performance.

Applications of BRASH[®] to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days. Sorghum growing under conditions of stress such as high moisture, low fertility, and abnormal temperature may be more sensitive to applications of BRASH[®].

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture, hay or silage, refer to the **Pasture and Rangeland** section of the **Food/Feed Crop-Specific Information** section of this label for livestock grazing and feeding restrictions.

Sorghum Restrictions:

Do not apply BRASH[®] to sorghum grown for seed production.

18/28

The preharvest interval (PHI) is 30 days.

Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.

Sorghum Restrictions, Postemergence:

Do not use surfactants or oils with postemergence applications of BRASH[®] on sorghum crops.

Do not use BRASH[®] if the potential for sorghum injury is not acceptable.

Limited to 1 application per crop cycle.

Maximum of 2.78 pints product/acre per application.

SORGHUM Tank Mixes: BRASH[®] may be applied in tank mixes with one or more of the following herbicides: Atrazine, Basagran[®], BROX[™] 2EC Herbicide or Buctril[®], Laddok[®] S-12, Paramount[®], Peak[®], or Permit[®].

SUGARCANE

Applications of BRASH[®] can be made any time after the weeds have emerged and are actively growing but prior to the close-in stage of sugarcane. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage. Application rates and timing are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

- For control of listed ANNUAL broadleaf weeds, apply 2 pints of BRASH[®] per treated acre.
- For suppression of listed PERENNIALS, apply 1 – 5.6 pints of BRASH[®] per treated acre.

Sugarcane Restrictions:

Do not harvest cane prior to crop maturity.

Do not apply more than 11 pints/acre per crop cycle.

Sugarcane Restrictions, Preemergence:

Limited to one application per crop cycle.

Maximum of 5.6 pints product/acre per application.

Sugarcane Restrictions, Postemergence:

Limited to one application per crop cycle.

Maximum of 5.6 pints product/acre per application.

SUGARCANE Tank Mixes: BRASH[®] may be tank mixed with one or more of the following herbicides: Asulox[®], Atrazine, Evik[®], Sencor[®], or Sinbar[®].

WHEAT

(Fall and Spring-seeded)

If small grains are grown for pasture or hay only, refer to the **Pastures, Rangeland and Grass (Hay, Silage)** section of this label.

Do not graze or harvest for livestock feed prior to crop maturity. Do not use BRASH[®] in wheat underseeded to legumes.

EARLY SEASON APPLICATIONS:

Apply 0.5-1 pint of BRASH[®] per acre to wheat unless using one of the wheat specific programs below.

Early season applications to spring-seeded wheat must be made after tillering and before wheat reaches the 6-leaf stage.

Early season applications to fall-seeded wheat must be made after tillering and prior to the jointing stage. Care should be taken in staging early developing wheat varieties such as TAM 107, Madison, or Wakefield to be certain that the application occurs prior to the jointing stage.

SPECIFIC USE PROGRAMS FOR FALL-SEEDED WHEAT ONLY:

Up to 1.33 pints of BRASH[®] per acre may be applied on fall-seeded wheat after the wheat begins to tiller for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. Periods of extending stresses such as cold and wet weather may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

PREHARVEST APPLICATIONS:

BRASH[®] can be used to control weeds that may interfere with harvest of wheat. Apply up to 1.4 pints of BRASH[®] per acre as a broadcast or spot treatment to annual broadleaf weeds when wheat is in the hard dough stage and the green color is gone from the nodes (joints) of the stem. Best results will be obtained if application can be made when weeds are actively growing but before weeds canopy.

Do not use preharvest-treated wheat for seed unless a germination test is performed on the seed with an acceptable result of 95% germination or better.

For control of additional broadleaf weeds or grasses, BRASH[®] may be tank mixed with other herbicides such as Ally[®], Gly Star[™] Plus or Roundup[®] Ultra that are registered for preharvest use in wheat.

Preharvest use of BRASH[®] is not registered for use in California.

Wheat Restrictions:

Do not graze or harvest for livestock feed prior to crop maturity.

Do not use Brash[®] in wheat underseeded with legumes.

The preharvest interval (PHI) is 14 days.

Limited to 4.9 pints product/acre per crop cycle.

Wheat Restrictions, Postemergence:

Limited to one postemergence application per crop cycle.

Maximum of 3.5 pints product per application.

Wheat Restrictions, Preharvest:

Limited to one postemergence application per crop cycle.

Maximum of 1.4 pints product per application.

WHEAT Tank Mixes

For control of grasses or additional broadleaf weeds, BRASH® may be tank mixed with the herbicides listed in the table below. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

Tank Mix Partner	Rate per Acre
Aim™	0.3 ounce
Ally®	0.05-0.1 ounce ¹
Amber®	0.14-0.28 ounce ¹
BROX™-M Herbicide or Bronate®	0.75-1.5 pints
BROX™ 2EC Herbicide or Buctril®	1-1.5 pints
Canvas®	0.2-0.4 ounce ¹
Curtail™	2 - 2.67 pints
Dakota®	16 fluid ounces
Express®	0.083-0.167 ounce ¹
Finesse®	0.167-0.33 ounce ¹
Glean®	0.167 ounce
Harmony® Extra	0.167-0.33 ounce ¹
Karmex® ³	0.5-1.5 pounds
2,4-D amine	4-20 fluid ounces ⁴
Sencor®, (Metribuzin ³)	0.25-0.375 pound a.i.
Peak® ¹	0.25-0.38 ounce
Stinger™	4 – 5.33 fluid ounces
Tiller® ²	1-1.7 pints

¹Do not use low rates of sulfonylurea herbicides such as Ally®, Amber®, Canvas®, Express®, Finesse®, Glean®, Harmony® Extra, and Peak® on more mature weeds or on dense vegetative growth.

²Do not use BRASH® as a tank mix treatment with Dakota® or Tiller® in Durum wheat. Do not tank mix Tiller® if wild oat is the target weed.

³Tank mixes with Karmex® and metribuzin are for use in fall-seeded wheat only.

⁴BRASH® contains 0.36 pounds a.e. of 2,4-D per pint. When tank mixing with 2,4-D, do not exceed a combined total of 1.0 pound a.e. per acre of 2,4-D and do not exceed 0.5 pound a.e. of 2,4-D unless injury to wheat is acceptable.

BETWEEN CROP APPLICATIONS, CONSERVATION RESERVE PROGRAMS, GENERAL FARMSTEAD AND FALLOW SYSTEMS

These uses are considered Food/Feed Crops when harvested, grazed or foraged. Refer to the **ADDITIVES** section for information on adjuvant restrictions and the **NON-FOOD/FEED USE (LAND NOT HARVESTED, GRAZED OR FORAGED) – SPECIFIC INFORMATION** section for specific use directions.

Restrictions:

Plant only labeled crops within 29 days following application.

Limited to 2 applications per year.

Maximum of 5.6 pints product/acre per application.

Minimum of 30 days between applications.

NON-FOOD/FEED USE (LAND NOT HARVESTED, GRAZED OR FORAGED) – SPECIFIC INFORMATION

Between Crop Applications

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL:

BRASH[®] can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply BRASH[®] as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

Refer to the **Crop Rotational Restrictions** and **General Restrictions and Limitations** for the recommended interval between application and planting to prevent crop injury.

Rates and Timings:

Apply 0.5-5.6 pints of BRASH[®] per acre. Refer to the **ANNUAL WEEDS** and **BIENNIAL AND PERENNIAL WEEDS Application Rate and Timing** tables to determine the use rates for specific targeted weed species. Retreatments may be made as needed; however, do not exceed a total of 8 pints of BRASH[®] per treated acre during a growing season. For best performance, apply BRASH[®] when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if BRASH[®] is applied when the majority of weeds have at least 4-6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for BRASH[®]. For seedling control, a follow-up program or other cultural practices could be instituted.

Fallowland (crop stubble on idle land, or postharvest to crops, or between crops) Restrictions:

Plant only labeled crops within 29 days following application.

Limited to 2 applications per year.

Maximum of 5.6 pints product/acre per application.

Minimum of 30 days between application.

Between Crop Tank Mixes

Apply 0.5-2 pints of BRASH[®] per acre in tank mixes with one or more of the following herbicides for control of annual weeds, or 2-8 pints of BRASH[®] per acre for control of biennial and perennial weeds:

Aim [™]	Glyphosate
Ally [®]	Gramoxone [®] Extra
Amber [®]	Kerb [™]
Atrazine	Landmaster [®] BW
Curtail [™]	Paramount [®]
Cyclone [®]	Sencor [®]
Distinct [®]	Tordon [™] 22K
Fallow Star [™]	Touchdown [®]
Fallowmaster [®]	2,4-D
Finesse [®]	

CONSERVATION RESERVE PROGRAMS AND GENERAL FARMSTEAD

BRASH[®] is recommended for use for Conservation Reserve Programs, general farmstead (non-cropland only), weed and brush control, or use in State Recognized Noxious Weed areas (non-cropland areas).

Refer to the **ANNUAL WEEDS** and **BIENNIAL AND PERRENIAL WEEDS Application Rate and Timing** tables for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

Rates above 4 pints of BRASH[®] per acre are for spot treatments only. Do not exceed a total of 8 pints of BRASH[®] per treated acre during a growing season.

Grasses in Conservation Reserve Program Areas

Annual Broadleaf Weeds – Apply when weeds are actively growing. Use higher rates on older weeds. Excessive injury may result if applied to young grasses with fewer than 6 leaves or prior to grasses being well established.

Biennial and Perennial Broadleaf Weeds – BRASH[®] may be used to suppress or control biennial and perennial broadleaf weeds in established grasses. Apply to actively growing weeds. Treat biennial weeds when they are in the seedling to rosette stage and before flower stalks become apparent. Treat perennial weeds in the bud to bloom stage.

Grasses in Conservation Reserve Program Areas Restrictions:

The preharvest interval (PHI) is 7 days (cut forage for hay).

Postemergence:

Limited to 2 applications per year.

Maximum of 5.6 pints product/acre per application.

Minimum of 30 days between application.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Farmstead and Fencerow Treatment

Application Instructions

BRASH[®] may be applied using water or oil and water emulsions in spot application to control undesirable vegetation using handgun or similar types of application equipment. In addition to weed species listed in the **ANNUAL WEEDS** and **BIENNIAL AND PERRENIAL WEEDS Application Rate and Timing** tables, these treatments may be used to control or suppress woody plant species listed below.

The following list of trees and vines can be controlled on farmsteads and fencerows as foliar, basal, or cut surface treatments:

Alder	Dogwood	Kudzu	Rose, McCartney
Ash	Elm	Locust, Black	Rose, Multiflora
Aspen	Grape	Maple	Sagebrush, Fringe
Basswood	Greenbriar	Mesquite	Sassafras
Beech	Hawthorn (Thornapple)	Oak	Spruce
Blackberry	Hemlock	Oak, Poison	Sumac
Blackgum	Hickory	Olive, Russian	Sweetgum

Cedar	Honeylocust	Persimmon, Eastern	Sycamore
Cherry	Honeysuckle	Pine	Tarbrush
Chinquapin	Hornbeam	Plum, Sand (Wild Plum)	Willow
Cottonwood	Huckleberry	Poplar	Witchhazel
Creosotebush	Huisache	Rabbitbrush	Yaupon
Dewberry	Ivy, Poison	Redcedar, Eastern	Yucca

To prepare oil and water emulsions, mix in the order and proportions indicated below. The solution should remain milky colored without an oily layer on top when under agitation. If an oily layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

Do not exceed 40 gallons of spray solution per treated acre per application. Forty gallons of spray solution contains 1.0 lb a.e. dicamba and 2.87 lb a.e. 2,4-D. Spray individual plants to wet. Do not allow this spray mixture to come into contact with desirable vegetation.

To control brush, briars, and weeds along fencerows surrounding pasture and ranch lands, and fallow fields, use a tank mix of 2.5% BRASH[®], 87.5% water, 10% diesel fuel, and sufficient emulsifier (to mix the diesel and emulsifier). The diesel oil in this tank mix will damage or kill desirable grasses and should not be used in pastures or where damage to desirable species cannot be tolerated.

1. **Water:** Begin by agitating a thoroughly clean sprayer tank with the desired quantity of clean water. Maintain constant agitation during complete mixing procedure.
2. **Emulsifier:** Add 0.5% volume to volume
3. **BRASH[®]:** Add 2.5 gallons per 100 gallons of total intended solution.
4. **Diesel Oil:** Add 10 gallons per 100 gallons of total intended solution.

Maintain constant agitation during application. Under good agitation, the spray solution should be milky white with no oily layer on top. If an oily layer forms, increase the amount of emulsifier or change to a more effective emulsifier.

CRP, Farmstead and Fencerow Treatment Restrictions, Postemergence – annual and perennial weeds:

Limited to 2 applications per year.

Maximum of 5.6 pints product/acre per application.

Minimum of 30 days between application.

CRP, Farmstead and Fencerow Treatment Restrictions, Postemergence – woody plants:

Limited to 1 application per year.

Maximum of 11 pints/acre per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research.

For Spraying Foliar Applications

1. Spray when leaves have reached full size but have not hardened due to drought or maturity.
2. Spray individual plants to wet with handgun.
3. For larger stems (up to 3" in diameter) and hard to control species, direct spray stream to base of stems to wet the stem at soil surface in addition to wetting the foliage.
4. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

For Dormant Basal Applications

1. Increase diesel content to 15% or 15 gallons/100 gallons of total mixture.
2. Spray in late winter and early spring before plants break dormancy.
3. Spray the bottom 24" of stem to wet on all sides.
4. For larger stems (up to 3" in diameter) and hard to kill species, direct spray solution to base of stems to wet the soil at the stem/soil junction in addition to wetting the stem.
5. Do not apply under drip line of desirable trees or adjacent to desirable vegetation.

Cut Surface Treatments

BRASH® may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. Use BRASH® in an undiluted state.

FRILL OR GIRDLE TREATMENTS: Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with BRASH®.

STUMP TREATMENTS: Spray or paint freshly cut surface with BRASH®. The cambium layer (the layer adjacent to the bark) should be thoroughly wet. Treat stumps within 6 hours after cutting.

PREPLANT APPLICATION DIRECTIONS FOR BROADLEAF CONTROL IN CROPLAND ROTATED TO WHEAT (POST-HARVEST / FALLOW / STUBBLE / SET-ASIDE)

IMPORTANT

Before using this product, read and carefully observe all applicable directions, restrictions, and precautionary statements in this booklet and on the container label.

WEEDS CONTROLLED

BRASH®, when applied at the specified rates, will control the ANNUAL and BIENNIAL weeds and suppress the PERENNIAL weeds listed below.

ANNUALS		
Buckwheat, Wild	Mustards	Salsify, Western
Cockle, Cow	Nightshade, Black	Smartweed, Pennsylvania
Cocklebur, Common	Pigweed, Redroot (Carelessweed)	Sowthistle, Annual
Knotweed	Pigweed, Rough	Sunflower
Kochia	Purslane, Common	Tansymustard
Lambsquarters, Common	Ragweed, Common	Thistle, Russian
Mallow, Common	Sage, Lanceleaf	Velvetleaf
BIENNIALS		
Carrot, Wild	Starthistle, Yellow	Thistle, Musk
Ragwort, Tansy	Thistle, Bull	Thistle, Plumeless
PERENNIALS		
Bindweed, Field	Dock, Curly	Thistle, Canada

RATES AND TIMINGS

Application may be made to fallow land, wheat stubble or land to be rotated to wheat. Application should be made to emerged and actively growing weeds. Use higher rate when treating dense vegetative growth. Avoid disturbing treated areas for seven days following application.

Wheat injury may occur if the interval between application and planting is less than 10 days for each pint per acre of BRASH® used. Exclude days when ground is frozen.

Weed Type & Stage	Broadcast Rate per Treated Acre Amount
Annual	
Small, actively growing (less than 4 inches)	1.0 – 1.5 pints
Established weed growth (greater than 4 inches)	1.5 – 3.0 pints
Biennial	
Rosette diameter (3 inches or less)	1.5 – 2.0 pints
(3 inches or more)	2.0 – 4.0 pints
Greater than 4 inches, tillering bolted or flowering	4.0 pints
Perennial	
Suppression or top growth control	2.0 – 4.0 pints
Seasonal Control	4.0 – 5.6 pints

Add 0.5% v/v of an agriculturally approved surfactant to BRASH[®] when used alone or in a tank mix. The addition of a surfactant will enhance spray coverage and the herbicide's penetration of weed foliage.

Retreatments may be made as needed; however, do not exceed a total of 11 pints of BRASH[®] per treated acre.

Cropland Rotated to Wheat (Post-Harvest / Fallow / Stubble / Set-Aside) Restrictions:

Plant only labeled crops within 29 days following application.

Limited to 2 applications per year.

Maximum of 5.6 pints product/acre per application.

Minimum of 30 days between applications.

TANK MIX TREATMENTS

BRASH[®] may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic or other restrictions. Add 0.5% v/v of an agriculturally approved surfactant to all tank mixes.

Herbicide	Rate per Treated Acre (lb ai)
Atrazine	1/2 to 3.0
Chlorsulfuron (Glean [®])	0.016 to 0.024
Glyphosate (Gly Star [™] Original or Roundup [®])	1/4 to 2.0
Metribuzin	1/3 to 3/4
Paraquat	1/2 to 1.0

**RIGHTS-OF-WAY
(Roadways, Utility, Railroad, Highway, Pipeline)**

When used as directed, BRASH[®] will control or suppress many herbaceous broadleaf weeds (annual, biennial and perennial) as well as many unwanted woody plant and vine species. Regardless of the species to be controlled, spray volumes should be high enough to allow for good spray coverage. Make applications when weeds and brush are actively growing.

Refer to the **ANNUAL WEEDS** and **BIENNIAL AND PERENNIAL WEEDS Application Rate and Timing** tables for rate selection based on targeted weed or brush species. Some weed species will require tank mixes for adequate control.

The addition of surfactants can increase control. Biennials are best controlled in the rosette stage. Regrowth may occur in resistant species.

HERBACEOUS BROADLEAF WEED CONTROL: Apply 2 to 5 ½ pints of BRASH® in 20 to 100 gallons of water per treated acre. When using low volume application equipment, 3 to 20 gallons of water per acre is acceptable. 2 to 4 pints per acre of BRASH® is recommended for annuals, 3 to 5 pints per acre for biennials and 5 ½ pints per acre for established perennials.

BRUSH AND VINE CONTROL – High Volume Foliar Spot Applications: Mix 6 to 8 pints of BRASH® in enough water to make 100 gallons of spray mix. When using low-volume application equipment, 3 to 20 gallons of water per acre is acceptable. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 8 pints of product per treated acre. Direct the spray to treat all foliage, stems, and root collars to wet.

BRUSH AND VINE CONTROL – Broadcast Applications with Ground Equipment: Apply 6 to 8 pints of BRASH® in 20 to 100 gallons of water per treated acre. When using low-volume application equipment, 3 to 20 gallons of water per acre is acceptable. Spray volume applied will depend on the size and density of the brush to be treated, but do not apply more than 8 pints of product per treated acre. Spray all foliage, stems, and root collars to wet.

AERIAL APPLICATIONS: Aerial applications may be made to control either herbaceous or woody plants. Apply 1 to 3 quarts of BRASH® for herbaceous weeds or 6 to 8 pints for woody brush and vines in 5 to 40 gallons of water per acre. Coverage is important, so increase spray volume when treating dense stands of brush or weeds. Do not apply more than 8 pints of product per treated acre.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, AND OTHER RESTRICTIONS. For broader spectrum control, BRASH® may be tank mixed with one or more of the following herbicides for non-cropland use (e.g. railroad, highway, pipeline, etc.).

Amitrole	Diquat	Maleic hydrazide	Simazine
Asulam	Diuron	Mefluidide	Sulfometruon methyl
Atritol	Fenac	Metsulfuron methyl	Sulfosate
Bromacil	Fosamine ammonium	MSMA	Tebuthiuron
Clorflurecol	Glyphosate	Norflurazon	Triclopyr
Chlorsulfuon	Glufosinate	Paraquat	2,4-D
Clopyralid	Hexazinone	Pendimethalin	2,4-DP
Dalapon	Imazapyr	Picloram	
Dicamba	Imazameth	Prodiamine	

Due to variations in formulated products and water supplies, a compatibility test is recommended prior to actual tank mixing.

All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

RIGHTS-OF-WAY RESTRICTIONS:

- Postemergence (annual & perennial weeds): Do not make more than 2 applications per year.

- Postemergence (annual & perennial weeds): Do not apply more than 5 ½ pints per acre per application.
- Postemergence (annual & perennial weeds): Minimum spray interval between applications is 30 days.
- Postemergence (woody plants): Do not make more than 1 application per year.
- Postemergence (woody plants): Do not apply more than 8 pints per acre per application.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

BRASH[®] contains 0.36 pounds a.e. of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of a.e. per acre per year.

BRASH[®] contains 0.125 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

REGISTERED TRADEMARKS

BROX[™], Fallow Star[™] and Gly Star[™] are trademarks of Albaugh, Inc.
Basagran[®] and Laddok[®] are registered trademarks of BASF Ag Products.

Banvel[®], Clarity[®], Distinct[®], Fallowmaster[®] and Paramount[®] are registered trademarks of BASF Corporation.

Asulox[®], Bronate[®], Buctril[®], Dakota[®], Sencor[®] and Tiller[®] are registered trademarks of Bayer CropScience.

Curtail[™], Kerb[™], Stinger[™] and Tordon[™] are trademarks of Dow AgroSciences, LLC.

Ally[®], Canvas[®], Express[®], Finesse[®], Glean[®], Harmony[®] Extra, Karmex[®], and Sinbar[®] are registered trademarks of E.I. DuPont de Nemours & Company.

Aim[™] is a trademark of FMC Corporation.

Landmaster[®], Permit[®] and Roundup[®] are registered trademarks of the Monsanto Company.

Amber[®], Cyclone[®], Evik[®], Gramoxone[®], Peak[®], Rave[®] and Touchdown[®] are registered trademarks of Syngenta Crop Protection, Inc.

AgriSolutions[™] is a trademark and Brash[®] and Sterling[®] are registered trademarks of Winfield Solutions, LLC.