

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 30, 2014

Danielle Erding
Registration Specialist
Winfield Solutions, LLC
P.O. box 64589
MS 5705
St. Paul, MN 55164

Subject: Label Notification per PRNs 98-10 and 2001-5 – Adding Mode of Action code to page 1 (Group 4 A)
Product Name: Advise 2FL
EPA Registration Number: 1381-205
Application Date: 7/16/2014
Decision Number: 493592

Dear Ms. Erding:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notices (PRNs) 98-10 and 2001-5 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under these PRNs and finds that the action requested falls within the scope of PRNs 98-10 and 2001-5.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

If you have any questions, you may contact Autumn Metzger at 703-305-5314 or via email at metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Venus Eagle".

Venus Eagle, Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)
Office of Pesticide Programs

GROUP

4A

INSECTICIDE

ADVISE[®] 2FL

For uses in pest management and suppression of listed insects that may vector diseases and maintenance of plant health.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 21.4%

OTHER INGREDIENTS: 78.6%

TOTAL 100.0%

Contains 2 pounds of imidacloprid per gallon.

SHAKE WELL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTIFICATION

Jul 30, 2014

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
In case of medical emergency call toll free at 1-877-424-7452. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
Note to Physician: No specific antidote is available. Treat the patient symptomatically.	

SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.

EPA Reg. No. 1381-205

EPA Est. No. 42750-MO-01

Distributed By:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

NET CONTENTS: _____ gallon(s)
LOT NO. _____

1/1216/3 1/0716/4

WINFIELD[™]

AgriSOLUTIONS[™]

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

TAKE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar. Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>. Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See individual crops for specific pollinator protection application restrictions. If none exist under the specific crop, for foliar applications, follow these application directions for crops that are contracted to have pollinator services or for food/feed & commercially grown ornamentals that are attractive to pollinators:

1. FOR CROPS UNDER CONTRACTED POLLINATION SERVICES

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless the following condition has been met.

If an application must be made when managed bees are at the treatment site, the beekeeper providing the pollination services must be notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

2. FOR FOOD CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT





FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton, and
- Shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Use label language appropriate for container size and type.

Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident,
call CHEMTREC 1-800-424-9300.

Do not apply this product through any type of irrigation system unless it is specified in the crop-specific sections of this label.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions.

Avoiding spray drift is the responsibility of the applicator.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150 – 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions during Temperature Inversions:

Do not make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

For Aerial Applications:

Mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Airblast (Air Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. Follow specific drift practices:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

No-Spray Zone Requirements for Foliar Applications

Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

Mixing and Loading Requirements

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintained a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes or field drains.

No-Spray Zone Requirements for Soil Applications

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Runoff Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. Employ the best management practices for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

Endangered Species Notice

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

Resistance Management

Some insects may develop resistance to imidacloprid after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

This product contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

The Active ingredient in this product belongs to the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to this product. In order to maintain susceptibility to this class of chemistry in insect species with the high resistance development potential, it is recommended that for each crop season: 1) only a single, soil application of this product be made; 2) foliar applications of products from this same class not be made following a long residual, soil application of this product, or other neonicotinoid products.

Other Group 4A neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Clutch, Couraze, Intruder, Leverage, Pasada, Provado, Trimaz Pro, and Venom.

Other Group 4A neonicotinoid products used as soil treatments include: Admire Pro, Alias, Couraze, Platinum and Venom.

Contact your local extension specialists, certified crop advisor and /or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irc-online.org/>.

SOIL APPLICATION DIRECTIONS

Direct application of this product into the seed or root-zone of crop. Failure to place this product into root-zone may result in loss of control or delay in onset of activity. This product may be applied with ground or chemigation application. Make broadcast foliar applications to seedling flats or trays, or where product is intended to be washed from foliage to soil prior to drying on foliage.

Optimum activity of this product results from applications to the root-zone of plants to be protected. The earlier this product is available to a developing plant, the earlier the protection begins. This product is continuously taken into the roots over a long period of time and the systemic nature of this product allows movement from roots through the xylem tissue to all vegetative parts of the plant. This results in extended residual activity of this product, the control of insects and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of Advise 2FL applied affects the length of the plant protection period. Use higher rates when infestations occur later in crop development, or where pest pressure is continuous. This product will generally not control insects infesting flowers, blooms or fruit. Additional crop protection may be required for insects feeding in, or on these plant parts and for insects not listed in the crop-specific, pests controlled sections of this label. Additional, specific Advise 2FL directions are also provided in the crop-specific sections of this label.

Suppression, or less than complete control of certain diseases and insect pests including reduced feeding may also result from ADVISE 2FL applications. Control of these pests/diseases may require supplemental crop protection measures.

ADVISE 2FL use on crops grown for production of true seed intended for private or commercial planting is not permitted unless allowed under State specific 24(c) labeling. Contact your local Cooperative Extension Service, PCAs, consultants or local Winfield Solutions representative for application on these type crops.

Pre-mix ADVISE 2FL with water or other appropriate diluents prior to application. Maintain constant agitation to avoid settling.

Restriction:

Do not apply more than 0.50 lbs active ingredient per acre, per year, regardless of formulation or method of application unless specified within a crop specific section for a given crop. Do not apply with aerial equipment.

FOLIAR APPLICATION DIRECTIONS

Apply ADVISE 2FL as a directed or broadcast foliar spray. Thorough coverage of foliage without runoff is required for maximum insecticidal efficacy. Use of adequate spray volumes and correctly calibrated application equipment is critical. Use of a spray adjuvant may enhance thorough coverage. Lack of adequate coverage and retention of ADVISE 2FL on foliage and fruit can delay or lessen insect control. ADVISE 2FL may be applied with ground or aerial application equipment that has been properly calibrated.

Minimum rate spray volumes (unless otherwise specified on crop sections) are:

- 10 gallons/Acre by ground application
- 5 gallons/Acre through aerial equipment.

ADVISE 2FL may also be applied by overhead chemigation (see CHEMIGATION DIRECTIONS FOR USE

section below) if allowed in crop specific application section.

ADVISE 2FL application to crops grown for production of true seed intended for private or commercial planting may be allowed under State specific 24(c) labeling. Additional information on ADVISE 2FL uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants or local Winfield Solutions, LLC representatives.

Restrictions:

Do not exceed application of more than 0.5 lbs. active ingredient per acre, per year, regardless of formulation or method of application unless specified within a crop specific section for a given crop.

Do not apply ADVISE 2FL in enclosed structures such as greenhouses or planthouses.

MIXING INSTRUCTIONS FOR SOIL AND FOLIAR APPLICATIONS

To prepare the application mixture, add a portion of the required amount of water to the tank and with agitation add this product. Complete filling tank with the balance of water needed. Maintain sufficient agitation during both mixing and application. This product may also be used with other pesticides and/or fertilizer solutions. Please see Compatibility Information below. When tank mixture of this product and other pesticides are involved, prepare the tank mixture as specified above and follow suggested mixing order below.

Mixing Order: When pesticide mixtures are needed, add wettable powders first, this product and other flowable (suspension concentrate) products second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Compatibility Information: Unless the applicator has prior knowledge of the compatibility of the intended tank mixture, Winfield Solutions recommends a small scale test by adding proportionate amounts of each ingredient in the appropriate order, to a clear pint or quart sized jar. Cap and shake for 5 minutes, then let set for 5 minutes. Any visual indication of poor mixing or formation of precipitates that cannot be easily re-dispersed indicates incompatibility and the mixture must not be used.

USE IN CHEMIGATION SYSTEMS

Types of Irrigation for Soil Systems: Chemigation applications of ADVISE 2FL may only be made to crops through chemigation systems as specified in crop-specific sections and only through low-pressure irrigation systems unless specifically indicated for a given crop. Do not apply ADVISE 2FL through any other type of irrigation system.

Types of Irrigation Systems for Foliar Application:

Chemigation applications of ADVISE 2FL may be made to crops through overhead sprinkler chemigation systems if specified in crop-specific sections. Do not apply ADVISE 2FL through any other type of irrigation system. Make ADVISE 2FL foliar chemigation applications as concentrated as possible. Retention of ADVISE 2FL on target site of insect infestation is necessary for optimum activity. Do not use water volumes exceeding 0.10 inches/acre for chemigation applications.

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, there is a 12-month plant-back interval.

PLANTBACK INTERVAL	COMMENT
Immediate Plant-back:	Any crop listed on this label plus the following crops not on this label: barley, canola, cardoon, corn (field, sweet and pop), rapeseed, sorghum, sugar beet, and wheat.
30-Day Plant-back:	Cereals (including buckwheat, millet, oats, rice, rye and triticale), soybeans, safflower
12-Month Plant-back:	All other crops

*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

FIELD CROPS

COTTON (Soil Application)

To control Aphids, Plant bugs, Thrips, Whiteflies apply 1.3 fluid ounces per 1,000 row feet. Equivalent to 17.0 to 21.1 fluid ounces per acre depending on row spacing.

Apply label rate of ADVISE 2FL in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting;
3. Chemigation into root-zone through low-pressure drip or trickle irrigation.

Restrictions for Cotton (Soil Application):

- Maximum ADVISE 2FL allowed per year: 21.1 fluid ounces/Acre (0.33 lb ai/Acre)
- Apply no more than 0.5 lb active ingredient of imidacloprid per acre per year including seed treatment, soil and foliar uses.
- Do not apply more than a total of 6 applications of the active ingredient per season.
- Do not graze treated fields after any application of ADVISE 2FL.

Please see Resistance Management section of this label.

COTTON (Foliar Application)

Apply as foliar spray at specified rate per acre when insect pressure reaches economic threshold. Uniform coverage is required to achieve best control and a spray adjuvant may help improve coverage. Two applications may be required to achieve control when initial insect populations are high. Base retreatment on field scout reports. ADVISE 2FL may be tank mixed with other labeled insecticides to increase control or control pests not controlled by imidacloprid. Apply only through properly calibrated ground, aerial or chemigation application equipment insuring thorough coverage.

For control of Aphids, Flea hoppers, Bandedwinged whitefly, Plant bugs (east of Rocky Mountains), Green stink bug, Southern green stink bug, Bollworm/Budworm (ovicidal effect) apply 2.0 to 4.0 fluid ounces per acre (3.0 fluid ounces: CA only) and suppression of Lygus bug (west of Rocky Mountains) and Whiteflies (other than bandedwinged whitefly) apply 3.0 to 4.0 fluid ounces per acre, (3.0 fluid ounces per acre: CA only) as a broadcast or directed spray to infested area. Apply only through properly calibrated ground, aerial or chemigation application equipment insuring thorough coverage.

Tank Mix Information

For early season control of Thrips apply 2.0 to 3.0 fluid ounces of ADVISE 2FL per acre + 1.6 to 3.2 fluid ounces of Bidrin[®] 8* per acre as a broadcast or directed foliar spray to infested area. For mid to late season control of plant bugs, stink bugs (including Brown stink bug), Grasshoppers, Saltmarsh caterpillar and Cotton leafperforator apply 2.0 to 3.0 fluid ounces of ADVISE 2FL per acre + 4.0 to 8.0 fluid ounces of Bidrin[®] 8* per acre as a broadcast spray of directed foliar spray to infested area.

Restrictions for Cotton (Foliar Application):

- Pre-Harvest Interval (PHI): 14 days
- Minimum interval between applications : 7 days
- Maximum ADVISE 2FL allowed per year: 20.0 fluid ounces/Acre (0.31 lb AI/A); For CA: 17.6 fluid ounces/Acre (0.28 lb. AI/A)
- Maximum number of ADVISE 2FL applications per crop season: 6
- Do not graze treated fields after any application of ADVISE 2FL.

* Refer to the Bidrin[®] 8 product label for specific use rates; observe all restrictions and precautions that appear on the label.

POTATO (Soil Application)

To control Aphids, Colorado potato beetle, Flea beetles, Potato Psyllids apply 0.9 to 1.3 fluid ounces per 1,000 row feet, which is equivalent to 13.0 to 20.0 fluid ounces per acre depending on row spacing.

To suppress wireworms or in response to symptoms of Potato leaf roll virus (PLRV), Potato yellows, or Net necrosis (PLRV) apply 0.9 to 1.3 fluid ounces per 1,000 row feet, which is equivalent to 13.0 to 20.0 fluid ounces per acre depending on row spacing.

SOIL APPLICATION

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. In-furrow spray during planting directed on the seed pieces or seed potatoes;
2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil;
3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil;
4. In the bedding operation (7 days or less) before planting, apply a narrow band directly below the eventual seed row. To be effective applications must be placed below soil-surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of ADVISE 2FL may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered.

Restrictions for Potato (Soil Application):

- Maximum ADVISE 2FL allowed per year: 20.0 fluid ounces/acre (0.31 lb AI/Acre.)

POTATO (Foliar Application)

For control of Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Psyllids apply 3.0 fluid ounces per acre as a broadcast or directed spray to infested area. Apply only through properly calibrated ground, aerial or chemigation application equipment insuring thorough coverage.

Restrictions for Potatoes (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum Interval between applications: 7 days
- Maximum ADVISE 2FL allowed per crop season: 12.1 fluid ounces/Acre (0.19 lb. AI/A)

POTATO (Seed Piece Treatment)

To control Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Potato Psyllids, and Wireworms apply 0.4 to 0.8 fluid ounces per 100 lbs. seed. Equivalent to 8.0 to 16.0 fluid ounces per acre depending on row spacing. (Based on a seeding rate of 2000 lbs/acre)

In response to symptoms of Potato leaf roll virus (PLRV), Potato yellows, or Net necrosis (PLRV) apply 0.8 fluid ounces per 1,000 row feet, which is equivalent to 16.0 fluid ounces per acre depending on row spacing.

Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part ADVISE 2FL. Maintain agitation during application. Fungicidal dust treatments may be applied after ADVISE 2FL application. Make applications only in areas with adequate ventilation. Plant seed-pieces as soon as possible after treating. Do not expose seed-pieces treated with ADVISE 2FL to sunlight.

Restrictions for Potato Seed Piece Treatment:

- Maximum ADVISE 2FL allowed per crop season: 20.0 fluid ounces/Acre (0.31 lb AI/Acre)
- Do not use treated seed-pieces for food, feed, or fodder.
- Do not apply any subsequent application of ADVISE 2FL or any other imidacloprid based product (in-furrow), following an ADVISE 2FL seed-piece treatment.

TOBACCO (Soil Application)

To control Aphids and Flea beetles apply 1.0 fluid ounces per 1,000 plants as seedling tray drench or 1.4 fluid ounces per 1,000 plants as an in-furrow or transplant-water.

To control Mole Crickets, Whiteflies, Wireworms apply 1.4 to 2.8 fluid ounce per 1,000 plants as seedling tray

drench or 1.8 to 2.8 fluid ounces per 1000 plants as an in-furrow or transplant water.

To suppress cutworms or in response to symptoms of Tomato spotted wilt virus (TSWV) apply 1.4 to 2.8 fluid ounce per 1,000 plants as seedling tray drench or 1.8 to 2.8 fluid ounces per 1000 plants as an in-furrow or transplant water.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting. Follow as soon as possible with overhead irrigation to wash ADVISE 2FL from foliage into potting media. Failure to wash ADVISE 2FL from foliage may result in a reduction in pest control. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.
2. In-furrow spray or transplant-water drench during setting.
3. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Proper drench applications to plants in trays with ADVISE 2FL are generally the best method of application. However, the specified rate of ADVISE 2FL may be applied as combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of ADVISE 2FL into the plant and a delay in control.

To control Aphids and Flea beetles apply 1.0 fluid ounce per 1,000 plants as seedling tray drench or 1.4 fluid ounces per 1000 plants as an in-furrow or transplant water.

Restrictions for Tobacco (Soil Application):

- Pre-Harvest Interval (PHI): 14 days
- Maximum ADVISE 2FL allowed per year is 32.0 fluid ounces/Acre (0.50 lb ai/acre)

TOBACCO (Foliar Application)

For control of Aphids apply 1.6 to 3.2 fluid ounces per acre as a broadcast or directed spray to infested area. Use higher rate when insect pressure is heavy.

For control of Flea beetles and Japanese beetle apply 3.2 fluid ounces per acre as a broadcast or directed spray to infested area.

Restrictions for Tobacco (Foliar Application):

- Pre-Harvest Interval (PHI): 14 days
- Minimum interval between applications : 7 days
- Maximum ADVISE 2FL allowed per year: 17.9 fluid ounces/Acre (0.28 lb. AI/A)

Apply only through properly calibrated ground, aerial or chemigation application equipment insuring thorough coverage.

VEGETABLE AND SMALL FRUIT CROPS

For Foliar Application uniform coverage is required to achieve best control and a spray adjuvant may help improve coverage. Two applications may be required to achieve control when initial insect populations are high. Base retreatment on field scout reports. ADVISE 2FL may be tank mixed with other labeled insecticides to increase control or control pests not controlled by imidacloprid.

Apply only through properly calibrated ground, aerial or chemigation application equipment insuring thorough coverage.

CUCURBIT VEGETABLES (Soil Application Only)

Includes: Chayote (fruit), Chinese waxgourd, (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourds (edible, includes hyotan, cucuzza, hechima, Chinese okra), Momordica spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of Cucumis melo including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash,

calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

To control Aphids, Cucumber beetles, Leafhoppers, Thrips (Foliar-feeding thrips only), Whiteflies apply 16.0 to 24.0 fluid ounces per acre.

For suppression in response to Bacterial wilt (as vectored by various cucumber beetles) and Leaf silvering resulting from whitefly feeding apply 16.0 to 24.0 fluid ounces per acre.

Apply specific dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
2. In-furrow spray directed on or below seed
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting
5. Post-seeding drench, transplant-water drench or hill drench
6. Subsurface side-dress on both sides of each row. ADVISE 2FL must be incorporated into root-zone

Restrictions for Cucurbit Vegetables (Soil Application):

- Preharvest Interval (PHI): 21 days
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)

CUCURBIT VEGETABLES (Planthouse Application)

To control Aphids and Whiteflies apply 0.1 fluid ounce per 1000 plants.

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash ADVISE 2FL from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash ADVISE 2FL from foliage may result in reduced pest control
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.

Important: Not all varieties of fruiting vegetables have been tested for tolerance to ADVISE 2FL applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

Restrictions for Cucurbit Vegetables (Planthouse Application):

- Maximum amount ADVISE 2FL applied in the planthouse: 0.1 fluid ounces (0.00156 pound AI)/1,000 plants, and maximum number of applications in planthouse: 1.
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

FRUITING VEGETABLES (Soil Application)

Includes: Eggplant, Ground Cherry, Okra, Pepper (Including Bell peppers, Chili peppers, Cooking peppers, Pimentos and Sweet peppers), Tomato, Pepinos, Tomatillo

To control Aphids, Colorado potato beetle, Flea beetles, Leafhoppers, Thrips (Foliar-feeding thrips only), Whiteflies apply 16.0 to 24.0 fluid ounces per acre. For Okra and Pepper apply 16.0 to 32.0 fluid ounces per acre.

For suppression in response to symptoms of Tomato mottle virus, Tomato spotted wilt virus, Tomato yellow leaf curl virus apply 16.0 to 24.0 fluid ounces per acre. For Okra and Pepper apply 16.0 to 32.0 fluid ounces per acre.

Apply specific dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
2. In-furrow spray directed on or below seed
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2 " with sufficient irrigation within 24 hours of application
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting
5. Post seeding drench, transplant-water drench, or hill drench.
6. Subsurface side-dress on both sides of each row. ADVISE 2FL must be incorporated into root-zone.

Restrictions for Fruiting Vegetables (Soil Application):

- Pre-Harvest Interval (PHI): 21 days
- Maximum ADVISE 2FL allowed on pepper and okra crops per season: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Maximum ADVISE 2FL allowed on other fruiting vegetable crops per season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

FRUITING VEGETABLES (Foliar Application)

Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet), Tomato, Pepinos, Tomatillo

For control of Aphids, Colorado potato beetle, Leafhoppers, Whiteflies apply 3.0 fluid ounces per acre.

For control of Pepper weevil (Pepper only) apply 5.0 fluid ounces per acre. Make applications prior to a damaging population becoming established.

Restrictions for Fruiting Vegetables (Foliar Application):

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 5 days
- Maximum ADVISE 2FL allowed per crop season: 15.2 fluid ounces/Acre (0.24 lb. AI/A)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling

Applications of ADVISE 2FL must be part of a full-season resistance management program that uses alternate applications products from multiple classes of chemistry and different modes of action. For pepper weevil, apply specified dosage of ADVISE 2FL by ground equipment only, timing applications prior to a damaging population becoming established. Good coverage of foliage and fruit is necessary for optimum control.

FRUITING VEGETABLES (Planthouse application)

To control Aphids and Whiteflies apply 0.1 fluid ounce per 1000 plants.

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash ADVISE 2FL from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash ADVISE 2FL from foliage may result in reduced pest control.
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.

Restrictions for Fruiting Vegetables (Planthouse Application):

- Maximum amount ADVISE 2FL applied in the planthouse: 0.1 fluid ounces (0.00156 lb AI)/1000 Plants
- Maximum number ADVISE 2FL applications in planthouse: 1

Important: Not all varieties of fruiting vegetables have been tested for tolerance to ADVISE 2FL applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

GLOBE ARTICHOKE (Foliar Application Only)

For control of Aphids and Leafhoppers apply 3.2 to 8.0 fluid ounces per acre. Use higher listed rates when pest pressure more severe.

Restrictions for Globe Artichoke:

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 14 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb. AI/A)

GREENHOUSE VEGETABLES (Soil Application Only)

(mature plants in production greenhouses)

Cucumber, Tomato Only

To control Aphids and Whiteflies apply 1.4 fluid ounces per 1000 plants.

Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Do not apply to immature plants since phytotoxicity may occur.

Make applications when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (*Orius* sp.) can occur when ADVISE 2FL is applied.

Many varieties of vegetables have been tested for tolerance to ADVISE 2FL and show good safety. However, certain varieties may show more sensitivity to ADVISE 2FL. Therefore, treatment of a few plants is recommended before treating the whole greenhouse.

Restrictions for Greenhouse Vegetables:

- Pre-Harvest Interval (PHI): 0 days
- Maximum number ADVISE 2FL applications per crop season: 1
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

HEAD and STEM BRASSICA VEGETABLES (Soil Application)

Includes: Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai Choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard spinach, Rape greens, Turnip tops (leaves)

AND

LEAFY GREEN VEGETABLES (Soil Application)

Including: amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

To control Aphids and Whiteflies apply 10.0 to 24.0 fluid ounces per acre (based on 36 inch row spacing).

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
2. In-furrow spray directed on or below seed
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting
5. Post-seeding drench, transplant-water drench or hill drench
6. Subsurface side-dress on both sides of each row. ADVISE 2FL must be incorporated into root-zone

Restrictions for Head and Stem Vegetables and Leafy Green Vegetables (Soil Application):

- Pre-Harvest Interval (PHI): 21 days
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

HEAD and STEM BRASSICA VEGETABLES (Foliar Application)

Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai Lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens

AND

LEAFY GREEN VEGETABLES (Foliar Application)

Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden). Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter). Raddicchio (red chicory). Spinach (Including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only. Applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

For control of Aphids, Flea beetles, Leafhoppers, Whiteflies apply 3.0 fluid ounces per acre.

Restrictions for Head and Stem Brassica Vegetables and Leafy Green Vegetables (Foliar Applications):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum ADVISE 2FL allowed per crop season: 15.2 fluid ounces/Acre (.24 lb. AI/A)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling

HERBS (Soil Application)

Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Loveage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

To control Aphids, Flea beetles, Leafhoppers, and Whiteflies apply 16.0 to 24.0 fluid ounces per acre.

For suppression in response to Thrips (foliage feeding thrips only) apply 16.0 – 24.0 fluid ounces per acre.

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Winfield Solutions, LLC strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.

Apply specific dosage of ADVISE 2FL in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. In-furrow spray or transplant-water drench during setting or transplanting;
3. Shanked-into or below eventual seed-line;
4. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Restrictions for Herbs (Soil Application):

- Pre-Harvest Interval (PHI): 14 days
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)

HERBS (Foliar Application)

Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Loveage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

To control Aphids, Flea beetles, Leafhoppers, and Whiteflies apply 2.8 fluid ounces per acre.

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Winfield Solutions, LLC strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.

Apply specific dosage of ADVISE 2FL through properly calibrated ground or aerial application equipment. Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage and control.

Restrictions for Herbs (foliar application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum ADVISE 2FL allowed per crop season: 8.4 fluid ounces/Acre (0.13 lb ai/acre)

LEAFY PETIOLE VEGETABLES (Soil Application Only)

Including: Cardoon, Celery, Chinese celery (fresh leaves and stalk only), Celtuce, Florence fennel including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard

To control Aphids, Leafhoppers, Whiteflies apply 10.0 to 24.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
2. In-furrow spray directed on or below seed
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting
5. Post-seeding drench, transplant-water drench or hill drench
6. Subsurface side-dress on both sides of each row. ADVISE 2FL must be incorporated into root-zone

Restrictions for Leafy Petiole Vegetables:

- Pre-Harvest Interval (PHI): 45 days
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

LEGUME VEGETABLES (except soybean, dry) (Soil Application)

Including: Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

Bean (Lupinus spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)
Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)
Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)
Other Beans and Peas (Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (Hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean)

To control Aphids, Leafhoppers, Thrips (foliar feeding only), Whiteflies apply 16.0 to 24.0 fluid ounces per acre.

For suppression In response to symptoms of Bean common mosaic virus (BCMV), Bean golden mosaic virus (BGMV), Beet curly top hybrigeminivirus (BCTV) apply 16.0 to 24.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment
2. In-furrow spray at planting directed on or below seed
3. In a narrow (2" or less) surface band over seed-line during planting incorporated into a depth of 1 to 1½ inch with sufficient irrigation within 24 hours following application
4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting
5. As a post-seeding drench, transplant drench, or hill drench.

Restrictions for Legume Vegetables (Soil Application):

- Preharvest Interval (PHI): 21 days
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

LEGUMES VEGETABLES (except soybean, dry) (Foliar Application)

Edible Podded and Succulent Shelled Pea/ and Bean and Dried Shelled Pea and Bean
Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)
Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)
Bean (*vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean. Southern pea, urd bean, yardlong bean)
Pea (*Pisum* spp. Includes dwarf pea, edible pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snappea)
Other Beans and Peas (Broad been (fava), chickpea (garbanzo bean), Guar, Jackbean. Lablab bean (hyacinth bean), lentil. Pigeon pea, soybean (immature seed), Sword bean)

For control of Aphids, Leafhoppers, Whiteflies apply 2.8 fluid ounces per acre.

Restrictions for Legume Vegetables (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 7 days
- Maximum ADVISE 2FL allowed per crop season: 8.3 fluid ounces/Acre (0.13 lb. AI/A)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

ROOT VEGETABLES (Soil Application)

Including: Beet (garden)¹, Carrot¹, Burdock (edible)¹, Celeriac¹, Chervil (turnip-rooted)¹, Chicory¹, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip¹, Radish¹, Oriental radish (daikon)¹, Rutabaga¹, Salsify (oyster plant), Salsify (black)¹, Salsify (Spanish), Skirret, Turnip¹

¹ Tops or greens from these crops may be utilized for food or feed.

To control Aphids, Flea beetles, Leafhoppers, Whiteflies apply 0.7 to 1.7 fluid ounces per 1000 row feet, which is equivalent to 10.0 to 24.0 fluid ounces per acre depending on row spacing.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting;
3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting.

Important: Rate applied affects the length of control. Use higher listed rates where infestations occur later in crop development, or where pest pressure is continuous. ADVISE 2FL rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. ADVISE 2FL treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

Restrictions for Root Vegetables:

- Pre-Harvest Interval (PHI): 21 days.
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lbs AI/Acre).
- Maximum ADVISE 2FL applications per crop season: 1.
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling

TUBEROUS AND CORM VEGETABLES (Soil Application)

Includes: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Casava (bitter and sweet)¹, Chayote (root), Chufa, Dasheen (taro)¹, Ginger, Leren, Sweet potato, Tanier (cocoyam)¹, Turmeric, Yam bean (jicama, manioc pea), Yam (true)¹
(For applications on potato see Field Crop section)

¹ Tops or greens from these crops may be utilized for food or feed.

To control Aphids, Flea beetles, Leafhoppers, Whiteflies apply 0.7 to 1.7 fluid ounces per 1000 row feet, which is equivalent to 10.0 to 24.0 fluid ounces per acre depending on row spacing.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. In-furrow spray (rate specified per 1000 row-feet) over planting material (hulis) or shanked-in 1 to 2 inches below hulis depth at planting;
2. Side-dress not more than 0.6 fluid ounces/1000 row-feet no later than 45 days after-planting. Observe same PHI as above.

Rate applied affects the length of control. Use higher listed rates where infestations occur later in crop development, or where pest pressure is heavy or continuous. ADVISE 2FL rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. ADVISE 2FL treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

Restrictions for Tuberous and Corm Vegetables (Soil Application):

- Pre-Harvest Interval (PHI) from planting application: 3 days (leaves); 125 days (corms).
- Maximum ADVISE 2FL allowed per crop season: 24.0 fluid ounces/Acre (0.38 lbs AI/Acre).
- Maximum ADVISE 2FL applications per crop season: 1.
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

ROOT, TUBEROUS and CORM VEGETABLES (Foliar Application)

Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden)¹, Burdock (edible)¹, Canna (edible, Queensland arrowroot), carrot¹, Cassava (bitter & sweet)¹, Celeriac¹, Chayote (root), Chervil (turnip-rooted)¹, Chickory¹, Chufa, Dasheen (taro), Ginger, Ginseng, Horseradish, Leren, Parsley (turnip-rooted), Parsnip¹, Radish¹, Oriental radish (diakon)¹, Rutabaga¹, Salsify (black), Salsify (oyster plant), Salsify (Spanish), Skirret, Sweetpotato¹, Tanier (cocoyam)¹, Turmeric, Turnip¹, Yam bean (jicama, manioc pea), Yam (true)¹

¹Tops or greens from these crops may be utilized for food or feed.

For applications on potato see Field Crops section

For control of Aphids, Flea beetles, Leafhoppers, Whiteflies apply 2.8 fluid ounces per acre.

Restrictions for Root, Tuberous and Corm Vegetables (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum ADVISE 2FL allowed per crop season: 2.8 fluid ounces/Acre (0.044 lb. AI/A) on radish; 8.3 fluid ounces/Acre (0.13 lb. AI/A) on other crops.
- Maximum ADVISE 2FL applications per crop season: 1 on radish; 3 on other crops
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

STRAWBERRY (Soil Application) (annual and perennial crops)

To control Aphids and Whiteflies apply 24.0 to 32.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods.

1. Chemigation into root-zone through low pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening
2. As a plant material or plant hole treatment just prior to, or during transplant.

The rate applied affects the length of control. Use higher listed rates where infestations may occur later in crop development or where pest pressure is continuous.

Restrictions for Strawberries (annual and perennial crops)

- Pre-Harvest Interval (PHI): 14 days
- Maximum ADVISE 2FL allowed per crop season: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.
- Do not use both application methods on the same crop in the same season.
- Do not apply immediately prior to bud opening or during bloom or when bees are foraging.



STRAWBERRY (Soil Application) (Post-harvest Use on Perennial Crops)

To control White grub complex (grubs of Asiatic garden beetle, European and Masked Chafer, Japanese beetle, Oriental beetle) apply 16.0 to 24.0 fluid ounces/acre.

Apply a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified dosage of ADVISE 2FL in one of the following methods:

1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre
2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed.
3. As a chemigation application with 600 to 1000 gallons of water followed by 0.10 to 0.25 inches irrigation.

Important: All soil-surface applications must be followed by 0.25 inches of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate ADVISE 2FL into egg-deposition zone may result in decreased activity of beetle grubs.

Restrictions for Strawberry Post-harvest use on Perennial Crops:

- Pre-Harvest Interval (PHI): 14 days
- Maximum ADVISE 2FL allowed pre-season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Do not use both application methods on the same crop in the same season.

STRAWBERRY (Foliar Application)

For control of Aphids, Spittlebugs, Whiteflies apply 3.0 fluid ounces per acre.

Restrictions for Strawberries (Foliar Applications):



- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 5 days
- Maximum ADVISE 2FL allowed per crop season: 9.1 fluid ounces/Acre (0.14 lb. AI/A)
- Do not apply during bloom or within 10 days prior to bloom or when bees are foraging.

SUGARBEET (Soil Application Only)

To control Aphids, Leafhoppers, Whiteflies, Flea beetles apply 6.0 to 12.0 fluid ounces per acre.

For suppression in response to symptoms of Western yellows/Beet curly top hybrigeminivirus (BCTV) apply 6.0 to 12.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in the following method:

1. Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of other pests listed.

Restrictions for Sugarbeets:

- Maximum ADVISE 2FL allowed per year: 12.0 fluid ounces/Acre (0.18 lb ai/acre)
- Maximum imidacloprid allowed per season: 0.18 lb ai/acre (from any formulation) on any row spacing
- For use only in California
- Not for use on crops grown for seed unless allowed by state-specific 24(c) labeling.

CONVERSION CHART FOR LINEAR APPLICATION								
RATE Fluid ounces/Acre	Rate in Fluid Ounces / 1000 Row Feet Based On Average Row Spacing (in inches):							
	10	15	20	25	30	35	40	45
10	.19	.29	.38	.48	.57	.67	.76	.86
12	.23	.34	.46	.57	.69	.80	.92	1.03
14	.27	.40	.54	.67	.80	.94	1.07	1.21
16	.31	.46	.61	.77	.92	1.07	1.22	1.38
18	.34	.52	.69	.86	1.03	1.21	1.38	1.55
20	.38	.57	.76	.96	1.15	1.34	1.53	1.72
22	.42	.63	.84	1.05	1.26	1.47	1.68	1.89
24	.46	.69	.92	1.15	1.38	1.61	1.84	2.07
26	.50	.75	.99	1.24	1.49	1.74	1.99	2.24
28	.54	.80	1.07	1.34	1.61	1.87	2.14	2.41
30	.57	.86	1.15	1.43	1.72	2.01	2.29	2.58
32	.61	.92	1.22	1.52	1.84	2.14	2.45	2.75

Important: The ADVISE 2FL rate applied affects the length of control and the degree of control. Row-spacing X ADVISE 2FL rate combinations in shaded blocks may not provide long-term, residual control. Use higher listed rates where pest pressure may occur later in crop development or where pest pressure is severe or continuous. Winfield Solutions offers no warranty for use of ADVISE 2FL at rates below 0.7 fluid ounces/1000 row-feet.

TREE, VINE AND FRUIT CROPS

For Foliar Application uniform coverage is required to achieve best control and a spray adjuvant may help improve coverage. Two applications may be required to achieve control when initial insect populations are high. Base retreatment on field scout reports. ADVISE 2FL may be tank mixed with other labeled insecticides to increase control or control pests not controlled by imidacloprid.

BUSHBERRY (Soil Application)

Includes: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

To control Japanese beetle (adults, feeding on foliage), White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle) apply 16.0 to 32.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. 18 inch band on each side of the row followed with 0.25 inches of irrigation immediately after application.

For optimal grub control, apply ADVISE 2FL to control 1st or 2nd instar larvae. Application may be made post-bloom up to 7 days prior to harvest, or post-harvest until October 1st. For optimum control of Japanese beetle larvae, make applications from June 1 to July 15. Do not apply during bloom.

Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root-zone will help protect berry plant roots from grub feeding.

Apply ADVISE 2FL to moist soil. If necessary, apply one hour of irrigation water immediately before application of ADVISE 2FL. To ensure maximum efficacy, ½ to 1 inch of irrigation water or rainfall must be applied or received within 24 hours of application of ADVISE 2FL to facilitate movement into the soil and into the root-zone

Restrictions for Bushberry (Soil Application):

- Pre-Harvest Interval (PHI): 7 days
- Maximum ADVISE 2FL allowed per season: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



BUSHBERRY (Foliar Application)

Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

For control of Aphids and Leafhoppers/Sharpshooters apply 2.4 to 3.2 fluid ounces per acre. Use higher rates when pest pressure more severe.

For control of Japanese beetles (adults), Blueberry Maggot and Thrips apply 4.8 to 6.4 fluid ounces per acre. Use higher listed rates when pest pressure more severe.

For control of Blueberry maggot apply 6.4 fluid ounces per acre.

Restrictions for Bushberries (Foliar Application):

- Pre-Harvest Interval (PHI): 3 days
- Minimum interval between applications : 7 days
- Maximum ADVISE 2FL allowed per crop season: 32.0 fluid ounces/Acre (0.5 lb. AI/A)
- Maximum number of ADVISE 2FL applications per crop season: 5
- Minimum application volume (water): 20.0 GPA - ground; 5.0 GPA - aerial.
- Do not apply pre-bloom or during bloom or when bees are foraging.



CANEBERRY (Soil Application)

BLACKBERRY (Rubus spp. – including Andean Blackberry, Arctic blackberry, Bingleberry, Black satin berry, Boysenberry, Brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyene blackberry, Common blackberry, Coryberry, Darrowberry, Dewberry, Dirksen thornless berry, Evergreen blackberry, Himalayaberry, Hullberry, Lavacaberry, Loganberry, Lowberry, Lucretiaberry, Mammoth blackberry, Marionberry, Moras, Mures deronce, Nectarberry, Northern dewberry, Olallieberry, Oregon evergreen berry, Phenomenalberry, Rangeberry, Ravenberry, Rossberry, Shawnee blackberry, Southern dewberry, Tayberry, Youngberry, Zarzamora, and varieties and/or hybrids of these).

RASPBERRY (Rubus spp. – including Bababerry, Black raspberry, Thimbleberry, Tulameen, Yellow raspberry, and varieties and/or hybrids of these, and Wild raspberry)

To control Rootgrubs (Scarabaeidae) and Rootworms (Chrysomelidae) apply 16.0 to 32.0 fluid ounces per acre.

Apply post-bloom, immediately after bees are removed. Target applications to early instar larvae.

ADVISE 2FL has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the ADVISE 2FL and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

Apply ADVISE 2FL to moist soil.

Immediately upon application, ADVISE 2FL must be incorporated into root-zone by 0.1 – 0.3 inches water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.

Apply specific dosage of ADVISE 2FL in one of the following methods:

1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre;
2. As a chemigation application with 600 to 1000 gal water.

Restrictions for Caneberry (Soil Applications):

- Pre-Harvest Interval (PHI): 7 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.5 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



CANE BERRY (Foliar Application)

BLACKBERRY (Rubus spp. – including Andean Blackberry, Arctic blackberry, Bingleberry, Black satin berry, Boysenberry, Brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyene blackberry, Common blackberry, Coryberry, Darrowberry, Dewberry, Dirksen thornless berry, Evergreen blackberry, Himalayaberry, Hullberry, Lavacaberry, Loganberry, Lowberry, Lucretiaberry, Mammoth blackberry, Marionberry, Moras, Mures deronce, Nectarberry, Northern dewberry, Olallieberry, Oregon evergreen berry, Phenomenalberry, Rangeberry, Ravenberry, Rossberry, Shawnee blackberry, Southern dewberry, Tayberry, Youngberry, Zarzamora, and varieties and/or hybrids of these).

RASPBERRY (Rubus spp. – including Bababerry, Black raspberry, Thimbleberry, Tulameen, Yellow raspberry, and varieties and/or hybrids of these, and Wild raspberry)

To control Aphids, Leafhoppers, and Thrips apply 6.4 fluid ounces per acre.

Restrictions for Caneberry (Foliar Application):

- Pre-Harvest Interval (PHI): 3 days
- Minimum interval between applications: 7 days



- Maximum ADVISE 2FL allowed per crop season: 19.2 fluid ounces/Acre (0.3 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.

CITRUS (Field) (Soil Application)

Includes: calamondin, citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

To control Aphids, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhopper/Sharpshooters, Mealybugs, Scales, Termites (FL only), Whiteflies apply 16.0 to 32.0 fluid ounces per acre.

For suppression of Thrips (foliage feeding thrips only) and in response to symptoms of Citrus tristeza virus (CTV) through vector control and Citrus yellows apply 32.0 fluid ounces per acre.

Applications

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, tickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. Soil should be lightly prewetted to break soil surface tension prior to applications of ADVISE 2FL. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move ADVISE 2FL into root-zone. Allow 24 hours before initiating subsequent irrigations.
2. Soil surface band spray on both sides of the tree. Overlap bands at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less;
3. Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only use on trees up to 8 feet tall.

For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk.

Restrictions for Citrus (Field) (Soil Application):

- Pre-Harvest Interval (PHI): 0 day
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb ai/acre)

CITRUS (Foliar Application)

Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, and other cultivars and/or hybrids of these

To control Aphids, Asian citrus psyllid, Black fly, Leafhoppers/Sharpshooters, Leafminers, Mealybugs, Scales, Whiteflies apply 8.0 to 16.0 fluid ounces per acre, depending on tree size, target pest and infestation pressure. For suppression of Thrips, apply 8.0 to 16.0 fluid ounces per acre.

Restrictions for Citrus (Foliar Application):

- Pre-Harvest Interval (PHI): 0 days
- Minimum interval between applications: 10 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.5 lb. AI/A)
- Do not apply during bloom or within 10 days prior to bloom or when bees are foraging.



Applications

Aerial application of ADVISE 2FL may result in slower activity and reduced control to results from ground application.

Scales - time applications to the crawler stage. Treat each generation. Where concentrated applications are appropriate, increase the spray solution concentration to apply an equivalent rate per acre to that applied in the

diluted application. The 16.0 fluid ounce/Acre rate is based on full sized trees. This rate may be reduced proportionally for smaller trees.

CITRUS (containerized)

Includes: calamondin, citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

To control Aphids, Asian citrus psyllid, Blackfly, Citrus leafminer, Leafhoppers/Sharpshooters), Mealybugs, Scales, Whiteflies apply 0.75 milliliter per cubic foot of container media.

To control Citrus root weevil (larval complex) apply 1.25 to 2.50 milliliters per cubic foot of container media.

To suppress Citrus thrips apply to 2.50 milliliters per cubic foot of container media.

Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of ADVISE 2FL per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, make treatments at planting prior to insect infestation. Repeat if necessary. For control of larvae of the citrus root weevil complex, make application prior to neonate larvae entering potting media. Utilize higher dosage for heavy infestation.

Restrictions for Citrus (containerized):

- Pre-Harvest Interval (PHI): 0 days
- Maximum ADVISE 2FL allowed per application: 0.5 mLs/0.1 ft³ container media.
- Maximum ADVISE 2FL allowed per crop season: 3.5 mLs/plant.
- Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.



CRANBERRY (Soil Application Only)

To control Rootgrubs (Scarabaeidae), Rootworms (Chrysomelidae) apply 16.0 to 32.0 fluid ounces per acre.

Apply ADVISE 2FL to moist soil. Apply specified dosage of ADVISE 2FL in one of the following methods:

1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre.
2. As a chemigation application with 600 to 1000 gal water

Immediately upon application, ADVISE 2FL must be incorporated into root-zone by 0.1 to 0.3 inches water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.

Rootgrubs and Rootworms:

Apply post bloom (after bees are removed) to coincide with the early instar larvae stage.

ADVISE 2FL has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the ADVISE 2FL and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

Restrictions for Cranberry:

- Pre-Harvest Interval (PHI): 30 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



GRAPE (Soil Application)

Including: American bunch grape, Muscadine grape, and Vinifera grape

To control Mealybugs, Leafhoppers/Sharpshooters, Phylloxera* spp. Apply 16.0 to 32.0 fluid ounces per acre.

To suppress Pierce's Disease apply 24.0 to 32.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation.
3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

For optimum results, make application(s) between bud-break and the pea-berry stage.

*Repeated and regular use of ADVISE 2FL over several, consecutive growing seasons controls existing Phylloxera infestations over time or prevents Phylloxera from becoming established.

Restrictions for Grape (Soil Application):

- Pre-Harvest Interval (PHI): 30 days
- Maximum ADVISE 2FL allowed per season: 32.0 fluid ounces/Acre (0.50 lb ai/acre)

GRAPE (Foliar Application)

American bunch grape, Muscadine grape and Viniferous grape

For control of Leafhoppers/Sharpshooters and Mealybugs apply 2.4 to 3.2 fluid ounces per acre. Use higher rates when pest pressure more severe.

For control of Grapeleaf skeletonizer apply 3.0 fluid ounces per acre. Control can usually be achieved with ground applications that provide more thorough coverage of foliage. Aerial applications may only provide suppression due to lack of thorough coverage.

Restrictions for Grapes (Foliar Application):

- Pre-Harvest Interval (PHI): 0 days
- Maximum interval between applications: 14 days
- Maximum ADVISE 2FL allowed per year: 6.0 fluid ounces/Acre (0.1 lb. AI/A)

HOPS (Soil Application)

To control Aphids apply 19.2 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation.
3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

Restrictions for Hops (Soil Application):

- Pre-Harvest Interval (PHI): 60 days
- Maximum ADVISE 2FL allowed per year: 19.2 fluid ounces/Acre (0.30 lb ai/acre)

HOPS (Foliar Application)

For control of Aphids apply 6.4 fluid ounces per acre.

Restrictions for Hops (Foliar Application):

- Pre-Harvest Interval (PHI): 28 days
- Maximum interval between applications: 21 days
- Maximum ADVISE 2FL allowed per year: 19.2 fluid ounces/Acre (0.30 lb. AI/A)

POME FRUIT (Soil Application)

Including: apple, crabapple, loquat, mayhaw, pear (including Oriental pear), quince

To control Aphids (including woolly apple aphid) and Leafhoppers, apply 16.0 to 24.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.



Restrictions for Pome Fruit (Soil Application):

- Pre-Harvest Interval (PHI): 21 days
- Maximum ADVISE 2FL allowed per season: 24.0 fluid ounces/Acre (0.38 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.

POME FRUIT (Foliar Application)

Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

The amount of ADVISE 2FL required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. To calculate the rate needed on smaller trees, multiply the pest specific rate (e.g., for aphid control, 1.6 fluid ounces/100 gallons) times the number of 100 gallons of spray solution required to thoroughly wet foliage just prior to the point of runoff, on one acre of the trees being treated. For concentrate sprays, apply the same amount of ADVISE 2FL per acre as would be applied in a dilute spray based on tree size and foliage volume.

To control Leafhoppers apply 3.2 to 6.4 fluid ounces of ADVISE 2FL per acre. Apply low rate for low to moderate populations of white apple leafhoppers and high rate for high populations or for other leafhopper species. Apply ADVISE 2FL while most leafhoppers are in the nymphal stage.

To control Aphids (except wooly apple aphid), Leafminers, San Jose scale, apply 6.4 fluid ounces of ADVISE 2FL per acre. For first generation leafminer control, make application after pollination is complete and bees are no longer present in the orchard. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, better control will be obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. ADVISE 2FL will not control late instar larvae. For rosy apple aphid - apply prior to leafrolling caused by rosy apple aphid. For San Jose scale - time applications to the crawler stage. Treat each generation.

To control Mealybugs and Pear psylla on pear only, apply 16.0 fluid ounces of ADVISE 2FL per acre. For mealybugs - apply maximum gallonage for tree with ground equipment. Ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of mealybugs.

Restrictions for Pome Fruits (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Maximum interval between applications: 10 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.5 lb. AI/A)
- Do not apply pre-bloom or during bloom or when bees are foraging.



Aerial application of ADVISE 2FL may result in slower activity and reduced control compared to ground application due to less thorough coverage.

POMEGRANATE (Soil Application)

To control Aphids, Leafhoppers/Sharpshooters, and Whiteflies apply 16.0 to 24.0 fluid ounces per acre.

Apply specific dosage of ADVISE 2FL in one of the following methods:

Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Restrictions for Pomegranate (Soil Application):

- Pre-Harvest Interval (PHI): 0 days
- Maximum ADVISE 2FL allowed per crop season: 32.0 fluid ounces/Acre (0.5 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



POMEGRANATE (Foliar Application)

To control Aphids, Leafhoppers/Sharpshooters, and Whiteflies apply 6.4 fluid ounces per acre.

For suppression in response to Scales apply 6.4 fluid ounces per acre.

Restrictions for Pomegranate (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 7 days
- Maximum ADVISE 2FL allowed per crop season: 19.2 fluid ounces/Acre (0.3 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



STONE FRUIT (Soil Application)

Including: apricot, cherry (including sweet and tart), nectarine, peach, plum (including Chickasaw, Damson and Japanese), Plumcot, prune (fresh and dried)

To control Aphids (including woolly apple aphid) and Leafhoppers apply 16.0 to 24.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Restrictions for Stone Fruit (Soil Application):

- Maximum ADVISE 2FL allowed per season: 24.0 fluid ounces/acre (0.38lb ai/acre)
- Pre-Harvest Interval (PHI): 21 days
- Do not apply pre-bloom or during bloom or when bees are foraging.



STONE FRUIT (Foliar Application)

Apricot, Cherry (including sweet and tart), nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

To control Aphids, Green June beetle, Japanese beetle, Leafhoppers/Sharpshooters, Plant bugs, Rose chafer, San Jose scale apply 3.2 to 6.4 fluid ounces of ADVISE 2FL per acre. To control Cherry fruit fly (maggot of Eastern and Western) apply 4.8 to 6.4 fluid ounces of ADVISE 2FL per acre. For suppression of Plum curculio and Stink bugs apply 6.4 fluid ounces of ADVISE 2FL per acre.

Apply in a minimum application volume (water) of 50 gallons per acre for ground applications and 25 gallons per acre for aerial applications.

Aerial application of ADVISE 2FL may result in slower activity and reduced control relative to results from ground application

Restrictions for Apricot, Nectarine, Peach (Foliar Application):

- Pre-Harvest Interval (PHI): 0 day
- Minimum interval between applications: 7 days
- Maximum ADVISE 2FL allowed per year: 19.2 fluid ounces/Acre (0.30 lbs. AI/A)
- Do not apply pre-bloom or during bloom or when bees are foraging.



Restrictions for Cherries, Plums, Plumcot, Prune (Foliar Application):

- Pre-Harvest Interval (PHI): 7 day
- Minimum interval between applications: 10 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lbs. AI/A)
- Do not apply pre-bloom or during bloom or when bees are foraging.



STONE FRUIT (Preplant, Root Drip Application)

To control Black peach aphid (infesting roots) apply 2.0 fluid ounces per 10 gallons of root dip solution.

Mix ADVISE 2FL at 2.0 fluid ounces per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the ADVISE 2FL solution for up to 5 minutes. Allow solution to dry on

roots and transplant trees as soon as possible following treatment.

TREE NUTS (except Almond) (Soil Application)

Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

To control Aphids, Leafhoppers/Sharpshooters, Mealybugs, Spittlebugs, Termites and Whiteflies apply 16.0 to 32.0 fluid ounces per acre.

For suppression in response to Pecan scab (from reduction in honeydew deposition) apply 16.0 to 32.0 fluid ounces per acre. For Thrips (foliage-feeding thrips only) apply 32.0 fluid ounces per acre.

Use the higher rates within rate range when applied by shank or subsurface sidedress, used on larger trees, soils with high clay content, for high plant populations, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.

Apply specific dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent irrigation equipment. Pre-wet soil prior to applications of ADVISE 2FL and allow soil to dry following application and prior to subsequent irrigation;
2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site;
3. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Product should be applied in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area should follow within 48 hours to promote uptake by root system.
4. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18 – 24 inches to obtain optimum control. Allow soil to dry following treatment and prior to apply any irrigation.

Restrictions for Tree Nuts (Soil Application):

- Pre-Harvest Interval (PHI): 7 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.5 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



TREE NUTS (except Almonds) (Foliar Application)

Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut [black and English]

To control Aphids (except black pecan aphid), Leafhoppers/Sharpshooters, Phylloxera sp. (leaf infestations, Spittlebugs and Whiteflies apply 2.8 to 5.6 fluid ounces per acre.

To control Black pecan aphid, Mealybugs and San Jose scale apply 6.4 fluid ounces per acre.

Time applications for control of San Jose scale according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.

Restrictions for Tree Nuts (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 6 days
- Maximum ADVISE 2FL allowed per year: 22.8 fluid ounces/Acre (0.36 lb ai/acre)
- Minimum application volume (water): 50 GPA – ground application, 25 GPA – aerial application.
- Do not apply pre-bloom or during bloom or when bees are foraging.



TROPICAL FRUIT (Soil Application)

Including: acerola, avocado, black sapote, canistel, feijoa, jaboticaba, guava, longan, lychee, mamey sapote, mango, papaya, passionfruit, persimmon, pulasan, rambutan, sapodilla, Spanish lime, star apple, starfruit, wax jambu

To control Aphids, Leafhoppers, Whiteflies apply 24.0 to 32.0 fluid ounces per acre.

To suppress Scales apply 32.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Restrictions for Tropical Fruit (Soil Application):

- Preharvest Interval (PHI): 6 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



TROPICAL FRUIT (Foliar Application)

Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Pulasan, rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu

For control of Aphids, Leafhoppers /Sharpshooters, Thrips, and Whiteflies apply 6.4 fluid ounces per acre.

For suppression of Scales apply 6.4 fluid ounces per acre.

Restrictions for Tropical Fruit (Foliar Application):

- Pre-Harvest Interval (PHI): 7 days
- Minimum interval between applications: 10 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb, AI/A)
- Do not apply pre-bloom or during bloom or when bees are foraging.



Aerial application of ADVISE 2FL may result in slower activity and reduced control compared to ground application due to less thorough coverage.

OTHER CROPS

POPLAR/COTTONWOOD (Soil Application)

Includes: members of the genus *Populus* grown for pulp or timber

To control Aphids and Cottonwood leaf beetle apply 16.0 to 32.0 fluid ounces per acre.

To suppress *Phylloxera popularia* apply 16.0 to 32.0 fluid ounces per acre.

Apply specified dosage of ADVISE 2FL in the following method:

1. Chemigation through low-pressure drip irrigation.

For Cottonwood leaf beetle, protection against damage will occur when application is made early, when the beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake.

For *Phylloxera*, apply early in the year, from break of dormancy through May.

Restrictions for Poplar/Cottonwood (Soil Application):

- Maximum ADVISE 2FL allowed per season: 32.0 fluid ounces/Acre (0.50 lb ai/acre)
- Do not apply pre-bloom or during bloom or when bees are foraging.



POPLAR/COTTONWOOD (Foliar Application)

(includes members of the genus Populus grown for pulp or timber)

Apply as foliar spray at specified rate per acre when insect pressure reaches economic threshold. Uniform coverage is required to achieve best control and a spray adjuvant may help improve coverage. Two applications may be required to achieve control when initial insect populations are high. Retreatment should be based on field scout reports. ADVISE 2FL may be tank mixed with other labeled insecticides to increase control or control pests not controlled by imidacloprid.

For control of Aphids and Leaf beetles apply 3.2 to 6.4 fluid ounces per acre. Use higher listed rates when pest pressure more severe.

Restrictions for Poplar/Cottonwood (Foliar Application):

- Minimum interval between applications: 10 days
- Maximum ADVISE 2FL allowed per crop season: 32.0 fluid ounces/Acre (0.50 lb. AI/A)
- Do not apply pre-bloom or during bloom or when bees are foraging.



Aerial application of ADVISE 2FL may result in slower activity and reduced control compared to ground application due to less thorough coverage.

CHRISTMAS TREE (Foliar Application)

For control of Aphids, Adelgids and Sawflies apply 3.2 to 6.4 fluid ounces per acre. Use higher rate when pest pressure more severe.

Restrictions for Christmas Trees:

- Minimum interval between applications: 7 days
- Maximum ADVISE 2FL allowed per year: 32.0 fluid ounces/Acre (0.50 lb. AI/A)

Aerial application of ADVISE 2FL may result in slower activity and reduced control compared to ground application due to less thorough coverage.

Gall-forming adelgids - time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. After galls form, spraying will no longer be ineffective.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

AgriSolutions™ is a trademark and Advise® is a registered trademark of Winfield Solutions, LLC.