



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Avenue NW
Washington, D.C. 20460

EPA Reg. Number

1381-211

Date of Issuance

03/05/07

Term of Issuance:

Conditional

Name of Pesticide Product

Taiga Z Insecticide

NOTICE OF PESTICIDE:

Registration

Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Agriliance, LLC
P.O. Box 64089
St. Paul, MN 55164-0089

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA section 4.
2. Make the following labeling changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read: "EPA Registration No. 1381-211"

Signature of Approving Official:

George LaRocca
Product Manager (Team 13)
Insecticide Branch
Registration Division (7505P)

Date:

March 05, 2007

- b. Add the following to the Precautionary Statement: "***Remove and wash contaminated clothing before reuse***" and "***Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals***".
 - c. Revise the subtitle "Resistance" to read "***Resistance Management***".
 - d. In Specific Use Directions, delete crops under Fruiting Vegetables (Tomato, etc.); or provide appropriate data or citation(s) to support these uses.
 - e. In Specific Use Directions, you may delete "Tobacco variety descriptors (Air dried . . . etc.)".
3. Submit two (2) copies of the final printed label bearing the above stated revisions prior to releasing this product for sale.

Please note that you indicated in your cover letter, the inclusion of 3 Confidential Statements of Formula -CSF- (Basic, Alternates A and B). However, we received, reviewed and consider only the basic CSF for this registration.

If you fail to comply with the above stated conditions, this registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosed, please find a copy of the stamped label for your records. If you have any questions concerning this action, please contact Dr. B. A. Akinlosotu at (703) 605-0653.

Enclosure

3 8 36

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



Taiga™ Z Insecticide

Active Ingredient:

Lambda-cyhalothrin	
[1α(S*),3α(Z)]-(±)-cyano-(3-phenoxyphenyl)methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate	11.4%
Other Ingredients:	88.6%
Total:	100.0%

Taiga™ Z Insecticide contains 1 lb. of active ingredient per gal. and is a capsule suspension.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements, complete storage and disposal, and directions for use in booklet.

EPA Reg. No. 1381-

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 5 2007**

EPA Est. 070989-MO-001

Net Contents
PROP11/10/06

Distributed By
Agrilience, LLC
P.O. Box 64089
St. Paul, MN 55164-0089

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
1381-211**

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call 1-877-424-7452.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. May cause allergic skin reactions. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

Skin exposure may also result in a sensation described as a tingling, itching, burning, or prickly feeling. Onset may occur immediately to 4 hrs. after exposure and may last 2-30 hrs., without damage. Wash exposed areas once with soap and water. Relief from the skin sensation may be obtained by applying an oil-based cream.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton \geq 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL**Prohibitions**

Do not contaminate water, food, or feed by storage and disposal.

Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent); then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

Container Precautions

Before refilling RETURNABLE CONTAINERS, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions, and damaged or worn threads on closure devices.

REFILL ONLY WITH TAIGA™ Z INSECTICIDE. The contents of RETURNABLE CONTAINERS cannot be completely removed by cleaning. Refilling with materials other than Taiga™ Z Insecticide will result in contamination and may weaken container. After filling and before transporting, check for leaks.

Do not refill or transport damaged or leaking container.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Taiga™ Z Insecticide may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

RESISTANCE

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

SPRAY DRIFT PRECAUTIONS

Observe the Following Precautions When Spraying in the Vicinity of Aquatic Areas Such as Lakes; Reservoirs; Rivers; Permanent Streams, Marshes or Natural Ponds; Estuaries and Commercial Fish Farm Ponds.

- Do not apply by ground within 25 ft., or by air within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 ft. when ultra-low volume (ULV) application is made.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.
- Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.
- Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 ft. above the crop canopy should be avoided.
- Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3–10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.
- Risk of exposure to aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.
- Do not cultivate within 10 ft. of the aquatic area so as to allow growth of a vegetative filter strip.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.
- Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

In the State of New York, a 25 ft. vegetated, non-cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or stream that drains into a coastal salt marsh, for both aerial or ground application. For aerial applications, the 25 ft. vegetated non-cropped buffer strip for runoff protection would be part of the larger 150 ft. buffer strip (or 450 ft. buffer strip for ULV application) required for spray drift.

TANK MIX APPLICATION

When tank mixing with any other agricultural products, **always add Taiga™ Z Insecticide last**. Fill the tank with one half to two thirds volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of Taiga™ Z Insecticide to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While Taiga Z Insecticide has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture.

Taiga Z Insecticide is an aqueous based formulation. It is recommended that no type of non-emulsifiable oils be used in combination with Taiga Z Insecticide. If adjuvants are used, use only:

- Nonionic Surfactant (NIS) containing at least 75% surface agent, or
- Non-phytotoxic Crop Oil Concentrate (COC), including once refined Vegetable Oil Concentrate (VOC),
- Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier.

Adjuvants other than NIS or COC may be used providing the product meets the following criteria:

1. Contains only EPA exempt ingredients.
2. Is non-phytotoxic to the target crop.
3. Is compatible in mixture. (May be established through a jar test.)
4. Is supported locally for use with Taiga Z Insecticide on the target crop through proven field trials and through university and extension recommendations.

In addition, the following may be used as diluents:

Crop Oil Concentrate; Methylated Sunflower Oils; Urea-Ammonium Nitrate

It is recommended that the following not be used in combination Taiga Z Insecticide as diluents or adjuvants:

Non-emulsifiable oils; Diesel Fuel; Straight Mineral Oil

CHEMIGATION

Sprinkler Irrigation Application

Apply Taiga Z Insecticide at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types, (see TANK MIX APPLICATION) rates and mixing instructions. These recommendations should be proven, through university and extension field trials, to be effective with Taiga Z Insecticide applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Taiga Z Insecticide into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1–0.2 acre-inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Taiga Z Insecticide for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that Taiga Z Insecticide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions: Sprinkler Irrigation Application

- A. Apply this product only through (sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Do not apply through chemigation systems connected to public water systems.

12836

SPECIFIC USE DIRECTIONS

AGRICULTURAL USES

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
ALFALFA AND ALFALFA GROWN FOR SEED			
	Alfalfa Caterpillar Cutworm species Army Cutworm Green Cloverworm Looper species Velvetbean Caterpillar Webworm species Leafhopper species Threecornered Alfalfa Hopper	0.015-0.025	1.92-3.20
	Armyworm Corn Earworm Fall Armyworm ¹ Western Yellow-striped Armyworm Yellow-striped Armyworm Alfalfa Weevil Bean Leaf Beetle (Adult) Blister Beetle species Clover Leaf Weevil species Clover Root Borer (Adult) Clover Root Curculio species (Adult) Clover Stem Borer (Adult) Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle species (Adult) Egyptian Alfalfa Weevil Grape Colaspis (Adult) Green June Beetle (Adult) Japanese Beetle (Adult) Mexican Bean Beetle Pea Weevil (Adult) Sweet Clover Weevil (Adult) Whitefringed Beetle species (Adult) Meadow Spittlebug Plant Bug species including Lygus species ³ Stink Bug species Alfalfa Seed Chalcid (Adult) Blue Alfalfa Aphid Cowpea Aphid Green Peach Aphid ³ Pea Aphid Spotted Alfalfa Aphid Thrips species ⁴ Grasshopper species	0.02-0.03	2.56-3.84
	Beet Armyworm ^{1,3} Blotch Leafminer ³ Spider Mites ²	0.03	3.84

Remarks

- Apply only to fields planted to pure stands of alfalfa.
- Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals./A by air or 10 gals./A by ground. When foliage is dense and/or pest populations are high 5-10 gals./A by air or 20 gals./A by ground and higher use rates are recommended. Use higher rates for increased residual control.
- Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters.
- Do not apply more than 0.03 lb. a.i. (0.24 pts.)/A per cutting.
- Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season.
- Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.

¹ Use higher rates for large larvae.

² Suppression only.

³ See resistance statement under **General Directions for Use.**

⁴ Does not include Western Flower Thrips.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CANOLA	Cutworm species Looper species Armyworm species Diamondback Moth Flea Beetle Cabbage Seedpod Weevil Lygus Bug Grasshoppers	0.015-0.03	1.92-3.84
	Cabbage Aphid	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply a minimum of 2 gals. of water/A.
- Do not apply within 7 days of harvest.
- Do not apply more than 0.09 lb. a.i. (0.72 pts.)/A per year.

Crop	Target Pest	Rate	
		lb. a.i./A	fl.oz./A
CEREAL GRAINS			
Corn (At Plant): Field Corn Popcorn Seed Corn Sweet Corn	Wireworm species Cutworm species Seedcorn Maggot White Grub species Corn Rootworm Larvae Western Northern Southern Mexican Seedcorn Beetle Lesser Cornstalk Borer Red Imported Fire Ant ¹	0.005 lbs. a.i. per 1000 feet of row ²	0.66 fl. oz. per 1000 feet of row ²

Remarks

- **Banded Applications** – Apply at planting as a 5–7 inch T–band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel.
- **In–Furrow Applications** – Apply into the seed furrow through spray nozzles or microtubes, behind the planter furrow openers and in front of the press wheel.
- Apply a minimum of 3 gals. finished spray/A.
- Do not harvest or graze livestock or cut treated crops for feed within 21 days of at plant application.
- Do not apply more than 0.09 lb. a.i. (0.72 pts.)/A per crop at plant.
- For field corn, popcorn, and seed corn do not apply more than 0.12 lb. a.i./A per crop from at plant and foliar applications. For sweet corn do not apply more than 0.48 lb. a.i./A per crop from at plant and foliar applications.

¹Suppression only.

² Lbs. a.i. and fl. oz./A of Taiga™ Z INSECTICIDE Applied at 0.66 fl. oz./1000 ft. of Row for Various Row Spacings						
Row Spacing	40"	38"	36"	34"	32"	30"
Linear Ft./A	13,068	13,756	14,520	15,374	16,335	17,424
Lbs. a.i./A	0.067	0.07	0.075	0.079	0.084	0.09
Fl. oz./A	8.6	9.1	9.6	10.1	10.8	11.5

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CEREAL GRAINS			
Corn (Foliar) Field Corn Popcorn Seed Corn	Cutworm species Western Bean Cutworm ¹ Corn Earworm ¹ Green Cloverworm Meadow Spittlebug	0.015-0.025	1.92-3.20
	Tobacco Budworm ^{1,4} European Corn Borer ¹ Southwestern Corn Borer ¹ Stalk Borer ¹ Hop Vine Borer ¹ Lesser Cornstalk Borer Armyworm ² Fall Armyworm ² Yellow-striped Armyworm ² Webworm species Flea Beetle species Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Bean Leaf Beetle Cereal Leaf Beetle Japanese Beetle (Adult) Sap Beetle (Adult) Seedcorn Beetle Stink Bug species Grasshopper species Corn Leaf Aphid ³ Bird Cherry-Oat Aphid ³ English Grain Aphid ³	0.02-0.03	2.56-3.84
	Mexican rice Borer ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹ Beet Armyworm ⁴ Southern Corn Leaf Beetle ³ Chinch Bug Green Bug ^{3,4}	0.03	3.84

Remarks

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5-day intervals if needed. TAIGA™ Z Insecticide may only suppress heavy infestations and/or subsequent migrations.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 3.84 fl. oz./A (0.03 lb. a.i./A).
- Do not apply within 21 days of harvest.
- Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per crop from at plant and foliar applications.
- Do not apply more than 0.06 lb. a.i. (0.48 pts.) after silk initiation. Do not apply more than 0.03 lb. a.i. (0.24 pts.) after corn has reached the milk stage (yellow kernels with milky fluid).

¹For control before the larva bores into the plant stalk or ear.

²Use higher rates for large larvae.

³Suppression only.

⁴See resistance statement under General Directions for Use.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CEREAL GRAINS			
Sweet Corn (Foliar)	Corn Earworm Armyworm ¹ Fall Armyworm ¹ Southern Armyworm ¹ Beet Armyworm ^{1,3} Yellow-Striped Armyworm ¹ Cutworm species Western Bean Cutworm Webworm species European Corn Borer Southwestern Corn Borer Common Cornstalk Borer Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Japanese Beetle (Adult) Sap Beetle (Adult) Flea Beetle species Tarnished Plant Bug Stink Bug species Chinch Bug Aster Leafhopper Grasshopper species Aphid species ^{2,3} Spider Mite species ²	0.02-0.03	2.56-3.84
	Corn Silkfly (Adult) ²	0.03	3.84

Remarks

- Apply as required by scouting, or locally prescribed corn growth stages, usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be targeted for control before insects enter the stalk or ear.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present). When applying by air, apply in a minimum of 2 gals. of water/A.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 3.2 fl. oz./A (0.025 lb a.i./A).
- Do not apply within 1 day of harvest.
- Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment.
- Do not apply more than 0.48 lb. a.i. (3.84 pts./A) per crop from at plant and foliar applications.

¹Use higher rates for large larvae.

²Suppression only.

³See resistance statement under General Directions for Use.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CEREAL GRAINS			
Rice	True Armyworm Fall Armyworm Yellow-striped Armyworm Rice Water Weevil (Adult) Rice Stink Bug Chinch Bug Grasshopper species Leafhopper species Sharpshooter species Bird Cherry-Oat Aphid Yellow Sugarcane Aphid Green Bug Mexican Rice Borer ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹ European Corn Borer ¹ Rice Seed Midge	0.025-0.04	3.20-5.12

Remarks

- Apply as required by scouting. Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds. Determine the need for repeat applications, usually at intervals of 5-7 days, by scouting.
- Taiga Z Insecticide can be safely used when propanil products are being used for weed control.
- Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals of water (or a total carrier volume)/A but ensure sufficient volume is used to provide adequate coverage. In addition, adding an emulsifiable crop oil (e.g., 1 pt./A) when lower aerial application volumes are used is recommended to help improve coverage, reduce evaporation, and improve efficacy.
- For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- For control of rice water weevil in water seeded rice, make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.
- California: In addition to above directions for control of rice water weevil in water seeded rice Taiga Z Insecticide may be applied at the 1-3 leaf growth stage, with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults, based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field, or b) spray the entire field.
- Green bug is known to have many biotypes. Taiga Z Insecticide may only provide suppression. If satisfactory control is not achieved with the first application of Taiga Z Insecticide, a resistant biotype may be present. Use alternate chemistry for control.
- Do not release flood water within 7 days of an application.
- Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season.
- Do not apply more than 0.04 lb. a.i. (0.32 pts.)/A within 21 to 27 days of harvest.
- Do not apply within 21 days of harvest.
- Do not use treated rice fields for the aquaculture of edible fish and crustacea.
- Do not apply as an ultra-low volume (ULV) spray.

¹ For control before the larvae bores into the plant stalk.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CEREAL GRAINS			
Sorghum (Grain)	Cutworm species Sorghum Midge	0.015–0.02	1.92–2.56
	Amyworm Beet Armyworm ³ Fall Armyworm ¹ Yellow-striped Armyworm ¹ Corn Earworm Webworm species European Corn Borer ² Southwestern Corn Borer ² Lesser Cornstalk Borer ² Flea Beetle species Stink Bug species Grasshopper species	0.02–0.03	2.56–3.84
	Mexican Rice Borer ² Rice Stalk Borer ² Sugarcane Borer ² Chinch Bug	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A.
- For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5-day intervals if needed.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3-5-day intervals if needed. Taiga Z Insecticide may only suppress heavy infestations and/or subsequent migrations.
- Do not apply more than 0.08 lb. a.i. (0.64 pts./A) per season.
- Do not apply more than 0.06 lb. a.i. (0.48 pts./A) per season after crop emergence.
- Do not apply more than 0.02 lb. a.i. (0.16 pts./A) per season once crop is in soft dough stage.
- Do not apply within 30 days of harvest.

¹Use higher rates for large larvae.²For control before the larva bores into the plant stalk.³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CEREAL GRAINS			
Wheat Wheat Hay Triticale	Cutworm species Army Cutworm	0.015–0.025	1.92–3.20
	Armyworm Fall Armyworm Yellow-striped Armyworm Flea Beetle species Cereal Leaf Beetle Stink Bug species English Grain Aphid ¹ Russian Wheat Aphid ¹ Bird Cherry-Oat Aphid ¹ Grasshopper species Hessian Fly ⁴ Orange Blossom Wheat Midge	0.02–0.03	2.56–3.84
	Grass Sawfly	0.025–0.03	3.20–3.84
	Chinch Bug Greenbug ^{1,3} Corn Leaf Aphid ² Mite species ²	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- For chinch bug control, repeat applications at 3-5-day intervals if needed. Taiga Z Insecticide may only suppress heavy infestations and/or migrations.
- Greenbug is known to have many biotypes. Taiga Z Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.
- Do not apply within 30 days of harvest.
- Do not allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment. Do not feed treated straw to meat or dairy animals within 30 days after the last treatment.
- Do not apply more than 0.06 lb. a.i. (0.48 pts./A) per season.

¹Best control is obtained before insects begin to roll leaves. Once wheat has started to boot, Taiga Z Insecticide may provide suppression only. Higher rates and increased coverage will be necessary.

²Suppression only.

³See resistance statement under **General Directions for Use**.

⁴Make applications when adults emerge.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
COLE CROPS(Head and Stem Brassica)			
Broccoli Brussels Sprouts Cabbage Cavalo Broccoli Cauliflower Chinese Broccoli (gai lon) Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) Kohlrabi	Alfalfa Looper Cabbage Looper Imported Cabbageworm Southern Cabbageworm Cutworm species Cabbage Webworm	0.015–0.025	1.92–3.20
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow-striped Armyworm Corn Earworm Flea Beetle species Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper species Leafhopper species Plant Bug species including Lygus species ³ Stink Bug species Meadow Spittlebug Aphid species ^{2,3} Whitefly species ^{2,3} Thrips species ² Spider Mite species ²	0.02–0.03	2.56–3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 1 day of harvest.
- Do not apply more than 0.24 lb. a.i. (1.92 pts./A) per season.

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
COTTON	Cutworm species Tobacco Thrips Soybean Thrips	0.015–0.02	1.92–2.56
	Lygus Bug species ³ Pink Bollworm Cabbage Looper Cotton Leafperforator Saltmarsh Caterpillar Cotton Leafworm Cotton Fleahopper	0.02–0.03	2.56–3.84
	Cotton Bollworm Tobacco Budworm ³ Boll Weevil Fall Armyworm Beet Armyworm ^{1,3} European Corn Borer Brown Stink Bug Green Stink Bug Southern Green Stink Bug Twospotted Spider Mite ² Cotton Aphid ^{2,3} Bandedwing Whitefly ^{2,3} Sweetpotato Whitefly ^{2,3}	0.025–0.04	3.20–5.12

Remarks

- Apply as required by scouting, usually at intervals of 5–7 days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage.
- Applications may also be made with equipment adapted and calibrated for ULV sprays. Taiga Z Insecticide may be mixed with once-refined vegetable oil and applied in a minimum of at least one qt. of finished spray/A.
- Under light bollworm/budworm infestation levels, 0.02 lb. a.i./A may be applied in conjunction with intense field monitoring.
- For boll weevil control spray on a 3–5 day schedule.
- When applied according to label directions for control of cotton bollworm and tobacco budworm, Taiga Z Insecticide also provides ovicidal control of unhatched Heliiothine species eggs.
- Do not apply within 21 days of harvest.
- Do not graze livestock in treated areas.
- Do not apply more than 1.6 pts. (0.2 lb. a.i./A) per season.
- Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ammo® Insecticide, Asana® XL Insecticide, Baythroid® Emulsifiable Pyrethroid Insecticide, Capture® Insecticide/Miticide, Danitol® 2.4 EC Spray Insecticide/Miticide, Decis® Insecticide, Fury™ Insecticide, Karate® Insecticide, Karate® Insecticide with Zeon™ Technology, Mustang® Insecticide, Scout X-TRA® Insecticide, Taiga Z Insecticide, Warrior® Insecticide and Warrior Insecticide with Zeon™ Technology.

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under General Directions for Use.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
FRUITING VEGETABLES			
Tomato Tomatillo Peppers (bell and nonbell) Eggplant Ground cherry Pepino	Cabbage Looper Cutworm species Hornworm species	0.015–0.025	1.92–3.20
	Tomato Fruitworm Tobacco Budworm ³ Tomato Pinworm Beet Armyworm ^{1,3} Southern Armyworm ¹ Yellow-striped Armyworm ¹ Fall Armyworm ¹ European Corn Borer ⁴ Leafminer species ² Colorado Potato Beetle ³ Flea Beetle species Grasshopper species Leafhopper species Aphid species ^{2,3} Whitefly species ^{2,3} Meadow Spittlebug Stink Bug species Plant Bug species Stalk Borer ⁴ Blister Beetle species Japanese Beetle (Adult) Pepper Weevil (Adult) ² Vegetable Weevil (Adult) Tomato Psyllid ^{2,3} Spider Mite species ² Thrips ⁵ Cucumber Beetle species (Adult)	0.02–0.03	2.56–3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 5 days of harvest.
- Do not apply more than 0.36 lb. a.i. (2.88 pts./A) per season.

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under **General Directions for Use**.

⁴For control before the larva bores into the plant stalk or fruit.

⁵Does not include Western Flower Thrips.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
LEGUME VEGETABLES (Beans and Peas)			
Edible Podded (Only)	Cutworm species Green Cloverworm Imported Cabbageworm Saltmarsh Caterpillar Velvetleaf Caterpillar Mexican Bean Beetle	0.015–0.025	1.92–3.20
<i>Canavalia gladiata</i> – sword bean			
<i>Canavalia ensiformis</i> – jackbean			
<i>Glycine max</i> – Soybean (immature seed)	Corn Earworm Painted Lady Butterfly (Larva) European Corn Borer Looper Species Western Bean Cutworm	0.02–0.03	2.56–3.84
Edible Podded, Succulent Shelled or Dried Shelled	Tobacco Budworm ⁴ Armyworm ² Fall Armyworm ² Yellow-Striped Armyworm ² Western Yellow-Striped Armyworm ² Bean Leafskeltonizer Webworm species Leaf-tier species Alfalfa Caterpillar Stalk Borer ¹ Cucumber Beetle species (Adult) Corn Rootworm Beetle species (Adult) Flea Beetle species (Adult) Curculio and Weevil species ¹ (foliage and pod feeding adults and larvae) Blister Beetle species Bean Leaf Beetle Japanese Beetle (Adult) Leafhopper species Flea Hopper species Three-Cornered Alfalfa Hopper Meadow Spittlebug Stink Bug species Plant Bug species Including Lygus species ⁴ Grasshopper species Thrips species ^{4,5} Aphid species ⁴		
<i>Phaseolus</i> species – includes: field, kidney, lima, navy, pinto, runner, snap, tepary and wax beans			
<i>Vigna</i> species – includes: adzuki, asparagus, moth, mung, rice, urd and yardlong beans, black- eye pea, catjang, Chinese longbean, cowpea, Crowder pea, and Southern pea			
<i>Pisum</i> species – includes: dwarf, edible-pod, English, field, garden, green, snow and sugar snap peas			
<i>Cajanus cajan</i> – Pigeon pea			

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
LEGUME VEGETABLES (Beans and Peas)			
(continued)			
Succulent Shelled or Dried Shelled	Beet Armyworm ^{3,4} Soybean Looper ^{3,4} Lesser Cornstalk Borer ³ Leafminer species ^{3,4} Whitefly species ^{3,4} Spider Mite species ³	0.03	3.84
<i>Vicia faba.</i> – broadbean (favabean)			
Dried Shelled (Only)			
<i>Lupinus</i> species – includes: grain, sweet, white and sweet white lupines			
<i>Cicer arietinum</i> – chickpea (garbonzo bean)			
<i>Cyamopsis tetragonoloba</i> – guar			
<i>Lablab purpureus</i> – Lablab bean (hyacinth bean)			
<i>Lens esculata</i> – Lentils			

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest.
- For dried shelled legume vegetables, do not apply within 21 days of harvest.
- Do not apply more than 0.12 lb. a.i. (0.96 pts./A) per season.
- For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay.

¹For control before the larva bores into the plant stalk or pods²Use higher rates for large larvae.³For suppression only.⁴See resistance statement under **General Directions for Use.**⁵ Does not include Western Flower Thrips.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
LEGUME VEGETABLES (SOYBEANS)			
Soybean	Corn Earworm Velvetbean Caterpillar Green Cloverworm Cabbage Looper Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Woollybear Caterpillar Cutworm species Bean Leaf Beetle Mexican Bean Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Three-Cornered Alfalfa Hopper Potato Leafhopper Thrips species ⁵ Soybean Aphid ⁴	0.015–0.025	1.92–3.20
	Armyworm ¹ Fall Armyworm ¹ Yellow-striped Armyworm ¹ Tobacco Budworm ³ Webworm species European Corn Borer Silverspotted Skipper Japanese Beetle (Adult) Blister Beetle species Stink Bug species Plant Bug species Grasshopper species	0.025–0.03	3.20–3.84
	Beet Armyworm ^{2,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite species ²	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Do not graze or harvest treated soybean forage, straw or hay for livestock feed.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 2.56 fl. oz./A (0.02 lb. a.i./A).
- Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pts./A) per season

¹Use higher rates for large larvae.²Suppression only.³See resistance statement under **General Directions for Use**.⁴Use lower rates for early season applications and/or lighter populations.⁵Does not include WesternFlower Thrips.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
LETTUCE (HEAD AND LEAF)			
	Alfalfa Looper Cabbage Looper Imported Cabbageworm Cutworm species Saltmarsh Caterpillar Green Cloverworm	0.015–0.025	1.92–3.20
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Southern Armyworm Corn Earworm Tobacco Budworm ³ European Corn Borer Flea Beetle species Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper species Leafhopper species Plant Bug species including Lygus species ³ Stink Bug species Meadow Spittlebug Aphid species ^{2,3} Whitefly species ^{2,3} Spider Mite species ²	0.02–0.03	2.56–3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 1 day of harvest.
- Do not apply more than 0.3 lb. a.i. (2.4 pts./A) per season.

¹For control of first and second instar only.

²Suppression only.

³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
ONION (BULB) AND GARLIC			
	Cutworm species Seedcorn Maggot (Adult) Onion Maggot (Adult) Leafminer species (Adult)	0.015–0.025	1.92–3.20
	Amyworm species ¹ Onion Thrips ³ Tobacco Thrips ³ Western Flower Thrips ^{2,3} Flower Thrips ^{2,3} Aphid species ² Plant Bug species Stink Bug species	0.02–0.03	2.56–3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Use the higher label rates as thrips population increases and avoid rescue situations.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage.
- Do not apply within 14 days of harvest. Do not apply more than 0.24 lb. a.i. (1.92 pts.)/A per season.

¹For control of the first and second instar only.²Suppression only.³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
PEANUT			
	Cutworm species Green Cloverworm Velvetbean Caterpillar Red-necked Peanut Worm Potato Leafhopper Three Cornered Alfalfa Hopper	0.015–0.025	1.92–3.20
	Corn Earworm Fall Armyworm ¹ Bean Leaf Beetle Southern Corn Rootworm (Adult) Vegetable Weevil Whitefringed Beetle (Adult) Stink Bug species Tobacco Thrips Grasshopper species	0.02–0.03	2.56–3.84
	Beet Armyworm ^{2,3} Soybean Looper ^{2,3} Lesser Cornstalk Borer ² Spider Mite species ² Aphid species ²	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.12 lb. a.i. (0.96 pts./A) per season.

¹Use higher rates for large larvae.

²Suppression only.

³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
POME FRUITS			
Apple Crabapple Loquat Mayhaw Oriental Pear Pear Quince	Leafroller species Omnivorous Leafroller Codling Moth Orange Tortrix Tufted Apple Budworm Oriental Fruit Moth Lesser Appleworm Green Fruitworm Tent Caterpillar species Webworm species Tentiform Leaf Miner species Apple Maggot (Adult) Cherry Fruit Fly species (Adult) Pear Sawfly Stink Bug species Leafhopper species Plum Curculio Japanese Beetle Tree Borer species Plant Bug species Periodical Cicada Apple Aphid Rosy Apple Aphid Spirea Aphid ¹ Pear Psylla ¹ San Jose Scale (fruit infestations only)	0.02-0.04	2.56-5.12

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds and IPM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gals. of water/per acre, but use higher volumes as appropriate for thorough coverage.
- Do not apply within 21 days of harvest.
- Do not apply more than 0.2 lb. a.i. (1.6 pts.)/A per year. Do not apply more than 0.16 lb. a.i. (1.28 pts.)/A per year post bloom.

¹Suppression only

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
STONE FRUITS			
Apricot Sweet and Tart Cherry Nectarine Peach Plum Chickasaw Plum Damson Plum Japanese Plum Plumcot Prune	Leafroller species Peach Twig Borer Oriental Fruit Moth Peachtree Borer species Green Fruitworm Tent Caterpillar species Codling Moth American Plum Borer Apple Maggot (Adult) Cherry Fruit Fly species (Adult) Pear Sawfly Plum Curculio Rose Chafer Japanese Beetle June Beetle Plant Bug species Stink Bug species Leafhopper species Thrips species Periodical Cicada Black Cherry Aphid	0.02-0.04	2.56-5.12

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold and IPM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 5 gals. of water/per acre, but use higher volumes as appropriate for thorough coverage.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.2 lb. a.i. (1.6 pts.)/A per year. Do not apply more than 0.16 lb. a.i. (1.28 pts.)/A per year post bloom.

Crop	Target Pest	Rate	
		lb. a.i./A	fl. oz./A
SUGARCANE			
	Mexican Rice Borer ¹ Sugarcane Borer ¹ Rice Stalk Borer ¹ Sugarcane Beetle (Adult) ² Sugarcane Aphid ³ Yellow Sugarcane Aphid ³ West Indian Crane fly Pygmy Mole Cricket	0.025-0.04	3.20-5.12

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gals. of water/A.
- Do not apply within 21 days of harvest.
- Do not apply more than 0.16 lb. a.i. (1.28 pts.)/A per season.

¹For control before the larva bores into the plant stalk.

²Suppression only of beetles active above ground.

³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
SUNFLOWER			
	Sunflower Beetle Cutworm species	0.015–0.025	1.92–3.20
	Sunflower Moth Banded Sunflower Moth Fall Armyworm ¹ Woollybear Caterpillar Spotted Cabbage Looper Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Stem Weevil (Adult) Head-Clipper Weevil (Adult) Japanese Beetle (Adult) Sunflower Maggot (Adult) Leafhopper species Meadow Spittlebug Stink Bug species Grasshopper species	0.02–0.03	2.56–3.84
	Beet Armyworm ^{2,3} Spider Mite species ²	0.03	3.84

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 45 days of harvest.
- Do not apply more than 0.12 lb. a.i. (0.96 pts.)/A per season. Do not apply more than 0.09 lb. a.i. (0.72 pts.)/A per season after bloom initiation.
- Do not apply as an ultra-low volume (ULV) spray.

¹Use higher rates for large larvae.²Suppression only.³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
TOBACCO (AIR DRIED); BURLEY TOBACCO AND FLUE-CURED TOBACCO			
	Tobacco Hornworm Tomato Hornworm Cabbage Looper Corn Earworm Cutworm species Tobacco Budworm ² Salt Marsh Caterpillar Armyworm species ¹ Webworm species Potato Tuberworm Tobacco Flea Beetle (Adult) Cucumber Beetle species (Adult) Blister Beetle species Vegetable Weevil (Adult) Japanese Beetle (Adult) Grasshopper species Tree Cricket species Katydid species Plant Bug species ³ Stinkbug species Tobacco Thrips species ² Tobacco Aphid species ^{2,3}	0.015-0.03	1.92-3.84

Remarks

- Apply as required by scouting, usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
- Do not apply within 40 days of harvest.
- Do not apply more than 0.09 lb. a.i. (0.72 pts.)/A per year.

¹For control of first and second instars only.

²Suppression only.

³See resistance statement under **General Directions for Use**.

Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
TREE NUTS			
Almond Beech Nut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (Hazlenut) Hickory Nut Macadamia Nut (Bush Nut) Walnut, Black Walnut, English (Persian)	Leafroller species Navel Orangeworm Codling Moth Filbertworm Peach Twig Borer Walnut Husk Fly species (Adult) Ants Plant Bug species Stink Bug species Chinch Bug Leaffooted Bug Walnut Aphid	0.02-0.04	2.56-5.12
Pecan	Hickory Shuckworm Pecan Casebearer species Pecan Weevil Pecan Aphid species Pecan Spittlebug Stink Bug species Pecan Phylloxera species	0.02-0.04	2.56-5.12

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gals. of water/per acre, but use higher rates as appropriate for thorough coverage.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.16 lb. a.i. (1.28 pts./A) per year. Do not apply more than 0.12 lb. a.i. (0.96 pts./A) per year post bloom.

NON-AGRICULTURAL USES			
Crop	Target Pests	Rate	
		lb. a.i./A	fl. oz./A
CONIFER AND DECIDUOUS TREES			
Plantations and Nurseries	Pine Tip Moth species Spruce Budworm Bagworm Tent Caterpillar species Leafroller species Gypsy Moth Webworm species Tussock Moth species Birch Leafminer Pine Sawfly species Sawfly species Pine Chafer Japanese Beetle May Beetle species June Beetle species Pine Colaspis Beetle European Elm Bark Beetle Leaf Beetle species Elm Leaf Beetle Pales Weevil Pine Weevil species Black Pine Weevil Pine Conelet Bug Spittlebug species Pine Leaf Chermid Balsam Woolly Aphid Balsam Twig Aphid Poplar Aphid species Pine Tortoise Scale Pine Needle Scale Mealybug species ¹	0.02-0.04	2.56-5.12

Remarks

- To control exposed foliage, flower, cone, seed and bark feeding insects, apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground equipment using sufficient water to obtain full coverage of target site. When applying by air, apply a minimum of 2 gals. of water/A.
- Do not apply more than 0.24 lb. a.i. (1.92 pts./A) per year

¹Suppression only.

Crop	Target Pest	Rate	
		lb. a.i./A	fl.oz./A
CONIFER AND DECIDUOUS TREES			
Seed Orchards	Coneworm species Seed Bug species Thrips species	See Remarks	See Remarks

Remarks

- For high volume sprayers, dilute 5.12 fl. oz. per 100 gals. of water and apply 5-10 gals. of finished spray per tree.
- For low volume sprayers, dilute 20 fl. oz. per 100 gals. of water and apply 100 gals. of finished spray/A.
- For aerial applications, apply 15 fl. oz./A in a minimum of 10 gals. finish spray/A.
- Do not apply more than 0.5 lb. a.i. (4 pts./A) per year.

Crop	Target Pest	Rates	
		lb. a.i./A	fl. oz./A
NON-CROPLAND (EXCLUDING PUBLIC LAND)			
	See Crop Outlets on this TAIGA™ Z INSECTICIDE label for target pest and rates.	See Crop Outlets	See Crop Outlets

Remarks

- Spray non-cropland adjacent to agricultural areas to control migratory insects, which may threaten crops.
- Follow General Use Directions, rates and spray recommendations found elsewhere in this label for the adjacent crop outlet and target pests.
- Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages.
- Repeat as necessary to maintain control.
- Do not exceed 0.2 lb. a.i. (1.6 pts.)/A per year.
- Do not graze livestock in treated areas.

Rate Conversion Chart

lb. a.i./A	fl. oz./A	pts./A	Treated Acres/gal.
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER THE MANUFACTURER OR SELLER SHALL BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW.

WARRANTY DISCLAIMER

THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. UNINTENDED CONSEQUENCES, INCLUDING

BUT NOT LIMITED TO CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER MANUFACTURER OR SELLER SHALL BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

Taiga™ is a trademark of Agriliance LLC.

Karate® Insecticide, Karate® Insecticide With Zeon™ Technology, Warrior® Insecticide, Warrior® Insecticide with Zeon™ Technology, E-Z Handler®, and the Syngenta logo are trademarks of a Syngenta Group Company

Ammo®, Capture®, Fury™, and Mustang® are trademarks of FMC Corporation

Asana® is a trademark of E. I. du Pont de Nemours & Co. (Inc.)

Baythroid®, DECIS® and SCOUT-XTRA® are trademarks of Bayer Crop Sciences.

Danitol® is a trademark of Valent.