

1381-211

8/22/2012

1/39



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF
CHEMICAL SAFETY
AND POLLUTION PREVENTION

Ms Alice Walker
Winfield Soutions LLC
P O Box 64589
St Paul MN 55164

Subject Grizzly Z Insecticide EPA Reg No 1381 211
 Date of Registrant Submission June 13 2012
 Decision 466695

AUG 22 2012

Dear Ms Walker

The labeling referred to above submitted in connection with registration under the Federal insecticide Fungicide and Rodenticide Act as amended is acceptable

Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment A stamped copy of the label is enclosed for your records If you have any questions you may contact Samantha Hulkower at (703) 603 0683 or Hulkower Samantha@epa gov

Sincerely

Mark Suarez
Product Manager (13)
Insecticide Branch
Registration Division (7505P)

Enclosure Copy of Label Stamped Accepted

RESTRICTED USE PESTICIDE
Due to Toxicity to Fish and Aquatic Organisms
 For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification

Grizzly[®] Z Insecticide

Active Ingredient

Lambda cyhalothrin

[1α(S) 3α(Z)] (±) cyano (3 phenoxyphenyl)methyl

3 (2 chloro 3 3 3 trifluoro 1 propenyl) 2 2

dimethylcyclopropanecarboxylate

Other Ingredients

Total

11.4%

88.6%

100.0%

Grizzly[®] Z Insecticide contains 1 lb of active ingredient per gal and is a capsule suspension

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you do not understand the label find someone to explain it to you in detail)

See additional precautionary statements complete storage and disposal and directions for use in booklet

FIRST AID	
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice Do not give any liquid to the person Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
If inhaled	<ul style="list-style-type: none"> Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment	
NOTE TO PHYSICIAN	
Contains petroleum distillate – vomiting may cause aspiration pneumonia Have the product container or label with you when calling a poison control center or doctor or going for treatment For 24 hour medical emergency assistance (human or animal) call 1 877 424 7452	


EPA Reg No 1381 211

EPA Est 070989 MO 001

Distributed By
 Winfield Solutions LLC
 P O Box 64589
 St Paul MN 55164 0589

Net Contents 2.5 gal ons
 Lot No _____
 1/0613/2

WINFIELD



AGRISOLUTIONS

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
WARNING/AVISO

May be fatal if swallowed Harmful if absorbed through skin Causes moderate eye irritation May cause allergic skin reactions Avoid contact with eyes skin or clothing Wash thoroughly with soap and water after handling and before eating drinking or using tobacco

Skin exposure may also result in a sensation described as a tingling itching burning or prickly feeling Onset may occur immediately to 4 hrs after exposure and may last 2-30 hrs without damage Wash exposed areas once with soap and water Relief from the skin sensation may be obtained by applying an oil based cream

Remove and wash contaminated clothing before reuse Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category F on an EPA chemical resistance category selection chart

Applicators and other handlers must wear

Long sleeved shirt and long pants

Chemical resistant gloves such as barrier laminate butyl rubber nitrile rubber or Viton \geq 14 mils

Shoes plus socks

Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate Do not reuse them Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

When handlers use closed systems enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Recommendations

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing
- As soon as possible wash thoroughly and change into clean clothing

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic organisms and toxic to wildlife For terrestrial uses do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark Do not apply when weather conditions favor drift from treated areas Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment wash water

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area

DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

This labeling must be in the possession of the user at the time of application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls

- Chemical resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal. Use label language appropriate for container size and type.

Nonrefillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Refillable container Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY Spill leak fire exposure or accident call
CHEMTREC 1 800-424 9300**

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals /A by air or 10 gals /A by ground unless otherwise specified in this label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, Grizzly[®] Z Insecticide may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

TANK MIX APPLICATION

When tank mixing with any other agricultural products, **always add Grizzly[®] Z Insecticide last**. Fill the tank with one half to two thirds volume of the mixing diluent. Make sure all other products are fully dispersed in the mixing diluent before adding the recommended rate of Grizzly[®] Z Insecticide to the tank. Add the remainder of the mixing diluent volume. It is recommended that mixing and spray equipment have continuous agitation for best results. Follow the precautions and limitations of the most restricted product in the tank mixture.

While Grizzly[®] Z Insecticide has good flexibility for tank mixing with other agricultural products, a jar test for physical compatibility is recommended for untried mixtures using proper ratios and mixing sequences of all ingredients to be included in the mixture.

Grizzly[®] Z Insecticide is an aqueous based formulation. It is recommended that no type of non emulsifiable oils be used in combination with Grizzly[®] Z Insecticide. If adjuvants are used, use only:
 Nonionic Surfactant (NIS) containing at least 75% surface agent, or
 Non phytotoxic Crop Oil Concentrate (COC) including once refined Vegetable Oil Concentrate (VOC)
 Methylated Sunflower Oils (MSO) containing a minimum of 17% emulsifier

Adjuvants other than NIS or COC may be used providing the product meets the following criteria

- 1 Contains only EPA exempt ingredients
- 2 Is non phytotoxic to the target crop
- 3 Is compatible in mixture (May be established through a jar test)
- 4 Is supported locally for use with Grizzly[®] Z Insecticide on the target crop through proven field trials and through university and extension recommendations

In addition the following may be used as diluents

Crop Oil Concentrate Methylated Sunflower Oils Urea Ammonium Nitrate

It is recommended that the following not be used in combination Grizzly[®] Z Insecticide as diluents or adjuvants

Non emulsifiable oils Diesel Fuel Straight Mineral Oil

CHEMIGATION Sprinkler Irrigation Application

Apply Grizzly[®] Z Insecticide at rates and timing described elsewhere in this label. As local recommendations differ, consult your local State Extension Service or other local experts for recommendations on adjuvant or diluent types (see TANK MIX APPLICATION) rates and mixing instructions. These recommendations should be proven through university and extension field trials to be effective with Grizzly[®] Z Insecticide applied by chemigation.

Check the irrigation system to insure uniform application of water to all areas. Thorough coverage of foliage is required for good control. Good agitation in the pesticide supply tank should be maintained prior to and during the entire application period.

Apply by injecting the recommended rate of Grizzly[®] Z Insecticide into the irrigation system using a metering device that will introduce a constant flow and by distributing the product to the target area in 0.1–0.2 acre inch of water. In general, use the least amount of water required for proper distribution and coverage. It is recommended that the product be injected into the main irrigation line ahead of a right angle turn in the line to insure adequate dispersion or mixing in the irrigation water. Once the application is completed, flush the entire irrigation and injection system with clean water before stopping the system.

In addition to the above recommendations, if application is being made during a normal irrigation set of a stationary sprinkler, the recommended rate of Grizzly[®] Z Insecticide for the area covered should be injected into the system only during the end of the irrigation set for sufficient time to provide adequate coverage and product distribution.

It is not recommended that Grizzly[®] Z Insecticide be applied through an irrigation system connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Use Precautions Sprinkler Irrigation Application

- A Apply this product only through (sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.
- B Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- D Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

- E A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise
- F The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- G The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- H The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- I The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops
- J The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- K Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock
- L Any alternatives to the above required safety devices must conform to the list of EPA approved alternative devices
- M Do not apply when wind speed favors drift beyond the area intended for treatment
- N Do not apply through chemigation systems connected to public water systems

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10 foot wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as but not limited to lakes reservoirs rivers permanent streams marshes or natural ponds estuaries and commercial fish farm ponds)

Only apply products containing Lambda cyhalothrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat

For guidance refer to the following publication for information on constructing and maintaining effective buffers

Conservation Buffers to Reduce Pesticide Losses Natural Resources Conservation Services USDA NRCS 2000 Fort Worth Texas 21 pp

www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf

Buffer Zone for Ground Application (groundboom overhead chemigation or airblast)

Do not apply within 25 feet of aquatic habitats (such as but not limited to lakes reservoirs rivers permanent streams marshes natural ponds estuaries and commercial fish ponds)

Buffer Zone for ULV Aerial Application

Do not apply within 450 feet of aquatic habitats (such as but not limited to lakes reservoirs rivers permanent streams marshes natural ponds estuaries and commercial fish ponds)

Buffer Zone for Non ULV Aerial Application

Do not apply within 150 feet of aquatic habitats (such as but not limited to lakes reservoirs rivers permanent streams marshes natural ponds estuaries and commercial fish ponds)

In the State of New York a 25 ft vegetated non cropped buffer strip untraversed by drainage tiles must be maintained between a treated field and a coastal salt marsh or permanent stream that drains into a coastal salt marsh for both aerial or ground application For aerial applications the 25 ft vegetated non cropped buffer strip for runoff protection would be part of the larger 150 ft buffer strip (or 450 ft buffer strip for ULV application) required for spray drift

SPRAY DRIFT REQUIREMENTS

Wind Direction and Speed

Only apply this product if the wind direction favors on target deposition

Do not apply when the wind velocity exceeds 15 mph

Temperature Inversion

Do not make aerial or ground applications into temperature inversions

Inversions are characterized by stable air and increasing temperatures with height above the ground Mist or fog may indicate the presence of an inversion in humid areas The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non ULV aerial application) according to ASAE (S572) definition for standard nozzles In conditions of low humidity and high temperatures applicators should use a coarser droplet size

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side immediately prior to application

For ground boom applications apply using a nozzle height of no more than 4 feet above the ground or crop canopy

For airblast applications turn off outward pointing nozzles at row ends and when spraying the outer two rows To minimize spray loss over the top in orchard applications spray must be directed into the canopy

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter

Flight speed and nozzle orientation must be considered in determining droplet size

Spray must be released at the lowest height consistent with pest control and flight safety Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety

When applications are made with a cross wind the swath will be displaced downwind The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind

SPECIFIC USE DIRECTIONS

AGRICULTURAL USES

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
ALFALFA AND ALFALFA GROWN FOR SEED			
	Alfalfa Caterpillar Cutworm species Army Cutworm Green Cloverworm Looper species Velvetbean Caterpillar Webworm species Leafhopper species Threecornered Alfalfa Hopper	0 015-0 025	1 92-3 20
	Armyworm Corn Earworm Fall Armyworm ¹ Western Yellow striped Armyworm Yellow striped Armyworm Alfalfa Weevil Bean Leaf Beetle (Adult) Blister Beetle species Clover Leaf Weevil species Clover Root Borer (Adult) Clover Root Curculio species (Adult) Clover Stem Borer (Adult) Cowpea Curculio (Adult) Cowpea Weevil (Adult) Cucumber Beetle species (Adult) Egyptian Alfalfa Weevil Grape Colaspis (Adult) Green June Beetle (Adult) Japanese Beetle (Adult) Mexican Bean Beetle Pea Weevil (Adult) Sweet Clover Weevil (Adult) Whitefringed Beetle species (Adult) Meadow Spittlebug Plant Bug species including Lygus species ³ Stink Bug species Alfalfa Seed Chalcid (Adult) Blue Alfalfa Aphid Cowpea Aphid Green Peach Aphid ³ Pea Aphid Spotted Alfalfa Aphid Thrips species ⁴ Grasshopper species	0 02-0 03	2 56-3 84
	Beet Armyworm ^{1 3} Blotch Leafminer ³ Spider Mites ²	0 03	3 84

Remarks

Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals /A by air or 10 gals /A by ground. When foliage is dense and/or pest populations are high 5-10 gals /A by air or 20 gals /A by ground and higher use rates are recommended. Use higher rates for increased residual control.

Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2-3 days following application. Avoid direct application to bee shelters.

Do not apply more than 0.03 lb a.i. (0.24 pts) /A per cutting.

Do not apply more than 0.12 lb a.i. (0.96 pts) /A per season.

Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay.

Use higher rates for large larvae.

² Suppression only.

See resistance statement under **General Directions for Use**.

Does not include Western Flower Thrips.

Crop	Target Pests	Rate	
		lb a.i./A	fl oz /A
CANOLA			
	Cutworm species Looper species Armyworm species Diamondback Moth Flea Beetle Cabbage Seedpod Weevil Lygus Bug Grasshoppers	0.015-0.03	1.92-3.84
	Cabbage Aphid	0.03	3.84

Remarks

- Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.

Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply a minimum of 2 gals of water/A.

Do not apply within 7 days of harvest.

- Do not apply more than 0.09 lb a.i. (0.72 pts) /A per year.

Crop	Target Pests	Rate	
		lb a.i./A	fl oz /A
CEREAL GRAINS			
Barley Buckwheat Oats Rye	Army Cutworm Cutworm species	0.015-0.025	1.92-3.2
	Armyworm Bird Cherry Oat Aphid ¹ Cereal Leaf Beetle English Grain Aphid ¹ Fall Armyworm Flea Beetle species Grasshopper species Hessian Fly ⁴ Orange Blossom Wheat Midge Russian Wheat Aphid ¹ Stink Bug species Yellowstriped Armyworm	0.02-0.03	2.56-3.84
	Grass Sawfly	0.025-0.03	3.20-3.84

Chinch Bug Corn Leaf Aphid ¹ Greenbug ^{1,3} Mite species ²	0.03	3.84
--	------	------

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre.

For chinch bug control, repeat applications at 3-5 day intervals, if needed. Grizzly[®] Z Insecticide may only suppress heavy infestations and/or migrations.

Greenbug is known to have many biotypes. Grizzly[®] Z Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.

Do not apply within 30 days of harvest.

Do not allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment. **Do not** feed treated straw to meat or dairy animals within 30 days after the last treatment.

Do not apply more than 0.06 lb a.i. (7.68 fl. oz. or 0.48 pts. of product) per acre per season.

¹ Best control is obtained before insects begin to roll leaves. Once crop has started to boot, Grizzly[®] Z Insecticide with Zeon.

Technology may provide suppression only. Higher rates and increased coverage will be necessary.

² Suppression only.

³ See Resistance statement under **General Directions for Use**.

⁴ Make applications when adults emerge.

Crop	Target Pest	Rate	
		lb a i /A	fl oz /A
CEREAL GRAINS			
Corn (At Plant) Field Corn Popcorn Seed Corn Sweet Corn	Wireworm species Cutworm species Seedcorn Maggot White Grub species Corn Rootworm Larvae Western Northern Southern Mexican Seedcorn Beetle Lesser Cornstalk Borer Red Imported Fire Ant ¹	0 005 lbs a i per 1000 feet of row ²	0 66 fl oz per 1000 feet of row ²

Remarks

Banded Applications – Apply at planting as a 5–7 inch T–band sprayed across the open seed furrow between the furrow openers and the press wheels or as a band application behind the press wheel

In–Furrow Applications – Apply into the seed furrow through spray nozzles or microtubes behind the planter furrow openers and in front of the press wheel

Apply a minimum of 3 gals finished spray/A

Do not harvest or graze livestock or cut treated crops for feed within 21 days of at plant application

Do not apply more than 0 09 lb a i (0 72 pts)/A per crop at plant

- For field corn popcorn and seed corn do not apply more than 0 12 lb a i /A per crop from at plant and foliar applications For sweet corn do not apply more than 0 48 lb a i /A per crop from at plant and foliar applications

¹Suppression only

² Lbs a i and fl oz /A of Grizzly® Z INSECTICIDE Applied at 0 66 fl oz /1000 ft of Row for Various Row Spacings						
Row Spacing	40	38	36	34	32	30
Linear Ft /A	13 068	13 756	14 520	15 374	16 335	17 424
Lbs a i /A	0 067	0 07	0 075	0 079	0 084	0 09
Fl oz /A	8 6	9 1	9 6	10 1	10 8	11 5

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
CEREAL GRAINS			
Corn (Foliar) Field Corn Popcorn Seed Corn	Cutworm species Western Bean Cutworm ¹ Corn Earworm ¹ Green Cloverworm Meadow Spittlebug	0 015-0 025	1 92-3 20
	Tobacco Budworm ^{1 4} European Corn Borer ¹ Southwestern Corn Borer ¹ Stalk Borer ¹ Hop Vine Borer ¹ Lesser Cornstalk Borer Armyworm ² Fall Armyworm ² Yellow striped Armyworm ² Webworm species Flea Beetle species Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Bean Leaf Beetle Cereal Leaf Beetle Japanese Beetle (Adult) Sap Beetle (Adult) Seedcorn Beetle Stink Bug species Grasshopper species Corn Leaf Aphid ³ Bird Cherry Oat Aphid ³ English Grain Aphid ³	0 02-0 03	2 56-3 84
	Mexican rice Borer ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹ Beet Armyworm ⁴ Southern Corn Leaf Beetle ³ Chinch Bug Green Bug ^{3 4}	0 03	3 84

Remarks

Apply as required by scouting or locally prescribed corn growth stages usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A.

- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small corn. Direct spray to the base of corn plants. Repeat applications at 3-5 day intervals if needed. GRIZZLY[®] Z Insecticide may only suppress heavy infestations and/or subsequent

migrations

For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 3.84 fl oz /A (0.03 lb a.i./A)

Do not apply within 21 days of harvest

- Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment. Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment
- Do not apply more than 0.12 lb a.i. (0.96 pts)/A per crop from at plant and foliar applications. Do not apply more than 0.06 lb a.i. (0.48 pts) after silk initiation. Do not apply more than 0.03 lb a.i. (0.24 pts) after corn has reached the milk stage (yellow kernels with milky fluid)

¹For control before the larva bores into the plant stalk or ear

²Use higher rates for large larvae

³Suppression only

⁴See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a.i./A	fl oz /A
CEREAL GRAINS			
Sweet Corn (Foliar)	Corn Earworm Armyworm ¹ Fall Armyworm ¹ Southern Armyworm ¹ Beet Armyworm ^{1,3} Yellow Striped Armyworm ¹ Cutworm species Western Bean Cutworm Webworm species European Corn Borer Southwestern Corn Borer Common Cornstalk Borer Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Japanese Beetle (Adult) Sap Beetle (Adult) Flea Beetle species Tarnished Plant Bug Stink Bug species Chinch Bug Aster Leafhopper Grasshopper species Aphid species ^{2,3} Spider Mite species ²	0.02–0.03	2.56–3.84
	Corn Silkyfly (Adult) ²	0.03	3.84

Remarks

- Apply as required by scouting or locally prescribed corn growth stages usually at intervals of 4 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds or other locally recommended methods and should be

targeted for control before insects enter the stalk or ear

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage and ears (if present) When applying by air apply in a minimum of 2 gals of water/A

For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program use a minimum of 3.2 fl oz /A (0.025 lb a.i./A)

- Do not apply within 1 day of harvest
 - Do not allow livestock to graze in treated areas or harvest treated corn forage as feed for meat or dairy animals within 1 day after last treatment Do not feed treated corn fodder or silage to meat or dairy animals within 21 days after last treatment
- Do not apply more than 0.48 lb a.i. (3.84 pts.)/A per crop from at plant and foliar applications

¹Use higher rates for large larvae

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
CEREAL GRAINS			
Rice	True Armyworm Fall Armyworm Yellow striped Armyworm Rice Water Weevil (Adult) Rice Stink Bug Chinch Bug Grasshopper species Leafhopper species Sharpshooter species Bird Cherry Oat Aphid Yellow Sugarcane Aphid Green Bug Mexican Rice Borer ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹ European Corn Borer ¹ Rice Seed Midge	0.025-0.04	3.20-5.12

Remarks

- Apply as required by scouting Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds Determine the need for repeat applications usually at intervals of 5-7 days by scouting
Grizzly[®] Z Insecticide can be safely used when propanil products are being used for weed control Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage When applying by air apply in a minimum of 2 gals of water (or a total carrier volume)/A but ensure sufficient volume is used to provide adequate coverage In addition adding an emulsifiable crop oil (e.g. 1 pt./A) when lower aerial application volumes are used is recommended to help improve coverage reduce evaporation and improve efficacy
For control of rice water weevil in dry seeded rice make a foliar application as indicated by scouting for the presence of adults and/or feeding scars usually within a time frame of 0-5 days after permanent flood establishment Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present Adults may also be treated at later stages of rice development to reduce overwintering populations
For control of rice water weevil in water seeded rice make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline Under conditions of prolonged migration into the field start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and

if needed apply a second application within 7–10 days of the first application Adults may also be treated at later stages of rice development to reduce overwintering populations

- California In addition to above directions for control of rice water weevil in water seeded rice Grizzly® Z Insecticide may be applied at the 1–3 leaf growth stage with the majority at the 2 leaf growth stage Adults are vulnerable on levees and in the water Larvae are vulnerable while feeding on the leaf prior to entering the soil Monitor for adults based upon field history and density of population Monitor field edges and levee areas for adults Treat in the following manner a) spray the inside perimeter of the field or b) spray the entire field
Green bug is known to have many biotypes Grizzly® Z Insecticide may only provide suppression If satisfactory control is not achieved with the first application of Grizzly® Z Insecticide a resistant biotype may be present Use alternate chemistry for control
 - Do not release flood water within 7 days of an application
Do not apply more than 0.12 lb a.i. (0.96 pts)/A per season
Do not apply more than 0.04 lb a.i. (0.32 pts)/A within 21 to 27 days of harvest
Do not apply within 21 days of harvest
 - Do not use treated rice fields for the aquaculture of edible fish and crustacea
 - Do not apply as an ultra low volume (ULV) spray
- ¹ For control before the larvae bores into the plant stalk

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
CEREAL GRAINS			
Wild Rice	Bird Cherry Oat Aphid Chinch Bug Fall Armyworm Grasshopper species Green Bug Leafhopper species Rice Stink Bug Riceworm Rice Water Weevil (Adult) Sharpshooter species True Armyworm Yellow Sugarcane Aphid Yellowstriped Armyworm	0.025-0.04	3.20-5.12
	European Corn Borer ¹ Mexican Rice Borer ¹ Rice Seed Midge ¹ Rice Stalk Borer ¹ Sugarcane Borer ¹	0.03-0.04	3.84-5.12

Remarks

Apply as required by scouting Timing and frequency of application should be based upon insect populations reaching locally determined economic thresholds Determine the need for repeat applications usually at intervals of 5-7 days by scouting
Grizzly® Z Insecticide can be safely used when propanil products are being used for weed control
Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage When applying by air apply in a minimum of 2 gals. of water (or a total carrier volume) per acre but ensure sufficient volume is used to provide adequate coverage In addition adding an emulsifiable crop oil (e.g. 1 pt per acre) when lower aerial application volumes are used is recommended to help improve coverage reduce evaporation and improve efficacy
For control of rice water weevil in dry seeded rice make a foliar application as indicated by scouting for the presence of adults and/or feeding scars usually within a time frame of 1-5 days after permanent flood establishment **Do not** exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present Adults may also be treated at later stages of rice development to reduce overwintering populations

For control of rice water weevil in water seeded rice make the first foliar application after pinpoint flood as indicated by scouting for the presence of adults and/or feeding scars usually when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and if needed apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.

California: In addition to above directions for control of rice water weevil in water seeded rice Grizzle Z Insecticide with Zeon Technology may be applied at the 1-3 leaf growth stage with the majority at the 2 leaf growth stage. Adults are vulnerable on levees and in the water. Larvae are vulnerable while feeding on the leaf prior to entering the soil. Monitor for adults based upon field history and density of population. Monitor field edges and levee areas for adults. Treat in the following manner: a) spray the inside perimeter of the field or b) spray the entire field.

Greenbug is known to have many biotypes. Grizzly[®] Z Insecticide may only provide suppression. If satisfactory control is not achieved with the first application of Grizzly[®] Z Insecticide a resistant biotype may be present. Use alternate chemistry for control.

For control of stem borers scout fields when rice growth is near panicle differentiation for early symptoms of damaging populations exhibited as discoloration (orange tan) around the junction of the leaf sheath and leaf blade which is caused by feeding of young larvae within the sheath. Applications must be made before larvae bore into rice stems. Make the first application at panicle differentiation to 2 inch panicle for partial control. Make the second application at boot to heading for maximum control. All rice varieties are susceptible to stem borer damage but Cocodrie and Priscilla are particularly susceptible.

Mixers/loaders supporting aerial applications to wild rice at a rate of 0.04 lb a.i. per acre and treating 1200 acres (or more) per day must wear dust mist respirator.

Do not release flood water within 7 days of an application.

Do not apply more than 0.12 lb a.i. (15.36 fl. oz. or 0.96 pts. of product) per acre per season.

Do not apply more than 0.04 lb a.i. (5.12 fl. oz. or 0.32 pt. of product) per acre within 21 to 27 days of harvest.

Do not apply within 21 days of harvest.

Do not use treated rice fields for the aquaculture of edible fish and crustacea.

Do not apply as an ultra low volume (ULV) spray.

¹ For control before the larvae bores into the plant stalk

Crop	Target Pests	Rate	
		lb a.i./A	fl oz /A
CEREAL GRAINS			
Sorghum (Grain)	Cutworm species Sorghum Midge	0.015-0.02	1.92-2.56 ⁶
	Armyworm Beet Armyworm ³ Fall Armyworm ¹ Yellow striped Armyworm ¹ Corn Earworm Webworm species European Corn Borer ² Southwestern Corn Borer ² Lesser Cornstalk Borer ² Flea Beetle species Stink Bug species Grasshopper species	0.02-0.03	2.56-3.84
	Mexican Rice Borer ² Rice Stalk Borer ² Sugarcane Borer ² Chinch Bug	0.03	3.84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of target location. When applying by air, apply in a minimum of 2 gals. of water/A. For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 5 day intervals if needed.
- For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of sorghum plants. Repeat applications at 3-5 day intervals if needed. Grizzly® Z Insecticide may only suppress heavy infestations and/or subsequent migrations.
- Do not apply more than 0.08 lb a.i. (0.64 pts.)/A per season.
- Do not apply more than 0.06 lb a.i. (0.48 pts.)/A per season after crop emergence.
- Do not apply more than 0.02 lb a.i. (0.16 pts.)/A per season once crop is in soft dough stage.
- Do not apply within 30 days of harvest.

¹Use higher rates for large larvae

²For control before the larva bores into the plant stalk

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
CEREAL GRAINS			
Wheat	Cutworm species	0.015-0.025	1.92-3.20
Wheat Hay	Army Cutworm		
Triticale	Armyworm	0.02-0.03	2.56-3.84 ¹
	Fall Armyworm		
	Yellow striped Armyworm		
	Flea Beetle species		
	Cereal Leaf Beetle		
	Stink Bug species		
	English Grain Aphid ¹		
	Russian Wheat Aphid		
	Bird Cherry Oat Aphid ¹		
	Grasshopper species		
	Hessian Fly ⁴		
	Orange Blossom Wheat Midge		
	Grass Sawfly	0.025-0.03	3.20-3.84
	Chinch Bug	0.03	3.84
	Greenbug ^{1,3}		
	Corn Leaf Aphid ²		
	Mite species ²		

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.

For chinch bug control, repeat applications at 3-5 day intervals if needed. Grizzly® Z Insecticide may only suppress heavy infestations and/or migrations.

- Greenbug is known to have many biotypes. Grizzly® Z Insecticide may provide suppression only. In this situation, a second application using an alternative chemistry may be needed.

Do not apply within 30 days of harvest

Do not allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment **Do not** feed treated straw to meat or dairy animals within 30 days after the last treatment

- Do not apply more than 0.06 lb a.i. (0.48 pts)/A per season

¹Best control is obtained before insects begin to roll leaves. Once wheat has started to boot Grizzly® Z Insecticide may provide suppression only. Higher rates and increased coverage will be necessary.

²Suppression only

³See resistance statement under **General Directions for Use**

⁴Make applications when adults emerge

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
COLE CROPS(Head and Stem Brassica)			
Broccoli Brussels Sprouts Cabbage Cavalo Broccoli Cauliflower Chinese Broccoli (gai lan) Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) Kohlrabi	Alfalfa Looper Cabbage Looper Imported Cabbageworm Southern Cabbageworm Cutworm species Cabbage Webworm	0.015–0.025	1.92–3.20
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Yellow striped Armyworm Corn Earworm Flea Beetle species Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper species Leafhopper species Plant Bug species including Lygus species ³ Stink Bug species Meadow Spittlebug Aphid species ^{2,3} Whitefly species ^{2,3} Thrips species ² Spider Mite species ²	0.02–0.03	2.56–3.84

Remarks

- Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A. Do not apply within 1 day of harvest.
- Do not apply more than 0.24 lb a.i. (1.92 pts)/A per season.

¹For control of first and second instar only

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
COTTON	Cutworm species Tobacco Thrips Soybean Thrips	0 015-0 02	1 92-2 56
	Lygus Bug species ³ Pink Bollworm Cabbage Looper Cotton Leafperforator Saltmarsh Caterpillar Cotton Leafworm Cotton Fleahopper	0 02-0 03	2 56-3 84
	Cotton Bollworm Tobacco Budworm ³ Boll Weevil Fall Armyworm Beet Armyworm ^{1 3} European Corn Borer Brown Stink Bug Green Stink Bug Southern Green Stink Bug Twospotted Spider Mite ² Cotton Aphid ^{2 3} Bandedwing Whitefly ^{2 3} Sweetpotato Whitefly ^{2 3}	0 025-0 04	3 20-5 12

Remarks

- Apply as required by scouting usually at intervals of 5 7 days Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds Apply with ground or air equipment using sufficient water to obtain full coverage of foliage Applications may also be made with equipment adapted and calibrated for ULV sprays Grizzly[®] Z Insecticide may be mixed with once refined vegetable oil and applied in a minimum of at least one qt of finished spray/A
- Under light bollworm/budworm infestation levels 0 02 lb a i /A may be applied in conjunction with intense field monitoring
- For boll weevil control spray on a 3 5 day schedule
- When applied according to label directions for control of cotton bollworm and tobacco budworm Grizzly[®] Z Insecticide also provides ovicidal control of unhatched Heliothine species eggs
- Do not apply within 21 days of harvest
Do not graze livestock in treated areas
- Do not apply more than 1 6 pts (0 2 lb a i)/A per season
Do not make more than a total of 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season Synthetic pyrethroid products include Ammo[®] Insecticide Asana[®] XL Insecticide Baythroid[®] Emulsifiable Pyrethroid Insecticide Capture[®] Insecticide/Miticide Danitol[®] 2 4 EC Spray Insecticide/Miticide Decis[®] Insecticide Fury[™] Insecticide Karate[®] Insecticide Karate[®] Insecticide with Zeon[™] Technology Mustang[®] Insecticide Scout X TRA[®] Insecticide Grizzly[®] Z Insecticide Warrior[®] Insecticide and Warrior Insecticide with Zeon[™] Technology

¹For control of first and second instar only

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a /A	fl oz /A
CUCURBIT VEGETABLES			
Chayote (fruit)	Armyworm species ¹	0 02 0 03	2 56 3 84
Chinese Waxgourd (Chinese preserving melon)	Blister Beetle species		
Citron Melon	Cabbage Looper		
Cucumber	Corn Earworm		
Gherkin	Cricket species		
Gourd (edible)	Cucumber Beetle species (adults)		
<i>Lagenaria</i> species – includes hyotan	Cutworm species		
cucuzza	Flea Beetle species		
<i>Luffa acutangula</i> <i>L. cylindrical</i> – Includes hechima Chinese okra	Grasshopper species		
<i>Momordica</i> species – includes balsam	June Beetle species		
apple balsam pear bitter melon	Leaffooted Bug		
Chinese cucumber	Leafhopper species		
Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i>) – includes true cantaloupe cantaloupe casaba crenshaw	Lygus Bug species ¹		
melon golden pershaw melon honeydew	Melonworm		
melon honey balls mango melon	Pickleworm		
Persian melon pineapple melon	Plant Bug species		
Santa Claus melon snake melon	Rindworm species complex		
Pumpkin	Saltmarsh Caterpillar		
Squash summer (<i>Cucurbita pepo</i> var <i>melopepo</i>) – includes crookneck squash	Squash Beetle		
scallop squash straightneck squash vegetable marrow zucchini	Squash Bug species		
Squash winter (<i>Cucurbita maxima</i> <i>C. moschata</i>) – includes butternut squash	Squash Vine Borer species		
calabaza hubbard squash (<i>C. mixta</i> <i>C. pepo</i>) – includes acorn squash spaghetti squash	Stink Bug species		
Watermelon – includes hybrids and/or varieties of <i>Citrullus lanatus</i>	Thrips species ^{1 2}		
	Tobacco Budworm ¹		
	Webworm species		
	Aphid species ¹	0 03	3 84
	Leafminer species ^{1 3}		
	Spider Mite species ³		
	Whitefly species ^{1 3}		

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all plant parts. When applying by air, apply in a minimum of 2 gals total solution per acre. When applying by ground, a minimum of 10 gals total solution per acre is recommended.

Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual.

Insects that bore or tunnel into leaves, vines, stems or fruit must be controlled before penetration. Only exposed insects (larvae and/or adults) can be controlled with foliar applications of Grizzly[®] Z Insecticide.

Do not apply more than 0.18 lb a.i. (23 fl. oz. or 1.44 pts. of product) per acre per season. Do not apply within 1 day of harvest.

¹ See Resistance statement under **General Directions for Use**

² Does not include Western Flower Thrips

³ Suppression only

Crop	Target Pests	Rate	
		lb a t /A	fl oz /A
GRASS FORAGE FODDER AND HAY			
Pasture and Rangeland Grass Grass Grown for Hay or Silage and Grass Grown for Seed	Army Cutworm Cutworm species Essex Skipper Range Caterpillar Striped Grass Looper	0 015 0 025	1 92 3 2
	Beet Armyworm Billbug species ³ Bird Cherry Oat Aphid ¹ Black Grass Bug Black Turfgrass Beetle (adult) Blue Stem Midge Cereal Leaf Beetle Chinch Bug Crane Fly species Cricket species English Grain Aphid ¹ Fall Armyworm Flea Beetle species Grass Mealybug Grass Sawfly (adult) Grasshopper species Green June Beetle (adult) Greenbug ^{1 2} Japanese Beetle (adult) Katydid species Leafhopper species Mite species ³ Russian Wheat Aphid ¹ Southern Armyworm Spittlebug species Stink Bug species Sugarcane Aphid Thrips species Tick species True Armyworm Webworm species Yellowstriped Armyworm	0 02 0 03	2 56 3 84

Remarks

Apply as required by scouting Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage When applying by air apply in a minimum of 2 gals total solution per acre When applying by ground a minimum of 7 gals total solution is recommended

Use higher application volumes and rates when foliage is dense pest populations are high larvae are large and/or weather conditions are adverse Use higher rates for longer residual

For chinch bug control Grizzly[®] Z Insecticide may only suppress heavy infestations and/or migrations In this situation a second application using an alternative chemistry may be needed

Greenbug is known to have many biotypes Grizzly[®] Z Insecticide may provide suppression only In this situation a second application using an alternative chemistry may be needed

Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application **Do not** cut grass to be dried and harvested for hay until 7 days after the last application

Grass grown for seed Straw and mature seed (seed screenings) may be used as feed 7 days after the last application Regrowth of grass grown for seed may be used for grazing cut for forage or cut to be dried and harvested for hay

Do not apply more than 0.03 lb a.i. (3.84 fl. oz. or 0.24 pts. of product) per acre per cutting for pastures, rangeland and grasses grown for seed. A minimum re-treatment interval (RTI) of 30 days is required for pastures and rangeland receiving 0.03 lb a.i./A which have not been cut between applications.
 Do not apply more than 0.09 lb a.i. (11.52 fl. oz. or 0.72 pts. of product) per acre per season.

- ¹ Best control is obtained before insects begin to roll leaves
- ² See Resistance statement under **General Directions for Use**
- ³ Suppression only

Crop	Target Pests	Rate	
		lb a.i./A	fl. oz./A
FRUITING VEGETABLES			
Tomato Tomatillo Peppers (bell and nonbell) Eggplant Ground cherry Pepino	Cabbage Looper Cutworm species Hornworm species	0.015–0.025	1.92–3.20
	Tomato Fruitworm Tobacco Budworm ³ Tomato Pinworm Beet Armyworm ^{1,3} Southern Armyworm ¹ Yellow striped Armyworm ¹ Fall Armyworm ¹ European Corn Borer ⁴ Leafminer species ² Colorado Potato Beetle ³ Flea Beetle species Grasshopper species Leafhopper species Aphid species ^{2,3} Whitefly species ^{2,3} Meadow Spittlebug Stink Bug species Plant Bug species Stalk Borer ⁴ Blister Beetle species Japanese Beetle (Adult) Pepper Weevil (Adult) ² Vegetable Weevil (Adult) Tomato Psyllid ^{2,3} Spider Mite species ² Thrips ⁵ Cucumber Beetle species (Adult)	0.02–0.03	2.56–3.84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
 Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
 Do not apply within 5 days of harvest.
 Do not apply more than 0.36 lb a.i. (2.88 pts.)/A per season.

- ¹For control of first and second instar only
- ²Suppression only
- ³See resistance statement under **General Directions for Use**
- ⁴For control before the larva bores into the plant stalk or fruit

⁵Does not include Western Flower Thrips

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
LEGUME VEGETABLES (Beans and Peas)			
Edible Podded (Only)	Cutworm species Green Cloverworm Imported Cabbageworm	0 015-0 025	1 92-3 20
<i>Canavalia gladiata</i> - sword bean	Saltmarsh Caterpillar Velvetleaf Caterpillar Mexican Bean Beetle		
<i>Canavalia ensiformis</i> - jackbean			
<i>Glycine max</i> - Soybean (immature seed)	Corn Earworm Painted Lady Butterfly (Larva) European Corn Borer Looper Species Western Bean Cutworm	0 02-0 03	2 56-3 84
Edible Podded Succulent Shelled or Dried Shelled	Tobacco Budworm ⁴ Armyworm ² Fall Armyworm ² Yellow-Striped Armyworm ² Western Yellow-Striped Armyworm ²		
<i>Phaseolus</i> species - includes field kidney lima navy pinto runner snap tepary and wax beans	Bean Leafskeletonizer Webworm species Leaftier species Alfalfa Caterpillar Stalk Borer ¹ Cucumber Beetle species (Adult) Corn Rootworm Beetle species (Adult)		
<i>Vigna</i> species - includes adzuki asparagus moth mung rice urd and yardlong beans black-eye pea catjang Chinese longbean cowpea Crowder pea and Southern pea	Flea Beetle species (Adult) Curculio and Weevil species ¹ (foliage and pod feeding adults and larvae) Blister Beetle species Bean Leaf Beetle Japanese Beetle (Adult) Leafhopper species Flea Hopper species Three-Cornered Alfalfa Hopper Meadow Spittlebug Stink Bug species Plant Bug species Including Lygus species ⁴		
<i>Pisum</i> species - includes dwarf edible-pod English field garden green snow and sugar snap peas	Grasshopper species Thrips species ^{4,5} Aphid species ⁴		
<i>Cajanus cajan</i> - Pigeon pea			

Crop	Target Pests	Rate	
		lb ai/A	fl oz/A
LEGUME VEGETABLES (Beans and Peas)			
(continued)	Beet Armyworm ^{3,4}	0.03	3.84
Succulent Shelled or Dried Shelled	Soybean Looper ^{3,4}		
	Lesser Cornstalk Borer ³		
	Leafminer species ^{3,4}		
<i>Vicia faba</i> – broadbean (favabean)	Whitefly species ^{3,4}		
	Spider Mite species ³		
Dried Shelled (Only)			
<i>Lupinus</i> species – includes grain sweet white and sweet white lupines			
<i>Cicer arietinum</i> – chickpea (garbanzo bean)			
<i>Cyamopsis tetragonoloba</i> – guar			
<i>Labiab pupureus</i> – Lablab bean (hyacinth bean)			
<i>Lens esculata</i> – Lentils			

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.

- For edible podded and succulent shelled legume vegetables, do not apply within 7 days of harvest.
- For dried shelled legume vegetables, do not apply within 21 days of harvest.
- Do not apply more than 0.12 lb ai (0.96 pts)/A per season.
- For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay.

¹For control before the larva bores into the plant stalk or pods.

²Use higher rates for large larvae.

³For suppression only.

⁴See resistance statement under **General Directions for Use**.

⁵Does not include Western Flower Thrips.

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
LEGUME VEGETABLES (SOYBEANS)			
Soybean	Corn Earworm Velvetbean Caterpillar Green Cloverworm Cabbage Looper Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Woollybear Caterpillar Cutworm species Bean Leaf Beetle Mexican Bean Beetle Western Corn Rootworm Beetle (Adult) Northern Corn Rootworm Beetle (Adult) Southern Corn Rootworm Beetle (Adult) Mexican Corn Rootworm Beetle (Adult) Three Cornered Alfalfa Hopper Potato Leafhopper Thrips species ⁵ Soybean Aphid ⁴	0 015-0 025	1 92-3 20
	Armyworm ¹ Fall Armyworm ¹ Yellow striped Armyworm ¹ Tobacco Budworm ³ Webworm species European Corn Borer Silverspotted Skipper Japanese Beetle (Adult) Blister Beetle species Stink Bug species Plant Bug species Grasshopper species	0 025-0 03	3 20-3 84
	Beet Armyworm ^{2 3} Soybean Looper ^{2 3} Lesser Cornstalk Borer ² Spider Mite species ²	0 03	3 84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Do not graze or harvest treated soybean forage, straw or hay for livestock feed.

Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.

For control of adult corn rootworm beetles (*Diabrotica* species) as part of an aerial applied corn rootworm control program, use a minimum of 2.56 fl. oz./A (0.02 lb. a.i./A).

Do not apply within 30 days of harvest. Do not apply more than 0.06 lb. a.i. (0.48 pts./A) per season.

¹Use higher rates for large larvae

²Suppression only

³See resistance statement under **General Directions for Use**

⁴Use lower rates for early season applications and/or lighter populations

⁵Does not include Western Flower Thrips

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
LETTUCE (HEAD AND LEAF)			
	Alfalfa Looper Cabbage Looper Imported Cabbageworm Cutworm species Saltmarsh Caterpillar Green Cloverworm	0.015–0.025	1.92–3.20
	Diamondback Moth ³ Armyworm Beet Armyworm ^{1,3} Fall Armyworm ¹ Southern Armyworm Corn Earworm Tobacco Budworm ³ European Corn Borer Flea Beetle species Japanese Beetle (Adult) Vegetable Weevil (Adult) Grasshopper species Leafhopper species Plant Bug species including Lygus species ³ Stink Bug species Meadow Spittlebug Aphid species ^{2,3} Whitefly species ^{2,3} Spider Mite species ²	0.02–0.03	2.56–3.84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
Do not apply within 1 day of harvest.
Do not apply more than 0.3 lb a.i. (2.4 pts.)/A per season.

¹For control of first and second instar only

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
ONION (BULB) AND GARLIC			
	Cutworm species Seedcorn Maggot (Adult) Onion Maggot (Adult) Leafminer species (Adult)	0.015–0.025	1.92–3.20
	Armyworm species ¹ Onion Thrips ³ Tobacco Thrips ³ Western Flower Thrips ^{2,3} Flower Thrips ^{2,3} Aphid species ² Plant Bug species Stink Bug species	0.02–0.03	2.56–3.84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Use the higher label rates as thrips population increases and avoid rescue situations.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water/A.

For thrips control by aerial application, the addition of 1% COC v/v, 1/4% NIS v/v or a silicone adjuvant (follow manufacturers use directions) may enhance the deposition of the spray and increase plant coverage.

Do not apply within 14 days of harvest. Do not apply more than 0.24 lb a i (1.92 pts)/A per season.

¹For control of the first and second instar only

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
PEANUT			
	Cutworm species Green Cloverworm Velvetbean Caterpillar Red necked Peanut Worm Potato Leafhopper Three Cornered Alfalfa Hopper	0 015-0 025	1 92-3 20
	Corn Earworm Fall Armyworm ¹ Bean Leaf Beetle Southern Corn Rootworm (Adult) Vegetable Weevil Whitefringed Beetle (Adult) Stink Bug species Tobacco Thrips Grasshopper species	0 02-0 03	2 56-3 84
	Beet Armyworm ^{2 3} Soybean Looper ^{2 3} Lesser Cornstalk Borer ² Spider Mite species ² Aphid species ²	0 03	3 84

Remarks

- Apply as required by scouting usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air apply in a minimum of 2 gals of water/A.
Do not apply within 14 days of harvest.
Do not apply more than 0 12 lb a i (0 96 pts)/A per season.

¹Use higher rates for large larvae

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
POME FRUITS			
Apple	Leafroller species	0 02-0 04	2 56-5 12
Crabapple	Omnivorous Leafroller		
Loquat	Codling Moth		
Mayhaw	Orange Tortrix		
Oriental Pear	Tufted Apple Budworm		
Pear	Oriental Fruit Moth		
Quince	Lesser Appleworm		
	Green Fruitworm		
	Tent Caterpillar species		
	Webworm species		
	Tentiform Leaf Miner species		
	Apple Maggot (Adult)		
	Cherry Fruit Fly species (Adult)		
	Pear Sawfly		
	Stink Bug species		
	Leafhopper species		
	Plum Curculio		
	Japanese Beetle		
	Tree Borer species		
	Plant Bug species		
	Periodical Cicada		
	Apple Aphid		
	Rosy Apple Aphid		
	Spirea Aphid ¹		
	Pear Psylla ¹		
	San Jose Scale (fruit infestations only)		

Remarks

- Apply as required by scouting usually at intervals of 5 or more days Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds and IPM recommendations
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area When applying by air apply in a minimum of 5 gals of water/per acre but use higher volumes as appropriate for thorough coverage
- Do not apply within 21 days of harvest
- Do not apply more than 0 2 lb a i (1 6 pts)/A per year Do not apply more than 0 16 lb a i (1 28 pts)/A per year post bloom

¹Suppression only

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
STONE FRUITS			
Apricot	Leafroller species	0.02-0.04	2.56-5.12
Sweet and Tart Cherry	Peach Twig Borer		
Nectarine	Oriental Fruit Moth		
Peach	Peachtree Borer species		
Plum	Green Fruitworm		
Chickasaw Plum	Tent Caterpillar species		
Damson Plum	Codling Moth		
Japanese Plum	American Plum Borer		
Plumcot	Apple Maggot (Adult)		
Prune	Cherry Fruit Fly species (Adult)		
	Pear Sawfly		
	Plum Curculio		
	Rose Chafer		
	Japanese Beetle		
	June Beetle		
	Plant Bug species		
	Stink Bug species		
	Leafhopper species		
	Thrips species		
	Periodical Cicada		
	Black Cherry Aphid		

Remarks

- Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold and IPM recommendations.
- Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 5 gals of water/per acre but use higher volumes as appropriate for thorough coverage.
- Do not apply within 14 days of harvest.
- Do not apply more than 0.2 lb a i (1.6 pts)/A per year. Do not apply more than 0.16 lb a i (1.28 pts)/A per year post bloom.

Crop	Target Pest	Rate	
		lb a i /A	fl oz /A
SUGARCANE			
	Mexican Rice Borer ¹	0.025-0.04	3.20-5.12
	Sugarcane Borer ¹		
	Rice Stalk Borer ¹		
	Sugarcane Beetle (Adult) ²		
	Sugarcane Aphid ³		
	Yellow Sugarcane Aphid ³		
	West Indian Crane fly		
	Pygmy Mole Cricket		

Remarks

Apply as required by scouting usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.

Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply a minimum of 2 gals of water/A.

- Do not apply within 21 days of harvest.
- Do not apply more than 0.16 lb a i (1.28 pts)/A per season.

¹For control before the larva bores into the plant stalk.

²Suppression only of beetles active above ground
³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a.i./A	fl oz/A
SUNFLOWER			
	Sunflower Beetle Cutworm species	0.015-0.025	1.92-3.20
	Sunflower Moth Banded Sunflower Moth Fall Armyworm ¹ Woollybear Caterpillar Spotted Cabbage Looper Painted Lady (Thistle) Caterpillar Seed Weevil (Adult) Stem Weevil (Adult) Head Clipper Weevil (Adult) Japanese Beetle (Adult) Sunflower Maggot (Adult) Leafhopper species Meadow Spittlebug Stink Bug species Grasshopper species	0.02-0.03	2.56-3.84
	Beet Armyworm ^{2,3} Spider Mite species ²	0.03	3.84

Remarks

Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
 Apply with ground or air equipment using sufficient water to obtain full coverage of sunflower heads and/or foliage. When applying by air, apply in a minimum of 2 gals. of water/A.
 Do not apply within 45 days of harvest.
 Do not apply more than 0.12 lb a.i. (0.96 pts)/A per season. Do not apply more than 0.09 lb a.i. (0.72 pts)/A per season after bloom initiation.
 Do not apply as an ultra low volume (ULV) spray.

¹Use higher rates for large larvae
²Suppression only
³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a l /A	fl oz /A
TOBACCO			
	Tobacco Hornworm	0 015-0 03	1 92-3 84
	Tomato Hornworm		
	Cabbage Looper		
	Corn Earworm		
	Cutworm species		
	Tobacco Budworm ²		
	Salt Marsh Caterpillar		
	Armyworm species ¹		
	Webworm species		
	Potato Tuberworm		
	Tobacco Flea Beetle (Adult)		
	Cucumber Beetle species (Adult)		
	Blister Beetle species		
	Vegetable Weevil (Adult)		
	Japanese Beetle (Adult)		
	Grasshopper species		
	Tree Cricket species		
	Katydid species		
	Plant Bug species ³		
	Stinkbug species		
	Tobacco Thrips species ²		
	Tobacco Aphid species ^{2 3}		

Remarks

Apply as required by scouting usually at intervals of 7 or more days Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold

Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage When applying by air apply in a minimum of 2 gals of water/A

Do not apply within 40 days of harvest

- Do not apply more than 0 09 lb a l (0 72 pts)/A per year

¹For control of first and second instars only

²Suppression only

³See resistance statement under **General Directions for Use**

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
TREE NUTS			
Almond	Leafroller species	0.02–0.04	2.56–5.12
Beech Nut	Navel Orangeworm		
Brazil Nut	Codling Moth		
Butternut	Filbertworm		
Cashew	Peach Twig Borer		
Chestnut	Walnut Husk Fly species (Adult)		
Chinquapin	Ants		
Filbert (Hazlenut)	Plant Bug species		
Hickory Nut	Stink Bug species		
Macadamia Nut (Bush Nut)	Chinch Bug		
Pistachio	Leaffooted Bug		
Walnut Black	Walnut Aphid		
Walnut English (Persian)			
Pecan	Hickory Shuckworm	0.02–0.04	2.56–5.12
	Pecan Casebearer species		
	Pecan Weevil		
	Pecan Aphid species		
	Pecan Spittlebug		
	Stink Bug species		
	Pecan Phylloxera species		

Remarks

- Apply as required by scouting usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold.
Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage or target area. When applying by air, apply in a minimum of 5 gals. of water/per acre, but use higher rates as appropriate for thorough coverage.
Do not apply within 14 days of harvest.
- Do not apply more than 0.16 lb a.i. (1.28 pts.)/A per year. Do not apply more than 0.12 lb a.i. (0.96 pts.)/A per year post bloom.

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
TUBEROUS AND CORM VEGETABLES (Potato Sweet Potato Yams and Related)			
Arracacha	Cutworm species	0 015 0 025	1 92 3 20
Arrowroot	Leafhopper species		
Artichoke (Chinese and Jerusalem only)	Saltmarsh Caterpillar		
Canna (edible)	Sweet Potato Hornworm		
Cassava (bitter and sweet)	Woollybear Caterpillar species		
Chayote (root)	Aphid species ¹	0 02 0 03	2 56 3 84
Chufa	Armyworm species ¹		
Dasheen	Blister Beetle species		
Ginger	Colorado Potato Beetle ¹		
Leren	Corn Earworm		
Potato	Cricket species		
Sweet Potato	Cucumber Beetle species (adults)		
Tanier	European Corn Borer		
Turmeric	Flea Beetle species (adults)		
Yam (bean and true)	Grasshopper species		
	Looper species ¹		
	Lygus Bug species ¹		
	Plant Bug species		
	Potato Psyllid		
	Potato Tuberworm		
	Stink Bug species		
	Sweet Potato Leaf Beetle (adults)		
	Sweet Potato Vine Borer		
	Thrips species ^{1 2}		
	Tortoise Beetle species		
	Webworm species		
	Weevil species (adults)		
	Leafminer species ^{1 3}	0 03	3 84
	Whitefly species ^{1 3}		
	Spider Mite species ³		

Remarks

Apply as required by scouting usually at intervals of 7 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of all plant parts. When applying by air, apply in a minimum of 2 gals. total solution per acre. When applying by ground, a minimum of 10 gals. total solution per acre is recommended.

Use higher application volumes and/or rates when foliage is dense, pest populations are high, larvae are large, weather conditions are adverse and/or as plant size increases. Use higher rates for longer residual.

Insects that bore or tunnel into leaves, vines, stems, tubers or corms must be controlled before penetration.

Only exposed insects (larvae and/or adults) can be controlled with foliar applications of Grizzly[®] Z Insecticide.

Do not apply more than 0.12 lb a.i. (15.36 fl. oz. or 0.96 pts. of product) per acre per season. Do not apply within 7 days of harvest.

¹ See Resistance statement under **General Directions for Use**

² Does not include Western Flower Thrips

³ Suppression only

NON-AGRICULTURAL USES

Crop	Target Pests	Rate	
		lb a i /A	fl oz /A
CONIFER AND DECIDUOUS TREES			
Plantations and Nurseries	Pine Tip Moth species	0.02-0.04	2.56-5.12
	Spruce Budworm		
	Bagworm		
	Tent Caterpillar species		
	Leafroller species		
	Gypsy Moth		
	Webworm species		
	Tussock Moth species		
	Birch Leafminer		
	Pine Sawfly species		
	Sawfly species		
	Pine Chafer		
	Japanese Beetle		
	May Beetle species		
	June Beetle species		
	Pine Colaspis Beetle		
	European Elm Bark Beetle		
	Leaf Beetle species		
	Elm Leaf Beetle		
	Pales Weevil		
	Pine Weevil species		
	Black Pine Weevil		
	Pine Conelet Bug		
Spittlebug species			
Pine Leaf Chermid			
Balsam Woolly Aphid			
Balsam Twig Aphid			
Poplar Aphid species			
Pine Tortoise Scale			
Pine Needle Scale			
Mealybug species ¹			

Remarks

To control exposed foliage, flower, cone, seed and bark feeding insects, apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

Apply with ground equipment using sufficient water to obtain full coverage of target site. When applying by air, apply a minimum of 2 gals. of water/A.

Do not apply more than 0.24 lb a i (1.92 pts.) /A per year.

¹Suppression only

Crop	Target Pest	Rate	
		lb a i /A	fl oz /A
CONIFER AND DECIDUOUS TREES			
Seed Orchards	Coneworm species Seed Bug species Thrips species	See Remarks	See Remarks

Remarks

For high volume sprayers dilute 5 12 fl oz per 100 gals of water and apply 5 10 gals of finished spray per tree

For low volume sprayers dilute 20 fl oz per 100 gals of water and apply 100 gals of finished spray/A

For aerial applications apply 15 fl oz /A in a minimum of 10 gals finish spray/A

Do not apply more than 0 5 lb a i (4 pts)/A per year

Crop	Target Pest	Rates	
		lb a i /A	fl oz /A
NON-CROPLAND (EXCLUDING PUBLIC LAND)			
	See Crop Outlets on this GRIZZLY [®] Z INSECTICIDE label for target pest and rates	See Crop Outlets	See Crop Outlets

Remarks

- Spray non-cropland adjacent to agricultural areas to control migratory insects which may threaten crops
Follow General Use Directions rates and spray recommendations found elsewhere in this label for the adjacent crop outlet and target pests
- Use highest labeled rates for dense/large foliage high insect populations and larger larval stages
Repeat as necessary to maintain control
- Do not exceed 0 2 lb a i (1 6 pts)/A per year
- Do not graze livestock in treated areas

Crop	Target Pest	Rates	
		lb a i /A	fl oz /A
NON-BEARING CITRUS ORCHARDS			
Calamondin Citron Citrus Hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sweet & sour) Pummelo Satsuma mandarin Tangerine Uniq Fruit Including all cultivars and/or hybrids of these	Asian Citrus Psyllid	0 03 0 04 lb a i /A	3 84 5 12 fl oz /A

Remarks

Apply as required by scouting Timing and frequency of applications should be based upon insects reaching locally determined economic thresholds and IPM recommendations

Apply with ground or air equipment using sufficient water to obtain full coverage of the foliage of higher volumes as appropriate for thorough coverage

- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Specific Use Restrictions

Apply to non bearing citrus trees only to within 1 year of harvest

Note To avoid possible illegal residues do not apply to citrus that will bear harvestable fruit within 12 months

Do not apply more than 0.24 lb a.i. (30.72 fl. oz. or 1.92 pts. of product) per acre per year

Not for use in citrus nurseries

Rate Conversion Chart			
lb a.i./A	fl oz/A	pts/A	Treated Acres/gal
0.015	1.92	0.12	66
0.02	2.56	0.16	50
0.025	3.20	0.20	40
0.03	3.84	0.24	33
0.04	5.12	0.32	25

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations on liability do not use the product and return it unopened to the Seller and the purchase price will be refunded

Grizzly® is a registered trademark of Winfield Solutions LLC

Karate® Insecticide Karate® Insecticide With Zeon™ Technology Warrior® Insecticide Warrior® Insecticide with Zeon™ Technology E Z Handler® and the Syngenta logo are trademarks of a Syngenta Group Company

Ammo® Capture® Fury™ and Mustang® are trademarks of FMC Corporation

Asana® is a trademark of E I du Pont de Nemours & Co (Inc)

Baythroid® DECIS® and SCOUT XTRA® are trademarks of Bayer Crop Sciences

Danitol® is a trademark of Valent