


1381-213

02/14/2007

111

 <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Reg. Number: 1381-213	Date of Issuance: FEB 14 2007
	Term of Issuance: Conditional	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)	Name of Pesticide Product: Daze 50 WP	
Name and Address of Registrant (include ZIP Code): Agriliance LLC P. O. Box 64089 St. Paul, MN 55164		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data. 2. Make the following label changes: <ol style="list-style-type: none"> a. Revise the EPA Registration Number to read, "EPA Reg. No. 1381-213. 3. Submit one copy of the revised final printed label for the record before you release the product for shipment. 4. Comply with Agency letter issued on this same date on one of your registered sources of active ingredient. 		
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	Date: FEB 14 2007	

2/11

page 2
EPA Reg. No. 1381-213

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

3/11



For Agricultural Use Only

DAZE™ 50WP

Cotton Defoliant Packed In Water Soluble Bags

ACTIVE INGREDIENT:	
Thidiazuron*	50%
INERT INGREDIENTS:	50%
TOTAL	100%

*N-phenyl-N'-1,2,3-thiadiazol-5-ylurea

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice
- Have person sip a glass of water if able to swallow
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency call toll free 1-877-424-7452.

Transportation or spill, call CHEMTREC 800-424-9300

EPA Reg. No. 1381-

Distributed By:
Agrilience, LLC
P.O. Box 64089
St. Paul, MN 55164-0089

EPA Est. No. 70989-AR-001
NET CONTENTS:
5 x 1 lb water soluble bag

PROP2-6-07

**ACCEPTED
with COMMENTS
in EPA Letter Dated:**

FEB 14 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No

1381-213

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. Do not contaminate food or feedstuffs.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

ENGINEERING CONTROLS STATEMENT

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240 (d)(6)].

USER SAFETY REQUIREMENTS STATEMENT

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains that product. A level, well-maintained vegetative buffer strip between areas to which this product is applied, and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination. This chemical has properties and characteristics associated with chemicals detected in ground water. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not store box under wet conditions. Handle box carefully when stored at temperatures of less than 50°F to avoid breakage of soluble bags.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty box into application equipment. Then dispose of empty box in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. DO NOT REUSE EMPTY CONTAINER

GENERAL INFORMATION

DAZE 50WP is designed to defoliate cotton (remove leaves and inhibit regrowth) to aid harvesting. DAZE 50WP is a wettable powder that can be applied by air or ground. Cotton leaf defoliation will take several days depending on weather conditions. Low temperatures require higher application rates and/or longer time to defoliate.

This box contains DAZE 50WP cotton defoliant in water soluble bags. Inner bags dissolve in water and contents disperse. After opening outer box, open foil liner, immediately dump entire unopened inner bags into the mix tank. Do not handle soluble bags or expose them to moisture, since this may cause breakage.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system.

Rainfall within 24 hours after application will reduce the effectiveness of DAZE 50WP. Use only freshly prepared sprays. Do not store spray mixture overnight.

Do not feed foliage from treated cotton plants or gin trash to livestock.

PLANT BACK INTERVALS	
CROP	INTERVAL
Small grain, sorghum, corn or root crops (except carrots, onions, and sugar beets)	2 weeks
Legumes (including alfalfa) or leafy vegetables (except lettuce and spinach)	2 months
Sugar beets	4 months
Carrots, onions, or spinach	9 months
Lettuce	9 months and only after deep plowing (12-15 inches)

- Do not use immature crops for food or feed.
- Do not allow spray drift to contact crops other than the target crop of mature cotton, or cotton which you desire to defoliate, as this product may injure or defoliate other crops.
- Particular care should be taken when applying DAZE 50WP adjacent to lettuce, citrus, or cantaloupe. Tank mixes with organophosphates may increase non-target crop phytotoxicity.
- Additionally, do not apply DAZE 50WP by air within one-half (1/2) mile of lettuce. Do not apply DAZE 50WP by ground equipment within 100 feet of lettuce.
- In addition, for citrus crops, do not apply DAZE 50WP by air when citrus in flush is within five (5) miles downwind of the point of application. Do not apply DAZE 50WP by ground when citrus in flush is within one-half (1/2) mile downwind of the point of application.

SPRAY DRIFT MANAGEMENT

Avoid spray drift to nearby crops as this product will cause modifications in plant growth. Small seeded crops such as lettuce, carrots, onions and spinach are especially sensitive. Plant injury or reduced yields will result.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

- 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

AERIAL DRIFT REDUCTION ADVISORY

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom length - For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application height - Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced windward. Therefore, on the up and down edges of the field, the applicator should compensate for the displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with the increasing drift potential (higher wind, smaller drops, etc.).

WIND

Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Thidiazuron products cannot be applied by air within ½ mile of lettuce or 5 miles downwind of the point of application when citrus is flush (burst of new growth, as in springtime) in the Rio Grande Valley of Texas. Ground applications are restricted to 100 feet away from lettuce and ½ mile downwind from the point of application for citrus. (In general, a decrease in droplet size or increase in wind speed at the time of application will result in risk to non-target organisms. Alternatively, if droplet size is coarser or wind speeds are lower, exposures due to drift would be reduced.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

General Use Rate Information (under normal conditions) for Cotton:

0.125 lbs. ai/A

Maximum application rate 0.3 lbs ai/A/year

Maximum of 2 applications, but not to exceed 0.3 lbs ai/A/year

General Use Rate Information (under variable conditions) for Cotton:

Maximum application rate of 0.2 lbs ai/A

Maximum application rate of 0.3 lbs ai/A/year

Maximum of 2 applications, but not to exceed 0.3 lbs ai/A/year.

For cutout and mature cotton under normal weather patterns, use Daze 50WP at 0.05 to 0.10 lb. active ingredient per acre. The 0.05 lb. rate is most effective when used in a tank mix with other cotton defoliant products. Increased rates of Daze 50 WP above 0.10 lb. of active ingredient per acre may be needed to defoliate and control regrowth during periods of rank growth/high fertilizer conditions, extreme weather conditions, such as extended periods of rain and/or low temperatures (60° – 65°F), and on full-season cotton varieties.

To Achieve an Application Rate of:	Use This Amount of Daze 50WP	At the Indicated Rate, One Gallon of Daze 50 WP Will Treat
0.07 lbs. ai/Acre	2.3 fl. oz./Acre	56 Acres
0.17 lbs. ai/Acre	5.4 fl. oz./Acre	23.5 Acres

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR

Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

TIME OF APPLICATION

Make application to mature cotton when all bolls to be harvested are mature and at least 5 days prior to harvest. Degree and speed of defoliation will depend on level of cotton growth activity, ambient temperature, humidity levels within 5 days after application and moisture content in cotton foliage.

Low temperatures (< 60°F) and drought stressed cotton can also adversely effect the level of defoliation and regrowth inhibition. If nightfall temperatures are < 60°F before and immediately after application, consideration should be given to use of a tank mix with another defoliant or desiccant approved for use on cotton and that is less affected by cooler temperatures.

USE OF ADJUVANTS (CROP OILS)

Tank mixing DAZE 50WP with EPA approved petroleum-based crop oils will help offset the effects of low temperatures, drought stress or low humidity levels typical in some areas (i.e. desert Southwest area). Refer to the crop oil label for specific application rates.

MIXING INSTRUCTIONS

1. Fill spray tank with 40-50% total amount of water required.
2. Start agitator.
3. Open foil bag and dump water-soluble packets. Use all contents of foil bags once opened.
4. Finish filling spray tank with balance of water with agitator running.

TANK MIX ADJUVANTS

Use of a surfactant or compatibility agent for tank mixes with other pesticides (especially organophosphates) will enhance tank clean out and increase defoliation. If applicator has no experience with a particular tank mix, a clear jar test of all ingredients should be performed. The jar test should duplicate the proposed tank mix in proper ratios and recommended order of addition.

Use only products that are exempt from tolerance under 40 CFR 180.1001.

Always follow the mixing instructions on the label of the appropriate adjuvant.

APPLICATION

Thorough coverage of cotton foliage is critical to maximize defoliation. Mix the labeled rate of DAZE 50WP in enough water to achieve complete coverage with spray equipment used. Spray mixture should be agitated constantly during application to insure consistent concentration. Apply in a minimum of 10 gallons per acre by ground and 2 gallons per acre by air.

DOSAGE

One box (5 x 1 lb. bags) of DAZE 50WP treats 12.5-25 acres (0.2-0.4 lbs. DAZE 50WP per acre).

Rate should be adjusted to current and future weather conditions. As noted in Timing section, higher rates are recommended at lower temperatures (< 65° F).

Two applications may be needed to defoliant rank cotton but do not exceed 0.6 lb./acre total.

BOTTOM DEFOLIATION FOR USE IN ARIZONA ONLY:

DAZE 50WP can be used to defoliate the lower third of cotton plants to promote air circulation in the field that will reduce boll rot disease pressures.

Apply 0.1 lb of DAZE 50WP per acre in a minimum of 10 gallons of water as a directed spray with ground equipment. Care should be taken to apply DAZE 50WP to prevent damage to the plants by the application equipment. Ground speed and appropriate dilution rate may need to be adjusted to minimize potential physical damage.

PRECONDITIONING FOR USE IN ARIZONA AND CALIFORNIA ONLY:

0.1 to 0.2 lb of DAZE 50WP per acre may be applied as a pre-conditioner 7-14 days before the use of another cotton defoliant. Apply in a minimum of 10 gallons of water per acre by ground equipment and 2 gallons of water per acre by air. Refer to other defoliant for any applicable precautions.

TANK MIX OF DAZE 50WP PLUS ETHEPHON 6 or PREP FOR BOLL OPENING:

A tank mix of 0.1 to 0.4 lb DAZE 50WP and 21 - 42 fluid ounces of ETHEPHON 6 (equivalent to 1.0 - 2.0 lbs active ingredient) per acre will enhance defoliation and accelerate boll opening. Use the higher rate of DAZE 50WP during cooler temperatures or if excessive regrowth is possible.

Do not use this tank mix before the optimal percentage of bolls in a field has matured. Premature application could result in a yield decrease. Certified cottonseed producers should confirm seed maturity before treatment.

Apply as a dilute spray as follows:

<u>Application Method</u>	<u>Gallons of Water/Acre</u>
Aerial	3-10
Ground	10-50

TANK MIX OF DAZE 50WP PLUS ETHEPHON 6 OR PREP FOR DEFOLIATION ENHANCEMENT:

Apply 0.1 to 0.4 lb of DAZE 50WP in a tank mix with 1/3 pint of ETHEPHON 6 (equivalent to 0.25 lb active ingredient per acre) will increase defoliation but without undesirable accelerated boll opening. Ethephon may still be applied subsequently for boll opening. Users should refer to ETHEPHON 6 label for any seasonal application limits for all ethephon use on cotton. Use the higher rate of DAZE 50WP during cooler temperatures or if excessive regrowth is possible.

TANK MIX OF DAZE 50WP PLUS DEF 6² OR DAZE 50WP PLUS FOLEX 6EC³:

Defoliation and regrowth inhibition will be improved during periods of cool weather (<60°F) with a tank mix of 0.1 to 0.4 lb of DAZE 50WP and 0.5 to 2.0 pints of Def⁶ 6 (or Folex⁶ 6EC) per acre. Apply tank mix in a minimum of 10 gallons of water per acre by ground and 5 gallons of water per acre by air.

Use the higher rate of DAZE 50WP if excessive regrowth is expected due to high soil moisture and/or higher temperatures.

Use the lower rate of Def⁶ 6 or Folex⁶ when nighttime temperatures remain above 65°F. To avoid plant desiccation, do not apply maximum rates of DAZE 50WP and phosphate based defoliant.

When mixing the tank mix spray solution, add the DAZE 50WP first (see Mixing Instructions) and then add the Def⁶ 6 or Folex⁶ after the DAZE 50WP has completely dispersed.

10/11

If a second application of this tank mix is required, do not exceed a total DAZE 50WP application of 0.6 lb per acre.

Refer to Def[®] 6 or Folex[®] labeling for additional use directions and cautions when using tank mixtures of DAZE 50WP and those products.

DO NOT USE TANK MIX OF DAZE 50WP PLUS DEF[®] 6 OR FOLEX[®] IN THE RIO GRANDE VALLEY COUNTIES OF STARR, HIDALGO, WILLACY, AND CAMERON.

IMPORTANT CLEANOUT INSTRUCTIONS

All spray equipment components containing spray solutions with DAZE 50WP should be cleaned as soon as possible after application. Care should be take to clean ALL components of equipment including (but not limited to) mix tanks, pumps, transfer lines, application tanks, sumps, booms and nozzles.

Timely flushing of entire system while still wet with a commercial grade tank cleaner is strongly advised.

If spray solution is allowed to dry in any equipment component, small quantities may carryover and be released in subsequent applications and damage other crops. If spray residue is allowed to dry, flooding system with a dilute commercial tank cleaner and allowing to soak for several days will be necessary before using equipment again.

SEQUENTIAL APPLICATION PRECAUTION

Application of DAZE 50WP to cotton subsequent to the application of a defoliant or desiccant spray that contains no thidiazuron will result in reduced activity.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER THE MANUFACTURER OR SELLER SHALL BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW.

WARRANTY DISCLAIMER

THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER MANUFACTURER OR SELLER SHALL BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT

11/11

DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

¹Prep is a trademark of Aventis Group

²Def 6 is a registered trademark of Bayer Corporation, Agricultural Chemical Division.

³Folex 6EC is a registered trademark of Aventis CropScience

DAZE is a trademark of Agrilience, LLC