

1381-216

05/08/2007

1/40

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOTICE OF PESTICIDE:

Registration
 Reregistration

MAY 8 2007

EPA Reg. Number: 1381-216

Term of Issuance: Conditional

Name of Pesticide Product: Pendant 3.3 EC Herbicide

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Within 18 months of the date of this registration, submit one-year storage stability and corrosion characteristics study for review and acceptance.

3. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 1381-216."
- b. Under "First Aid - If swallowed";

Replace "Have a person sip a glass if able to swallow" with "Do not give **any** liquid to the person." and "Do not give anything by mouth to an unconscious person."

c. Under "Environmental Hazards"

Add "or rinsate" after "washwaters."

d. Add the following section on Non-Target Species Concerns under "General Information":

Non-Target Species Concerns

If non-target plant species occur in proximity to the application site for bearing citrus fruit trees, bearing nut trees, carrots or mint, the following mitigation measures are required:

- o If applied by air, leave untreated buffer zones of 200 feet. Must use straight stream nozzles (such as D-6 to D-8). Wind can be no more than 8 mph, and release height must be 15 feet or less.
- o If applied by ground, leave untreated buffer zones of 80 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE

fine to medium/coarse nozzles. To determine whether your county has an endangered species, consult the website <http://www.epa.gov/espp/usa-map.htm>. Endangered Species Bulletins may also be obtained from Extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

- e. Under "Storage and Disposal";

Change "Storage" to "Pesticide Storage."

- f. Under "Use with other Products (Tank Mixes)";

Move this section to "Warranty Disclaimer." Also add "To the fullest extent permitted by law," before "the liability of Agrillance, LLC, shall in . . . purchased price of this product."

- g. Under "Chemigation – NonBearing Fruit and Nut Crops and Vineyards";

- o Add "follow other requirements specified in the Chemigation section of the label."
- o Delete "- B" on line 4 of item 2.
- o Move "C" to the center of the nominator on the equation on line 3 of item 4.
- o Delete "= S" on line 5 of item 4.

- h. Under "Peas";

Remove "It is a violation of . . . with this labeling." It appears that it is a general labeling requirement, it needs to be placed under "Directions for Use."

- i. Under "Rice"

Change " or disposal of wastes" to "equipment washwaters or rinsate" at the end of line 9.

- j. Under "Delayed (Late) Preemergence Applications - Rice";

Add a "period" at the end of item 6.

- k. Under "Soybeans";

"Round-up Ready" is not reference #5 at page 37 of this label.

- l. Add a line between paragraphs for easy reading and comprehension.

Submit one copy of the revised final printed label for the record before you release the product for shipment. If the above conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

CONCURRENCES

SYMBOL <	7505C	Final						
SURNAME <	Bien	<i>[Signature]</i>						
DATE <	5-02-07	5-8-07						

4/40
1



Pendant 3.3 EC Herbicide

FOR USE IN SELECTED CROPS

ACTIVE INGREDIENT:		
Pendimethalin (N-(1-ethylpropyl)-3, 4-dimethyl-2,6-dinitrobenzenamine).....	37.4%	
INERT INGREDIENTS*	62.6%	
TOTAL	100.0%	

(1 gallon contains 3.3 pounds of pendimethalin)
*Contains aromatic naphtha

KEEP OUT OF REACH OF CHILDREN

CAUTION! / ¡PRECAUCION!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

In case of an emergency endangering life or property involving this product, call day or night 800-832-HELP.

FOR USE IN COTTON, DRY BULB ONIONS, DRY BULB SHALLOTS, EDIBLE BEANS, CORN (FIELD, SEED, SWEET), FORAGE LEGUMES, GARLIC, GRAIN SORGHUM, NONBEARING FRUIT, NUT CROPS AND VINEYARDS; PEANUTS, PEAS, POPCORN, POTATOES, RICE, SOYBEANS, SUGARCANE, SUNFLOWERS, SWEET LUPINES, AND TOBACCO

PENDANT 3.3 EC SHOULD BE USED IN ACCORDANCE WITH THE DIRECTIONS IN THIS BOOKLET AND IN SUPPLEMENTAL LABELS, FOR REGISTERED USES NOT INCLUDED IN THIS BOOKLET, AVAILABLE THROUGH LOCAL DEALERS. READ ALL DIRECTIONS CAREFULLY BEFORE USING.

Read all information on pages 2-9 before referring to specific crop use.

FOR CONTROL OF MOST ANNUAL GRASSES AND CERTAIN BROADLEAF WEEDS

FIRST AID

If swallowed, Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.
If in eyes, Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses. If present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin, Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.
NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent, vomiting should be induced only under professional supervision. For additional information in case of emergency call toll free 1-877-424-7452.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION!

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

See Additional Precautionary Statements Inside Booklet

EPA Reg. No. 1381-

Distributed By:
Agrilience, LLC
P.O. Box 64089
St. Paul, MN 55164-0089

MAY 8 2007

ACCEPTED
with COMMENTS
in EPA Letter Dated

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended for the pesticide
registered under EPA Reg. No.
1381-216

EPA Est. No. 241-MO-1

Net Contents: 2.5 gallons
prop1-03-07

Table of Contents

	Page		Page		Page
Ingredients	1	Forage Legumes	17	Soybeans	28
First Aid Statement	1	General Instructions	17	General Instructions	28
Precautionary Statements	1	Use Methods and Timings	17	Use Methods and Timings	28
Environmental Hazards	3	Use Rates	17	Use Rates	29
Directions for Use	3	Garlic, Dry Bulb Onions and Shallots	17	Tank Mix and Sequential Programs	29
Agricultural Use Requirements	3	General Instructions	17	PENDANT 3.3 EC plus PURSUIT	29
Storage and Disposal	3	Use Directions for Mineral Soils	18	PENDANT 3.3 EC followed by	
Disclaimer	3	Use Directions for Muck Soils		RAPTOR® herbicide	29
General Information	4	(Onions Only)	18	PENDANT 3.3 EC plus SCEPTER®	
Mixing Instructions	4	Grain Sorghum	19	herbicide	29
General Application Instructions	5	CULTI-SPRAY™		PENDANT 3.3 EC followed by	
Rates	5	(Postemergence Incorporated)	19	SCEPTER® OT herbicide	30
Soil Textures	5	Early Postemergence	19	PENDANT 3.3 EC plus Command®	30
Chemigation	5	Nonbearing Fruit and Nut Crops		PENDANT 3.3 EC plus Canopy ¹	30
Weeds Controlled	6	and Vineyards	20	PENDANT 3.3 EC plus Dual	30
Spraying Instructions	8	General Instructions	20	PENDANT 3.3 EC plus Lasso	31
Incorporation Instructions	9	Use Methods and Timings	20	PENDANT 3.3 EC plus Lorox	31
Applications with Liquid Fertilizers	10	Use Rates	21	PENDANT 3.3 EC plus Lorox Plus ¹	31
Applications with Dry Bulk Fertilizers	10	Tank Mix Programs	21	PENDANT 3.3 EC plus Preview ¹	31
Cultural Practices Following Applications	11	Peanuts	21	PENDANT 3.3 EC plus	
Follow Crop Restrictions	11	General Instructions	21	Sencor/Lexone	31
Corn, Field	11	Use Methods and Timings	22	Special Weeds	31
General Instructions	11	Use Rates	22	Shattercane	32
Use Methods and Timings	12	Tank Mix Programs	22	Woolly Cupgrass	32
Use Rates	12	PENDANT 3.3 EC plus PURSUIT®		Red rice/itchgrass	32
CULTI-SPRAY™		herbicide	22	Rhizome Johnsongrass	32
(Postemergence Incorporated)	12	PENDANT 3.3 EC plus Dual	22	Sugarcane (except Hawaii)	33
Corn, Sweet, Seed and Popcorn	13	PENDANT 3.3 EC plus Vernam ⁵	22	General Instructions	33
General Instructions	13	Peas	22	Use Methods and Timings	33
Use Methods and Timings	13	General Instructions	22	Use Rates	33
Use Rates	13	Use Methods and Timings	22	Tank Mix Programs	33
Restrictions for Postemergence		Use Rates	23	Sugarcane (for Hawaii)	33
Applications	13	Potatoes	23	General Instructions	33
Cotton	14	General Instructions	23	Use Methods and Timings	33
General Instructions	14	Use Methods and Timings	23	Use Rates	33
Use Methods and Timings	14	Use Rates	23	Tank Mix Programs	33
Use Rates	14	Tank Mix Programs	23	Sunflowers	33
Tank Mixes and Sequential Programs	15	PENDANT 3.3 EC plus Eptam	24	General Instructions	33
PENDANT 3.3 EC plus fluometuron		PENDANT 3.3 EC plus Lorox ¹	24	Use Methods and Timings	34
(i.e. Cotoran ²)	15	PENDANT 3.3 EC plus		Use Rates	34
PENDANT 3.3 EC plus prometryn		Sencor ⁶ /Lexone ¹	24	Tank Mix Programs	34
(i.e. Caparol ²)	15	PENDANT 3.3 EC plus Eptam plus		PENDANT 3.3 EC plus Eptam	34
PENDANT 3.3 EC plus diuron		Sencor/Lexone	24	No-Till Sunflowers	35
(i.e. Karmex ¹)	15	PENDANT 3.3 EC plus Matrix ¹	25	Tobacco	35
PENDANT 3.3 EC plus Zorial ³	15	Rice	25	General Instructions	35
Rhizome Johnsongrass	16	General Instructions	25	Use Methods and Timings	35
Edible Beans [(Dry, Lima, Snap,		Early Postemergence Applications	25	Use Rates	36
Chickpeas (Garbanzo beans),		PENDANT 3.3 EC plus propanil	25	Regional Map for Rate Determination	36
Southern Peas (Cowpeas)] and		PENDANT 3.3 EC plus Arrosolo ⁵	25	Footnotes	36
Sweet Lupines	16	PENDANT 3.3 EC plus propanil plus			
General Instructions	16	Londax ¹	26		
Use Methods and Timings	16	PENDANT 3.3 EC plus Facet ⁷ 75 DF	26		
Use Rates	16	Delayed Preemergence Applications	27		
Tank Mix Programs	16	PENDANT 3.3 EC alone	27		
PENDANT 3.3 EC plus Dual	17	PENDANT 3.3 EC plus glyphosate	28		
PENDANT 3.3 EC plus Eptam ⁵	17	PENDANT 3.3 EC plus Facet 75 DF	28		
PENDANT 3.3 EC plus Lasso,		PENDANT 3.3 EC plus Bolero ⁹ EC	28		
Micro-Tech ⁴ or Partner ⁴ WDG	17	Spraying Instructions	28		

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or vitron ≥ 12 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [(40 CFR 170.240)(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

Observe all cautions and limitations in this label and the labels of products used in combination with **PENDANT 3.3 EC**. The use of **PENDANT 3.3 EC** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide protection.

DO NOT use in manufacturing products for application to turf and ornamentals.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated areas if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or vitron ≥ 12 mils.
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: **DO NOT STORE BELOW 40°F**. Extended storage at temperatures below 40°F can result in the formation of crystals on the bottom of the container. If crystallization does occur, store the container on its side at room temperature (70°F) and rock occasionally until crystals re-dissolve.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: For Five Gallons and Under: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For Bulk and Mini-Bulk: Return empty container to point of purchase for repackaging or recycling.

Uses with other Products (Tank Mixes)

If this product is used in combination with any other product except as specifically recommended in writing by Agrilience, LLC, then Agrilience, LLC shall have no liability for any loss, damage or injury arising out of its use in any such combination not so specifically recommended. If used in combination recommended by Agrilience, LLC, the liability of Agrilience, LLC shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Agrilience, LLC product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

GENERAL INFORMATION

Special Precautions

PENDANT 3.3 EC herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Destroy existing weeds before applying **PENDANT 3.3 EC** (except as recommended in specific postemergence combination treatments). Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Over-application can result in crop stand loss, crop injury, or soil residues.

Uneven application or improper soil incorporation can decrease weed control or cause crop injury. Soil incorporation deeper than recommended can reduce weed control.

Seedling diseases, cold weather, excessive moisture, shallow or deep planting, low or high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of crop damage from **PENDANT 3.3 EC**. Under these conditions, crop yields can be reduced.

Use of **PENDANT 3.3 EC** herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with the mitotic inhibiting mode of action. Other herbicides with the mitotic inhibiting mode of action include other dinitroaniline herbicides such as Treflan¹¹, TRI-4^e herbicide and Sonolan¹¹. If naturally occurring mitotic inhibiting resistant biotypes are present in a field, **PENDANT 3.3 EC** and/or any other mitotic inhibiting mode of action herbicide should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control.

*A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants. See your Agrilience, LLC representative for additional information.

MIXING INSTRUCTIONS

- 1. Fill tank one-half to three-quarters full with clean water or liquid fertilizer and agitate. Prior to mixing **PENDANT 3.3 EC** or **PENDANT 3.3 EC** tank mixtures in liquid fertilizer, refer to appropriate label sections (see table of contents for page number) for recommended uses in liquid fertilizer, application instructions, and compatibility determinations.

2. PENDANT 3.3 EC Alone

When using **PENDANT 3.3 EC** alone, add **PENDANT 3.3 EC** to the partially-filled tank while agitating and then fill the remainder of the tank with water or liquid fertilizer.

3. PENDANT 3.3 EC Tank Mixtures

Add the tank mixture ingredients in the order listed below prior to adding **PENDANT 3.3 EC**. (For tank mixtures with Butyrac 200, Gramoxone Extra or Roundup see mixing instructions at the end of this section):

(a) Wettable powder formulations

Make a slurry of the wettable powder in water (1 part WP + 2 parts water). Add the slurry slowly into the partially-filled tank while agitating. If tank mixing more than one wettable powder formulation, add the first one while agitating. Continue agitation and add second wettable powder only after complete mixing of the first wettable powder has occurred.

(b) Dry flowable/water dispersible granule formulations

Add the granules to the partially-filled tank while agitating. Make a slurry of the granules in water before adding to liquid fertilizer.

(c) Flowable formulations

Add the flowable to the partially-filled tank while agitating.

(d) Water soluble concentrate formulations

Add the water soluble concentrate to the partially-filled tank while agitating.

(e) Emulsifiable concentrate formulations

Add the emulsifiable concentrate to the partially-filled tank while agitating.

After complete mixing, add **PENDANT 3.3 EC** to the tank.

(f) NOTE: For tank mixes including Butyrac⁴, Gramoxone⁶ Extra or Roundup⁵

After complete mixing of **PENDANT 3.3 EC**, continue filling the sprayer with water and add Butyrac or Gramoxone Extra or Roundup near the end of the filling process.

If Gramoxone Extra is included in the tank mixture, add 8 oz. of Ortho X-77 spreader or similar non-ionic surfactant per 100 gallons of total spray mixture as the last ingredient in the tank.

Fill the remainder of the tank mix with water or liquid fertilizer while agitating.

- 4. MAINTAIN CONTINUOUS AGITATION WHILE ADDING HERBICIDES AND UNTIL SPRAYING IS COMPLETED.** If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

GENERAL APPLICATION INSTRUCTIONS

Uniformly apply recommended **PENDANT 3.3 EC** or **PENDANT 3.3 EC** tank mixture treatments in 10 or more gallons of water or 20 or more gallons of liquid fertilizer per acre with ground equipment. Use higher gallonage for fields with heavy weed infestations or excessive crop residues. Refer to Applications with Liquid Fertilizers section (see table of contents for page number) for liquid fertilizer recommended uses, application instructions and compatibility determinations. Apply in 5 or more gallons of water per acre with aircraft. Refer to Spraying Instructions section (see table of contents for page number) for spraying directions. Refer to Applications with Dry Bulk Fertilizers section (see table of contents for page number) for **PENDANT 3.3 EC**/dry bulk fertilizer applications.

RATES

Follow all recommendations in specific crop sections of this label regarding rates per acre of **PENDANT 3.3 EC** and herbicides used in combination with **PENDANT 3.3 EC**. When a rate range is recommended, unless other directions are given in a crop section, the high rate of the range should be used if heavy weed populations are anticipated or if extensive crop residues were present prior to seedbed preparation.

SOIL TEXTURES

The rate tables for recommended **PENDANT 3.3 EC** treatments for each crop refer to coarse, medium, and fine soils. Soil type should be determined since rates for **PENDANT 3.3 EC** treatments will change with soil texture. The following table lists soil types for coarse, medium and fine soils:

COARSE	MEDIUM	HIGH
sands	sandy clay loams*	silty clay loams*
loamy sands	sandy clays	silty clays
sandy loams	loams	clay loams
	silt loams	clays
	silts	

*Sometimes considered transitional soils and may be classified as either medium or fine textured soils.

PENDANT 3.3 EC is not recommended for use on peat or muck soils, unless otherwise specified.

CHEMIGATION

PENDANT 3.3 EC may be applied through sprinkler irrigation systems in cotton, field corn, garlic, dry bulb shallots, and direct-seeded and transplanted dry bulb onions, grain sorghum, nonbearing fruit and nut crops, nonbearing vineyards, peanuts, potatoes, soybeans, and sunflowers. Follow all label recommendations for these crops regarding rates per acre, timing of application, special instructions, and precautions.

Apply this product only through a sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of **PENDANT 3.3 EC** applied corresponds to the recommended rate.

Apply **PENDANT 3.3 EC** in ½ to ¾ inches of water during the first sprinkler set. When application is complete, flush the system with water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Special Precautions for Chemigation

1. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
2. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
3. The system must be free of leaks and clogged nozzles.
4. The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor weed control.
5. Agitation must be maintained in the nurse tank.
6. The sprinkler-chemigation system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. All chemigation systems connected to public water systems must also follow restrictions listed in the preceding section titled "CHEMIGATION".

WEEDS CONTROLLED

The following grass and broadleaf weeds are controlled by treatments of **PENDANT 3.3 EC** in all registered crops at the rates recommended for each soil texture in the respective crop section:

Grasses Controlled

Barnyardgrass	Goosegrass
Crabgrass	Johnsongrass, seedling
Crowfootgrass	Panicum, fall
Field Sandbur	Panicum, Texas
Foxtail (giant)	Signalgrass ¹
Foxtail (green)	Witchgrass
Foxtail (yellow)	

Broadleaves Controlled

Bugloss, small	Pigweed (<i>Amaranthus</i> spp.)
Carpetweed	Purslane
Florida pusley	Smartweed, Pennsylvania ²
Kochia	Spurge, annual
Lambsquarters, common	Velvetleaf ²
Lambsquarters, slimleaf	

¹In cotton and soybeans, effective management can be achieved by utilizing a program involving **PENDANT 3.3 EC** preplant incorporated followed by a labeled postemergence grass product.

Signalgrass populations should be managed in all crop rotations for effective control.

²Reduced competition only.

In addition to the weeds listed above as being controlled, PENDANT 3.3 EC (when applied as directed) will control the weeds in the crops indicated in the following table:

WEEDS CONTROLLED-Grasses	CROP
Bluegrass, annual	Nonbearing fruit and nuts and vineyards
Guineagrass	Sugarcane
Itchgrass (Raouigrass)	Soybeans ¹ Sugarcane ² (except in Hawaii)
Johnsongrass, rhizome	Cotton ¹ Soybeans ¹
Junglerice	Nonbearing fruit and nuts and vineyards Rice Sugarcane
Lovegrass	Nonbearing fruit and nuts and vineyards
Panicum, Browntop	Nonbearing fruit and nuts and vineyards Sugarcane
Red Rice	Soybeans ¹
Shattercane	Field Corn ³ Grain Sorghum ³ Soybeans ¹
Sprangletop (<i>Leptochloa</i> spp.)	Rice Nonbearing fruit and nuts and vineyards
Swollen fingergrass	Sugarcane
Volunteer Sorghum	Cotton ⁵ (except in California and Arizona)
Wild Proso Millet	Field ⁴ and Sweet Corn, Grain Sorghum ³ and Peas
Woolly cupgrass	Field ⁴ and Sweet Corn Nonbearing fruit and nuts and vineyards Soybeans ¹
WEEDS CONTROLLED-Broadleaves	CROP
Carpetweed	Sunflowers, no-till Nonbearing fruit and nuts and vineyards
Chickweed, common	Nonbearing fruit and nuts, peas and vineyards
Fiddleneck	Nonbearing fruit and nuts and vineyards
Henbit	Nonbearing fruit and nuts and vineyards
Knotweed, prostrate	Nonbearing fruit and nuts and vineyards
London rocket	Nonbearing fruit and nuts and vineyards
Smartweed, Pennsylvania	Field and Sweet Corn Nonbearing fruit and nuts and vineyards
Puncturevine	Nonbearing fruit and nuts and vineyards
Sheperdspurse	Nonbearing fruit and nuts, peas and vineyards
Stinging nettle	Potatoes
Velvetleaf	Field and Sweet Corn Nonbearing fruit and nuts and vineyards

¹Control or reduced competition (refer to appropriate section for use instructions; see table of contents for page number).

² PENDANT 3.3 EC must be applied at the 7.2 pint/acre broadcast rates for itchgrass control. Surface application (no mechanical incorporation) will provide partial itchgrass control. An additional application of 4.8 to 7.2 pints/acre may be made at layby.

³Control with CULTI-SPRAY applications only. Refer to appropriate section for use instructions; see table of contents for page number.

⁴PENDANT 3.3 EC will control wild proso millet when applied as a CULTI-SPRAY application. PENDANT 3.3 EC will reduce the competition of wild proso millet when applied preemergence. Effective management of wild proso millet can be achieved with a program involving herbicide applications and mechanical cultivation to eliminate weed escapes. PENDANT 3.3 EC plus Accent SP, Bladex or Extrazine II applied early postemergence (no later than when corn is in the 4-leaf stage) treatments have provided the most consistent suppression of wild proso millet because: (1) early postemergence treatments provide a burn-down of the initial germination flush of wild proso millet and (2) early postemergence treatments provide residual activity later into the growing season.

⁵Effective management of woolly cupgrass can be achieved with a program involving herbicide applications and mechanical cultivation to eliminate weed escapes. PENDANT 3.3 EC plus Accent, Bladex or Extrazine II applied early postemergence (no later than when corn is in the 4-leaf stage) treatments have provided the most consistent control of woolly cupgrass because: (1) early postemergence treatments provide a burn-down of the initial germination flush of woolly cupgrass and (2) early postemergence treatments provide residual activity later into the growing season.

⁶Use the high rate for each soil texture. Apply broadcast to flat land and mechanically incorporate (two passes at an angle to one another) prior to bedding up.

SPRAYING INSTRUCTIONS

GROUND APPLICATIONS

Use sprayers with appropriate nozzles that provide uniform spray distribution and minimize drift. Keep the bypass line on or near the bottom of the tank to minimize foaming. Nozzle and in-line screens must be no finer than 50 mesh. DO NOT apply PENDANT 3.3 during periods of gusty winds. As with all herbicides, windy conditions can cause uneven applications.

BROADCAST TREATMENT – Uniformly apply in 10 or more gallons of water or 20 or more gallons of liquid fertilizer as specified in the appropriate sections of this booklet.

BAND TREATMENT – Uniformly apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per acre} = \text{Band VOLUME per acre}$$

AERIAL APPLICATIONS

Uniformly apply in 5 or more gallons of water per acre. Exercise caution to minimize drift. DO NOT apply during periods of gusty winds or when wind conditions favor drifting. Spray drift can cause injury to sensitive crops. It is recommended that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information presented below.

INFORMATION ON DROPLET SIZE:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see WIND, TEMPERATURE AND HUMIDITY, AND TEMPERATURE INVERSIONS).

CONTROLLING DROPLET SIZE

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles – Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

INCORPORATION INSTRUCTIONS

Break up clods before a preplant incorporated application.

1. Flat-planted crop

PENDANT 3.3 EC is to be incorporated prior to planting and within 7 days of application. When **PENDANT 3.3 EC** is applied to flat land that will not be bedded, mechanical incorporation should be into the top 1 or 2 inches of soil. Mechanical incorporation of **PENDANT 3.3 EC** into the top 1 or 2 inches of soil can be achieved by the following:

- (a) Disk harrow set to cut 3 to 4 inches deep and operated in two different directions at 4 to 6 mph. The second pass should be made at an angle to the first.
- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 2 inches deep and operated one time at 4 mph or less.
- (c) Rolling cultivator set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. The second pass should be made at an angle to the first. Use only on coarse and medium textured soils.
- (d) Field cultivator set to cut 3 inches deep operated two times at more than 5 mph. The second pass should be made at an angle to the first. Equipment must have 3 or 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Do not use chisel points.
- (e) Lely-Roterra¹² set and operated according to the manufacturer's directions to thoroughly incorporate **PENDANT 3.3 EC** to a depth of 1 to 2 inches.
- (f) Do-All set and operated according to the manufacturer's directions to thoroughly incorporate **PENDANT 3.3 EC** to a depth of 1 to 2 inches.

NOTE: **PENDANT 3.3 EC** can be incorporated in a single pass when the soil is of good tilth with moderate moisture and relatively free of clods and trash. Implements must be properly set up to thoroughly incorporate **PENDANT 3.3 EC** into the top 1 or 2 inches of soil. Recommended implements include: (1) C-shank or S-shank field cultivators equipped with flexline drag or rolling basket attachment and (2) the combination disk/field cultivator implement equipped with flexline drag or rolling basket attachment. Field cultivators must have 3 or 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Implements must be operated at 6 to 8 miles per hour and set to cut 3 to 4 inches deep.

2. Bedded crop – Application Over Beds

PENDANT 3.3 EC is to be incorporated prior to planting and within 7 days of application. When **PENDANT 3.3 EC** is applied over beds, mechanical incorporation should be into the top 1 or 2 inches of soil. Mechanical incorporation of **PENDANT 3.3 EC** into the top 1 or 2 inches of soil can be achieved by the following:

- (a) Rolling cultivator set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. Use only on coarse and medium textured soils.
- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 2 inches deep and operated one time at 4 mph or less.
- (c) Bed conditioner set to cut 2 to 3 inches deep and operated one time at 4 to 6 mph.

During planting, or if beds must be reshaped prior to planting, avoid tillage that will bring untreated soil to the surface or expose untreated soil in the seedbed or in the furrow.

3. Bedded crop – Application Prior to Bedding

PENDANT 3.3 EC is to be incorporated prior to planting and within 7 days of application. When **PENDANT 3.3 EC** is applied to flat land prior to bedding, mechanical incorporation should be of sufficient depth so that bedding does not bring up untreated soil. Mechanical incorporation of **PENDANT 3.3 EC** can be achieved by the following:

- (a) Disk harrow set to cut 4 to 6 inches deep and operated in two different directions at 4 to 6 mph. The second pass should be made at an angle to the first.

- (b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 3 to 4 inches deep and operated one time at 4 mph or less. During planting, or if beds must be reshaped prior to planting, avoid tillage that will bring untreated soil to the surface or expose untreated soil on the seedbed or in the furrow.

APPLICATIONS WITH LIQUID FERTILIZERS

PENDANT 3.3 EC alone or in tank mix combinations with wettable powders (WP), dry flowables (DF), flowables (F), water soluble concentrates (S) or emulsifiable concentrates (EC) may not combine properly with some liquid fertilizer materials. Small quantities should always be tested before full scale mixing. Follow the testing procedure below to determine compatibility and if a compatibility agent is needed.

Follow all PENDANT 3.3 EC label recommendations regarding registered crops, rates per acre, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. DO NOT apply PENDANT 3.3 EC postemergence in liquid fertilizers.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the PENDANT 3.3 EC/liquid fertilizer mixture.

LIMITED FERTILIZER COMPATIBILITY DETERMINATIONS

If liquid fertilizer/herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of PENDANT 3.3 EC alone or with other herbicides in the specific liquid fertilizer to be used according to the following directions:

- 1. Add 1 pint of fertilizer to a quart jar.
- 2. Add 1 to 4 teaspoon(s) of the DF, WP, AS, F or L formulation (depending on mixing ratio required) to the liquid fertilizer. The number of teaspoons of the formulation to add can be determined by the following formula:

$$\frac{\text{lbs or pts or product/acre}}{\text{Gallons of fertilizer/acre}} \times 11.4 = \text{number of teaspoons of herbicide to add to 1 pint of fertilizer}$$

- 3. Close the jar and agitate until the herbicide(s) are evenly dispersed in the liquid fertilizer. If the materials do not disperse well, it may be necessary to slurry the chemicals in water before adding to the fertilizer.
- 4. After dispersing the materials (step 2) add appropriate number of teaspoons of PENDANT 3.3 EC (see formula in step 2) to the jar and shake well. Add water soluble concentrate herbicides to the mixture last and agitate. Let the mixture stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
- 5. Determine compatibility.
 - (a) If the herbicide(s) and liquid fertilizer mixture does not separate, use this mixture in your spray tank.
 - (b) If the mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (c) If separation of the mixture occurs and agitation does not correct this problem, a compatibility agent* is needed.
- 6. If the need for a compatibility agent* is demonstrated the following procedure is recommended: Using a clean quart jar repeat step one above and add ½ teaspoon of the compatibility agent to the liquid fertilizer. Mix well and repeat steps 2, 3 and 4. If separation or precipitation occurs with the compatibility agent, DO NOT use PENDANT 3.3 EC with that specific liquid fertilizer. An effective compatibility agent will cause the mixture to remain uniformly dispersed with little or no separation (oil rising to the surface) for one half hour or longer. If slight separation occurs, 2 to 3 inversions of the jar should be sufficient to redisperse the mixture uniformly.

Use a clean jar for each test. A compatible mixture will have a uniform appearance and will be relatively easy to redisperse with gentle agitation of the jar.

*Compex-Kalo Laboratories Incorporated, Kansas City, MO; Sponto 168-D-Witco Chemical Company, Houston, TX; Unite-Hopkins Agricultural Chemical Company, Madison, WI; or other comparable adjuvants.

APPLICATIONS WITH DRY BULK FERTILIZERS

PENDANT 3.3 EC may be impregnated on dry bulk fertilizers. When applied as directed, PENDANT 3.3 EC/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of PENDANT 3.3 EC applied in water.

Follow all PENDANT 3.3 EC label recommendations regarding rates per acre, registered crops, incorporation, special instructions and precautions. Apply PENDANT 3.3 EC/dry bulk fertilizer mixtures only with ground equipment.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the PENDANT 3.3 EC/dry bulk fertilizer mixture.

A minimum of 200 pounds of dry bulk fertilizer impregnated with the recommended amount of PENDANT 3.3 EC must be applied per acre.

DO NOT impregnate PENDANT 3.3 EC onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with PENDANT 3.3 EC. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Refer to appropriate crop section in PENDANT 3.3 EC herbicide booklet to determine the rate of PENDANT 3.3 EC per acre. Use the following table to determine the amount of PENDANT 3.3 EC to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer which will be applied per acre.

**Rate Chart for Impregnation of Dry Bulk Fertilizers with PENDANT 3.3 EC
(Pints of Pendant 3.3 EC per Ton of Fertilizer)**

Fertilizer Rate (lbs/acre)	PENDANT 3.3 3C Rate per Acre (recommended rate for crop and soil texture)				
	1.2 pint	1.8 pint	2.4 pint	3.0 pint	3.6 pint
200	12 pts/ton	18 pts/ton	24 pts/ton	30 pts/ton	36 pts/ton
250	9.6 pts/ton	14.4 pts/ton	19.2 pts/ton	24 pts/ton	28.8 pts/ton
300	8 pts/ton	12 pts/ton	16 pts/ton	20 pts/ton	24 pts/ton
350	6.9 pts/ton	10.3 pts/ton	13.7 pts/ton	17.1 pts/ton	20.6 pts/ton
400	6 pts/ton	9 pts/ton	12 pts/ton	15 pts/ton	18 pts/ton
450	5 1/3 pts/ton	8 pts/ton	10 2/3 pts/ton	13 1/3 pts/ton	16 pts/ton

For those rates not listed in this table, calculate the pints of PENDANT 3.3 EC to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times \text{Pints of PENDANT 3.3 EC per acre (recommended rate for crop and soil texture)} = \text{pints of PENDANT 3.3 3C per ton of fertilizer}$$

To impregnate PENDANT 3.3 EC on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of PENDANT 3.3 EC onto the fertilizer during mixing. Apply the PENDANT 3.3 EC/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The PENDANT 3.3 EC/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to Incorporation Instructions section (see table of contents for page number) for incorporation directions.

CULTURAL PRACTICES FOLLOWING APPLICATION

Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control. A PENDANT 3.3 EC treatment may be followed by any registered herbicide to control weeds not listed on the PENDANT 3.3 EC label.

FOLLOW CROP RESTRICTIONS

1. Land treated with PENDANT 3.3 EC may be planted to other crops the following year. See restrictions below for sugar beets, red beets, and spinach.
2. Winter wheat, winter barley
Winter wheat and winter barley may be planted in the fall 4 months after a PENDANT 3.3 EC application in any registered crop. Winter wheat and winter barley may be planted in the fall 3 months after a PENDANT 3.3 EC CULTI-SPRAY application in irrigated field corn or grain sorghum. The treated crop must be grown to maturity and harvested before planting wheat or barley. In areas where irrigation is necessary to produce the crop treated with PENDANT 3.3 EC, DO NOT plant winter wheat or winter barley as follow crops if crop failure/destruction occurs and land is fallowed during the summer as crop injury may result. DO NOT plant winter wheat or winter barley as follow crops in treated land if PENDANT 3.3 EC is applied at 4.8 pints or higher until the next growing season. DO NOT feed forage or graze livestock for 75 days after planting wheat or barley in treated land.
3. Sugar beets, red beets, spinach
To avoid crop injury, do not plant sugar beets, red beets or spinach for 12 months following a PENDANT 3.3 EC application. To insure thorough mixing of soil, land should be plowed using a moldboard plow to a depth of 12 inches prior to planting these crops.
4. When PENDANT 3.3 EC is used in tank-mix or sequential combinations, refer to label of other herbicides for additional follow crop restrictions.
Use of PENDANT 3.3 EC herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and therefore, rotational crop injury is always possible.

FIELD CORN

General Instructions

PENDANT 3.3 EC may be applied in conventional, minimum or no-till as a preemergence, postemergence or CULTI-SPRAY (postemergence incorporated) application in field corn.

DO NOT APPLY PREPLANT INCORPORATED or serious crop injury can result.

DO NOT apply PENDANT 3.3 EC in no-till in California.

With the exception of minimum or no-tillage systems (see below) plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

PENDANT 3.3 EC or PENDANT 3.3 EC tank-mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain corn seeds are below the tilled area.

When PENDANT 3.3 EC alone is used, if corn loss occurs due to weather conditions, corn or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area or crop injury may occur.

DO NOT exceed the maximum labeled rate for any soil type.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Field Corn Grown in Minimum or No-Tillage Systems

PENDANT 3.3 EC alone and PENDANT 3.3 EC tank mixtures may be used in field corn in minimum or no-tillage systems. **PLANT CORN AT LEAST 1 1/2 INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.** The use of no-till planters under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if PENDANT 3.3 EC contacts the germinating corn seed. Check equipment to ensure good seed coverage.

Use Methods and Timings

DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.

Preemergence – Apply PENDANT 3.3 EC after planting, but before weeds and crop emerge.

Postemergence – Apply PENDANT 3.3 EC postemergence until corn is 30 inches tall or has 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

DO NOT apply in liquid fertilizer.

CULTI-SPRAY (postemergence incorporated) – Apply PENDANT 3.3 EC alone or PENDANT 3.3 EC plus atrazine when corn is at least 4 inches tall until layby (last cultivation). Refer to page 12 for complete CULTI-SPRAY instructions.

Tank Mixes and Sequential Programs - PENDANT 3.3 EC may be tank mixed with any registered herbicide(s) labeled for use in field corn. PENDANT can be applied based on the timing indicated by the companion label or until corn is 30" tall or 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

DO NOT apply in liquid fertilizer.

When using tank mixtures or sequential applications with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

PENDANT 3.3 EC Use Rates in Field Corn

Recommended use rates for PENDANT 3.3 EC alone, in tank mix or sequential applications are given in the following table.

Broadcast Rate¹ per Acre of PENDANT 3.3 EC Preemergence or Postemergence in Field Corn (Pints per Acre)

Soil Texture	<1.5% Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	3.6 pts
Medium	2.4 to 3.6 pts	3.6 pts	3.6 to 4.8 pts
Fine	2.4 to 3.6 pts	3.6 to 4.8 pts	3.6 to 4.8 pts

¹Use the high rate for each soil classification when using PENDANT 3.3 EC alone.

CULTI-SPRAY (POSTEMERGENCE INCORPORATED)

General Instructions

PENDANT 3.3 EC or PENDANT 3.3 EC plus atrazine tank mixtures applied as a CULTI-SPRAY (postemergence incorporated) treatment will control most late season annual grasses and certain broadleaf weeds in field corn. PENDANT 3.3 EC treatments can be applied from the 4-inch growth stage to as late as the last cultivation (layby) of field corn. PENDANT 3.3 EC plus atrazine must be applied before the field corn reaches 12 inches in height. PENDANT 3.3 EC treatments will not control established weeds. DESTROY EMERGED WEEDS BY CULTIVATION PRIOR TO PENDANT 3.3 EC APPLICATION.

CULTI-SPRAY (postemergence incorporated) applications of PENDANT 3.3 EC or PENDANT 3.3 EC plus atrazine tank mixture can be applied in field corn previously treated with herbicides registered for use in field corn. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in field corn and for Follow Crop Restrictions.

CULTI-SPRAY (Postemergence Incorporated) APPLICATIONS

PENDANT 3.3 EC alone or PENDANT 3.3 EC plus Atrazine

1. Cultivate with a sweep-type or rolling cultivator operated at sufficient speed to throw at least one inch of soil over the bases of the field corn plants. This will kill small weed seedlings growing in the field corn row and will prevent direct contact of the zone of brace root formation by PENDANT 3.3 EC during application.
2. Apply broadcast with a ground sprayer when corn is at least 4 inches tall up to layby (last cultivation). Use drop nozzles if crop foliage will prevent uniform coverage of the soil surface within the rows.
3. Thoroughly and uniformly incorporate PENDANT 3.3 EC treatments into the soil (1) with a sweep-type or rolling cultivator set to provide thorough incorporation in the top 1 inch of soil OR (2) with adequate irrigation water or rainfall. For best results, incorporate PENDANT 3.3 EC treatments as soon as possible after application. Incorporation must be completed within 7 days after application. If adequate moisture is not received within 7 days after application, incorporate PENDANT 3.3 EC with a sweep-type or rolling cultivator.

Under situations of low rainfall or soil moisture, when deep germinating weeds such as shattercane or field sandbur are anticipated, mechanical incorporation will provide best results.

If cultivation is needed after application and incorporation of PENDANT 3.3 EC, the depth of cut should be no deeper than the depth of cut used to incorporate.

Broadcast Rate Per Acre of PENDANT 3.3 EC CULTI-SPRAY (Postemergence Incorporated) in Field Corn (Pints per Acre)

Soil Texture	Southern States ¹	Northern States ¹
Coarse	1.2 to 1.8 pts	1.8 to 2.4 pts
Medium	1.8 to 2.4 pts	2.4 to 3.6 pts
Fine	1.8 to 3.6 pts	2.4 to 3.6 pts

For PENDANT 3.3 EC plus tank mixtures, do not exceed 1.2 lb. a.i. per acre of atrazine.

¹See map at end of this label for specific states.

NOTE: Livestock can graze or be fed forage from treated field corn after 21 days following application.

SWEET CORN, SEED CORN OR POPCORN

General Instructions

PENDANT 3.3 EC may be used in conventional tillage sweet corn, seed corn or popcorn as a preemergence or postemergence application. **DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.**

DO NOT apply PENDANT 3.3 EC in reduced, minimum or no-till sweet corn, seed corn or popcorn.

Plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 ½ INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

PENDANT 3.3 EC alone will not control emerged weeds. PENDANT 3.3 EC can be applied as a tank mix with atrazine or other herbicides labeled for use in sweet corn, seed corn or popcorn. When using tank mixtures with PENDANT 3.3 EC always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. **Always follow the most restrictive label.**

In addition, follow precautions and restrictions including state and local use restrictions that may apply to specific products.

PENDANT 3.3 EC or PENDANT 3.3 EC tank-mix combination treatments are most effective in controlling weeds when mechanically incorporated or incorporated by adequate rainfall or overhead irrigation within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage.

When PENDANT 3.3 EC alone is used, if corn loss occurs due to weather conditions, any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area or crop injury may occur.

DO NOT exceed the maximum labeled rate for any soil type.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.

Preemergence – Apply PENDANT 3.3 EC after planting, but before weeds and crop emerge.

Postemergence – Apply PENDANT 3.3 EC postemergence until corn is 20-24 inches tall or has 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

DO NOT apply in liquid fertilizer.

Tank Mixes and Sequential Programs - PENDANT 3.3 EC may be tank mixed with any registered herbicide(s) labeled for use in sweet corn, seed corn and pop corn. PENDANT can be applied based on the timing indicated by the companion label or until corn is 20-24" tall or 8 visible leaf collars (V8) which ever is more restrictive. Use drop nozzles and apply as a directed spray if the corn canopy density prevents applications from reaching the soil and target weeds.

DO NOT apply in liquid fertilizer.

When using tank mixtures or sequential applications with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. **Always follow the most restrictive label.**

PENDANT 3.3 EC USE RATES FOR SWEET CORN, SEED CORN OR POPCORN

Recommended use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate¹ per Acre of PENDANT 3.3 EC Preemergence or Postemergence (Pints per Acre)

Soil Texture	<1.5% Organic Matter	1.5-3.0% Organic Matter	>3.0% Organic Matter
Coarse	1.8 to 2.4 pts	2.4 to 3.6 pts	3.6 pts
Medium	2.4 to 3.6 pts	3.6 pts	3.6 to 4.8 pts
Fine	2.4 to 3.6 pts	3.6 to 4.8 pts	3.6 to 4.8 pts

¹Use the high rate for each soil classification when using PENDANT 3.3 EC alone.

RESTRICTIONS FOR POSTEMERGENCE APPLICATIONS

To avoid serious crop injury with postemergence applications observe the following restrictions:

Plant into a seedbed that is firm. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. **PLANT CORN AT LEAST 1 ½ INCHES DEEP. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL.**

- DO NOT APPLY PREPLANT INCORPORATED or serious corn injury can result.
- DO NOT apply following PENDANT 3.3 EC preemergence applications.
- DO NOT apply with liquid fertilizer as the carrier.

COTTON
General Instructions

PENDANT 3.3 EC may be applied in conventional, minimum, stale seedbed or no-till as a preplant surface, preplant incorporated, preemergence or layby application in cotton.

DO NOT use in no-till cotton in California.

Preplant surface, preemergence and layby treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. A shallow cultivation is recommended if soil crusting or soil compaction occurs. If weeds begin to germinate or adequate moisture is not received within 7 days after application, use shallow tillage (rotary hoe or light harrow) and make sure cotton seeds are below tilled area. Otherwise, the use of a postemergence herbicide treatment may be required to control weed escapes at planting or following cotton emergence.

In the event of a crop loss due to weather conditions, cotton or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary. DO NOT rework the soil deeper than the treated zone. There must be an interval of at least 60 days between the last PENDANT 3.3 EC application and cotton harvest.

DO NOT feed forage or graze livestock in treated cotton fields.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

Preplant Surface – Apply PENDANT 3.3 EC up to 15 days prior to planting. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

Preplant Incorporated - Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

Preemergence – Apply PENDANT 3.3 EC at planting or up to 2 days after planting. Apply to a seedbed which is firm and free of clods. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

Preplant Incorporated followed by Preemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply overlay application of PENDANT 3.3 EC at planting or up to 2 days after planting. **Total amount of PENDANT 3.3 EC applied per acre cannot exceed the highest labeled rate for a given soil type.** Preplant incorporated and preemergence applications of PENDANT 3.3 EC may be applied with labeled tank mix herbicide(s).

Layby Application (at last cultivation) - Apply PENDANT 3.3 EC directly to the soil between rows as a directed spray following the last normal cultivation (layby).

Layby applications can be applied in cotton previously treated with PENDANT 3.3 EC or any herbicide(s) registered for use in cotton. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in cotton and for follow crop restrictions. **The total amount of PENDANT 3.3 EC applied per acre per season can not exceed the highest labeled rate for a given soil type.**

DO NOT APPLY as a broadcast spray over the top of the cotton or **SERIOUS CROP INJURY CAN RESULT. AVOID CONTACT OF THE SPRAY** to the non-woody portion of cotton stems and to cotton foliage or **SERIOUS CROP INJURY CAN RESULT.** To reduce the potential for crop injury caused by herbicide contact with cotton foliage and stems, use protective shields when conditions favoring spray drift occur.

Roundup Ultra (or other glyphosate containing products) may be applied at layby with PENDANT 3.3 EC in a tank-mixture for enhanced weed control layby in cotton with the Roundup Ready¹ gene. **DO NOT apply Roundup Ultra (or other glyphosate-containing products) at layby on non-Roundup Ready cotton.**

DO NOT APPLY PENDANT 3.3 EC AND ROUNDUP ULTRA TANK-MIX AS A BROADCAST SPRAY OVER TOP OF COTTON, OR CROP INJURY MAY RESULT.

Fall Application - PENDANT 3.3 EC may be applied for weed control in cotton in the fall, after October 15 (up to 140 days prior to planting cotton) in Arizona, California, Louisiana, New Mexico, Mississippi, Oklahoma and Texas. Apply PENDANT 3.3 EC at the broadcast rate of 1.8 pints on coarse soils, 2.4 pints on medium soils and 3.6 pints on fine soils.

PENDANT 3.3 EC Use Rates in Cotton

Recommended use rates for PENDANT 3.3 EC alone and in tank mix or sequential applications are given in the following table.

Broadcast Rate per Acre of PENDANT 3.3 EC in Cotton (Pints per Acre)

Soil Texture	Conventional or Minimum Tillage	No-Tillage ²
Coarse	1.2 to 2.4 ¹ pts	1.8 to 2.4 pts
Medium	1.8 to 2.4 pts	2.4 to 3.6 pts
Fine	2.4 to 3.6 pts	3.6 to 4.8 pts

¹DO NOT exceed 1.8 pts/A on coarse textured soils in California.

²This use is not recommended for soils with more than 3% organic matter.

For heavy clay soils, apply PENDANT 3.3 EC at the broadcast rate of 3.6 pints per acre.

The high rates for each soil texture above should be used if heavy weed populations are anticipated, extensive crop residues were present prior to seedbed preparation or in no-till.

The total amount of PENDANT 3.3 EC applied per acre per season can not exceed the highest labeled rate for a given soil type.

PENDANT 3.3 EC Tank Mixes and Sequential Programs in Cotton

PENDANT 3.3 EC may be applied in a tank mix or a sequential application with other herbicides registered for use in cotton. Refer to the companion label for weeds controlled in addition to PENDANT 3.3 EC alone.

When using tank mixtures or sequential applications with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

Follow additional use directions in this table for PENDANT 3.3 EC tank mixes.

(Refer to page 14 to determine PENDANT 3.3 EC use rate)

For no-till, PENDANT 3.3 EC alone or PENDANT 3.3 EC tank mixes may be used with Gramoxone Extra or Roundup herbicides to kill existing vegetation if present before planting. Refer to these labels for specific use recommendations, rates and weeds controlled.

<p>PENDANT 3.3 EC plus Fluometuron (i.e. Cotoran) (0.8 to 2.0 lbs ai/A)</p>	<p>Preplant Incorporated followed by preemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply fluometuron as an overlay preemergence application as directed on the fluometuron label (use the fluometuron alone preemergence rates). Preemergence – Apply immediately after planting (or crop injury may result) in water or nitrogen solution with ground equipment only. DO NOT apply tank mix preemergence in Arizona, California, Oklahoma, New Mexico and West Texas.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the fluometuron label. • Plant cotton seeds 1 inch or deeper below the soil surface. • The use of fluometuron as a preemergence application following the use of a systemic insecticide at planting, may result in injury to cotton. DO NOT plant crops other than cotton within 6 months of the last application of fluometuron or injury may result. • Maintain good agitation at all times until spraying is completed. • DO NOT feed foliage from treated cotton plants or gin trash to livestock.
<p>PENDANT 3.3 EC plus prometryn (i.e. Caparol) (1.2 to 2.4 lbs ai/A) For use in Arizona, California, New Mexico, and the Upper and Lower El Paso Valley of Texas.</p>	<p>Preplant Incorporated – Apply with ground equipment prior to listing or over partially finished or finished beds. Incorporate immediately after application.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the prometryn label. • DO NOT use on sands or loamy sands. • DO NOT use this tank mix in cut areas of newly leveled fields, in areas of excess salt or in areas where flooding over the bed is likely to occur as crop injury may result. Cotton should not be planted in tractor wheel depressions or crop injury may result. • DO NOT use this tank mixture when cotton is irrigated up as crop injury may result. On mulch planted cotton, water back only after cotton seedlings are well established. • Maintain good agitation at all times until spraying is completed to keep the material in suspension. Mechanical agitation should be used in Arizona and California. • If replanting is necessary, do not rework the soil.
<p>PENDANT 3.3 EC plus prometryn (i.e. Caparol) For use in specified areas of Oklahoma and Texas.</p>	<p>Preplant Surface – Apply up to 15 days before planting. Preemergence – Apply at planting or up to 2 days after planting before weeds and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the prometryn label. • See the prometryn label for use rates and specified areas of Oklahoma and Texas.
<p>PENDANT 3.3 EC plus Diuron (i.e. Karmex) (0.5 to 1.6 lbs ai/A)</p>	<p>Preplant Incorporated followed by preemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply diuron as an overlay preemergence application as directed on the diuron label (use the diuron alone preemergence rates).</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the diuron label. • The use of diuron as a preemergence application following the use of a systemic insecticide at planting, may result in injury to cotton.
<p>PENDANT 3.3 EC plus Zorial Rapid 80 or 80WP (1.25 to 2.5 lbs/A) DO NOT use in Arizona, California, New Mexico, Oklahoma and Texas.</p>	<p>Preplant Surface – Apply up to 15 days prior to planting. Preplant Incorporated – Apply up to 30 days prior to planting and incorporate within 7 days of application. DO NOT incorporate deeper than 2-3 inches with commonly used equipment. Preemergence – Apply tank mix with ground equipment immediately after planting or crop injury may result. Make sure cotton seeds are placed 1 inch or deeper below soil surface. Preplant Incorporated followed by preemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply Zorial as an overlay preemergence application as directed on the Zorial label (use the Zorial alone preemergence rates).</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Zorial label. • If the cotton plant is stressed during early development, application of PENDANT 3.3 EC plus Zorial at the label rate may result in temporary bleaching or chlorosis of the leaves from which the plant will recover. <p>Maintain good agitation at all times until application is complete.</p>

Preplant Incorporated Application of PENDANT 3.3 EC for Control of Rhizome Johnsongrass

PENDANT 3.3 EC applied preplant incorporated for two consecutive years will provide control of rhizome johnsongrass (*Sorghum halepense*) in cotton at the rates recommended for soil textures listed below. This use is not recommended for Arizona, New Mexico and California. Rhizome johnsongrass will be suppressed after the first year and controlled after the second year.

Before application, use a chisel plow or similar implement to bring johnsongrass rhizomes to the surface. Chop rhizomes into small pieces with a disk harrow set to cut 4 to 6 inches deep and operated in two different directions at 4 to 6 mph.

PENDANT 3.3 EC is to be incorporated into the soil within 7 days after application prior to planting. For maximum control of rhizome johnsongrass, incorporate PENDANT 3.3 EC as soon as possible after application. Deep and thorough incorporation of PENDANT 3.3 EC is necessary for control of rhizome johnsongrass. Mechanical incorporation can be achieved by the following methods:

- a) Disc harrow set to cut 4 to 6 inches deep and operated in two different directions at 4 to 6 mph.
 - b) PTO-driven equipment (tillers, cultivators, hoes) set to cut 3 to 4 inches deep and operated one time at 4 mph or less.
- For johnsongrass escapes during the crop season, cultivation and/or application of registered postemergence herbicides is recommended. Follow the directions for use on the labels of the respective herbicides.

**Broadcast Rate Per Acre of PENDANT 3.3 EC Preplant Incorporated for Control of Rhizome Johnsongrass in Cotton
APPLY FOR TWO CONSECUTIVE YEARS**

Soil Texture (Up to 3% organic matter ¹)	PENDANT 3.3 EC (pints)
Coarse	2.4
Medium	3.6
Fine	4.8

¹This use is not recommended for soils with more than 3% organic matter.

NOTE: DO NOT feed forage or graze livestock in treated cotton fields. If cotton crop loss occurs due to weather conditions, cotton or soybeans can be replanted the same year into treated soil without adverse effects. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.

EDIBLE Beans

Dry, Lima, Snap, Chickpeas (Garbanzo beans), Southern Peas (Cowpeas) and Sweet Lupines

General Instructions

PENDANT 3.3 EC may only be applied preplant incorporated in chickpeas (garbanzo beans), dry beans (such as navy, great northern, red kidney, black turtle, cranberry and small white type), lima beans, snap beans and southern peas (cowpeas).

PENDANT 3.3 EC may be applied preplant incorporated or preemergence in sweet lupines.

DO NOT feed lupine hay and forage or graze livestock in treated lupine fields.

In the event of a crop loss due to weather conditions, beans or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

Preplant Incorporated – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply PENDANT 3.3 EC tank mixes as specified under the tank mix section.

Preemergence – Apply PENDANT 3.3 EC to sweet lupines only at planting or up to 2 days after planting. Apply to a seedbed which is firm and free of clods. DO NOT APPLY PENDANT 3.3 EC PREEMERGENCE (surface treatments) to chickpeas, dry beans, lima beans, snap beans and southern peas (cowpeas) after planting or serious crop injury can result.

PENDANT 3.3 EC Use Rates in Edible Beans

Recommended use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate¹ per Acre of PENDANT 3.3 EC In Edible Beans (Pints per Acre)

Soil Texture	Southern States ¹	Northern States ¹	
		<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 1.8 pts	1.2 to 2.4 pts	2.4 pts
Medium	1.8 to 2.4 pts	1.8 to 3.0 pts	3.0 to 3.6 pts
Fine	1.8 to 3.6 pts	2.4 to 3.6 pts	3.6 pts

¹See map at the end of this label for specific states.

Use the 3.6 pint rate for heavy clay soils.

PENDANT 3.3 EC Tank Mixes in Edible Beans

PENDANT 3.3 EC may be applied in a tank mix with Dual, Eptam, Lasso or Micro-Tech. Refer to the companion label for weeds controlled in addition to PENDANT 3.3 EC alone.

When using tank mixtures with PENDANT 3.3 EC always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

Follow additional use directions in the table for PENDANT 3.3 EC tank mixes.
(Refer to page 16 to determine PENDANT 3.3 EC use rate)

<p>PENDANT 3.3 EC plus Dual (1.5 to 3.0 pts/A)</p> <p>For use in dry, lima or snap beans, chickpeas, and southern peas.</p>	<p>Preplant Incorporated – Apply up to 14 days prior to planting and incorporate within 7 days of application.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Dual label. • DO NOT apply preemergence to dry, lima or snap beans, chickpeas or southern peas. • DO NOT graze or feed forage or fodder. • DO NOT apply to sweet lupines.
<p>PENDANT 3.3 EC plus Eptam (2.5 to 4.5 pts/A)</p> <p>For use in Dry and Snap Beans only.</p>	<p>Preplant Incorporated – Apply with ground sprayer up to 2 days prior to planting. Incorporate immediately (within minutes) after application to prevent loss of Eptam. Whenever possible, application and incorporation should be done in the same application.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Eptam label. • <u>Incorporation Instructions</u> – Use power driven cultivation equipment set to cut to a depth of 2 to 3 inches or tandem disks set to cut to a depth of 4 to 6 inches, operated at 4 to 6 mph, followed by a spike-toothed harrow or some other leveling device which extends beyond the ends of the disks. For thorough mixing, disk in two different directions (cross disk). • DO NOT exceed 3.5 pints Eptam per acre on small white beans on coarse-textured soils. • DO NOT use this tank mixture on Adzuki beans, chickpeas, southern peas (cowpeas, including blackeye peas, blackeye beans), soybeans, lima beans or other flat podded beans. • DO NOT feed bean hay, vines and forage or graze livestock in treated bean fields. • For nutsedge control, use 4.5 pints of Eptam per acre.
<p>PENDANT 3.3 EC plus Lasso or Micro-Tech (2.5 to 3.0 qts/A) or Partner WDG (3.8 to 4.8 lbs/A)</p> <p>For use only in Dry Beans west of the Mississippi River. DO NOT use in California.</p>	<p>Preplant Incorporated – Apply within 7 days prior to planting and incorporate within 7 days of application.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Lasso or Micro-Tech label. • Apply in water with ground equipment only. • This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

FORAGE LEGUMES

General Instructions

PENDANT 3.3 EC may be used in forage legumes used as a cover crop in federal set-aside or conservation reserve program areas. Some stand reduction of the legume cover crop may occur with this use. Consult local county extension service or the local ASC committee for recommended cover crops.

If loss of cover crop occurs due to adverse weather conditions, any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted the same year into PENDANT 3.3 EC-treated soil without adverse effects. If replanting is necessary, DO NOT rework the soil deeper than the PENDANT 3.3 EC-treated zone.

DO NOT feed or graze legume cover crops established following PENDANT 3.3 EC application.

The cover crop residue should ultimately be destroyed by tillage or left on the surface to retard erosion or as directed by the local ASC committee.

Use Methods and Timings

PENDANT 3.3 EC may be applied preplant incorporated or preemergence for weed control in legume cover crops. Refer to **Mixing, Application and Incorporation Instructions, Weeds Controlled and Follow Crop Restrictions** sections (see table of contents for page numbers).

PENDANT 3.3 EC Use Rates in Forage Legumes

Recommended use rates for PENDANT 3.3 EC alone is given in the following table.

Broadcast Rate Per Acre of PENDANT 3.3 EC Preplant Incorporated in Forage Legumes (Pints per Acre)

Soil Texture	Rate
Coarse	1.2 to 1.8 pts
Medium	1.8 to 2.4 pts
Fine	2.4 to 3.0 pts

GARLIC, DRY BULB SHALLOTS, AND DIRECT-SEEDED AND TRANSPLANTED DRY BULB ONIONS

General Instructions

Uniformly apply in 10 or more gallons of water per acre with ground equipment, or 5 or more gallons of water per acre with aircraft.

CHEMIGATION: PENDANT 3.3 EC may only be applied through center pivot, solid set or hand move irrigation. DO NOT apply through other irrigation systems. Apply PENDANT 3.3 EC between the 2 to 9 true leaf stage (2 to 6 true leaf stage in California) only unless otherwise specified in the Additional Use sections below. DO NOT irrigate in excess of 0.5 inches of water. Refer to page 5 for CHEMIGATION directions. PENDANT 3.3 EC treatments are most effective when adequate rainfall or overhead irrigation is received within 7 days after application. DO NOT mechanically incorporate except as specified for use on dry bulb onions in Colorado and the Texas High Plains. DO NOT apply to green (bunching) onions or leeks.

Use Directions for Mineral Soils

Broadcast Rate per Acre of PENDANT 3.3 EC Herbicide for Use on Mineral Soils in All States (Pints per Acre)

Soil Texture	Rate
Coarse	1.2 to 1.8 pts
Medium	1.8 to 2.4 pts
Fine	1.8 to 3.6 pts

DO NOT exceed 3.6 pints per acre per crop (except Idaho, Oregon and Washington). DO NOT apply within 60 days of harvest in California and within 45 days of harvest in all other sales. DO NOT feed or graze these crops. If loss of treated crop occurs due to adverse weather conditions, any crop registered for preplant incorporated use of PENDANT 3.3 EC can be replanted the same year. If replanting is necessary, DO NOT rework the soil deeper than 2 inches.

GARLIC

PENDANT 3.3 EC may be applied to garlic in the following ways:

- (a) Preemergence after planting but before crop and weeds emerge.
- (b) Postemergence to the garlic at the 1 to 5 true leaf growth stage.
- (c) As a split application, apply at both preemergence and post-emergence timings.

DRY BULB SHALLOTS AND DIRECT-SEEDED AND TRANSPLANTED DRY BULB ONIONS

In All States Except California:

Apply PENDANT 3.3 EC as a broadcast treatment when onions or shallots have 2 to 9 true leaves.

Additional Use in Colorado, Kansas and Nebraska:

PENDANT 3.3 EC may be applied sequentially in seeded onions. Apply first application of PENDANT 3.3 EC at loop stage. Apply sequential application of PENDANT 3.3 EC early postemergence (2 to 9 true leaf stage). DO NOT exceed the maximum labeled rate for a given soil texture. DO NOT apply PENDANT 3.3 EC at loop stage through the 9 true leaf stage if heavy rains are expected or severe crop injury may result.

Additional Use in Colorado and the Texas High Plains:

For Transplanted onions only, apply and shallow incorporate (less than 2" deep) PENDANT 3.3 EC into preformed beds prior to transplanting.

Additional use in Idaho, Oregon, and Washington:

Apply PENDANT 3.3 EC as a broadcast treatment when onions or shallots are between the flag leaf to 9 true leaf stage.

PENDANT 3.3 EC may be used at 3.6 to 4.8 pints per acre for dodder control on medium and fine textured solids. DO NOT apply PENDANT 3.3 EC using chemigation at the dodder control rate.

PENDANT 3.3 EC may be applied in the fall or spring to the furrow area of land bedded in the fall in preparation for planting seed of dry bulb onions the following spring. Apply PENDANT 3.3 EC as a banded application at rates based on appropriate soil texture. Band width should be approximately 1/2 the width of the row spacing. Keep PENDANT 3.3 EC away from the area where onion seed will be planted. Harrow-off tops of beds following PENDANT 3.3 EC furrow applications prior to planting onions. For selective weed control in the onion row, apply PENDANT 3.3 EC as a banded postemergence application to flag leaf onions at the labeled rates based on soil texture. Apply PENDANT 3.3 EC only once to the furrow area and once to the onion row as a postemergence application.

Additional Use in Michigan:

For minerals soils containing >10% organic matter, follow the directions for muck soils (see below).

In California:

PENDANT 3.3 EC may only be applied as a single application when onions or shallots have 2 to 6 true leaves.

Use Directions for Muck Soils (Onions Only)

On MUCK SOILS In All States Except California:

PENDANT 3.3 EC may be applied sequentially to ONIONS ONLY on muck soils as follows:

Broadcast Rate per Acre and Application Timing of PENDANT 3.3 EC Herbicide for Muck Soils (Pints per Acre)

Application Timing and Growth Stage	Rate
Preemergence through loop Stage	2.4 to 4.8 pts
Early Postemergence (2 to 6 true leaf stage)	3.6 to 4.8 pts
Late Postemergence (6 to 9 true leaf stage)	3.6 to 4.8 pts

DO NOT apply more that 14.4 pints per acre per growing season on muck soils. To maximize crop safety, ensure good soil coverage during planting or transplanting and delay preemergence applications to the loop stage if possible.

DO NOT apply PENDANT 3.3 EC preemergence through the loop stage if heavy rains are expected or else severe crop injury may result. If irrigating immediately after PENDANT 3.3 EC application at the preemergence through loop stage, DO NOT irrigate in excess of 0.5 inches of water.

DO NOT apply within 45 days of harvest.

DO NOT plant sugar beets, red beets, spinach, winter wheat, or winter barley as rotational crops on muck soils for 12 months from the time of the last applications if more than 3.6 pints per acre of PENDANT 3.3 EC is applied to the onion crop; see the PENDANT 3.3 EC leaflet label for additional follow crop restrictions.

If loss of onion crop occurs due to adverse weather conditions, DO NOT replant any crop other than onions in muck soil during the same cropping year and DO NOT rework the soil deeper that 2 inches.

DO NOT use on muck soils in California.

GRAIN SORGHUM

PENDANT 3.3 EC or PENDANT 3.3 EC plus atrazine may be applied as a CULTI-SPRAY (post emergence incorporated) application in grain sorghum grown in all states.

In addition, PENDANT 3.3 EC plus atrazine may be applied early postemergence in grain sorghum grown in states east of the Mississippi River and in Arkansas, East Texas, Louisiana, and the Missouri 'boothel'.

CULTI-SPRAY (Postemergence Incorporated) For use in all states

General Instructions

PENDANT 3.3 EC or PENDANT 3.3 EC plus atrazine applied as a CULTI-SPRAY (postemergence incorporated) treatment will control most late season annual grasses and certain broadleaf weeds in grain sorghum.

PENDANT 3.3 EC treatments can be applied from the 4-inch growth stage to as late as the last cultivation (layby) of grain sorghum.

PENDANT 3.3 EC plus atrazine must be applied before the grain sorghum reaches 12 inches in height.

PENDANT 3.3 EC treatments will not control established weeds. DESTROY EMERGED WEEDS BY CULTIVATION PRIOR TO PENDANT 3.3 EC APPLICATION.

CULTI-SPRAY (Postemergence incorporated) applications of PENDANT 3.3 EC or PENDANT 3.3 EC plus atrazine tank mixture can be applied in grain sorghum previously treated with herbicides registered for use in grain sorghum. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in grain sorghum and for Follow Crop Restrictions.

DO NOT APPLY PENDANT 3.3 EC in grain sorghum preplant incorporated or preemergence as serious crop injury can result.

DO NOT APPLY PENDANT 3.3 EC in grain sorghum which is planted in double row beds since this cultural practice does not permit adequate soil coverage of the bases of grain sorghum plants with cultivation.

CULTI-SPRAY (Postemergence Incorporated) APPLICATIONS PENDANT 3.3 EC alone or PENDANT 3.3 EC plus Atrazine Tank Mixtures

1. Cultivate with a sweep-type or rolling cultivator operated at sufficient speed to throw at least one inch of soil over the bases of the grain sorghum plants. This will kill small weed seedlings growing in the grain sorghum row and will prevent direct contact of the zone of brace root formation by PENDANT 3.3 EC during application.
2. Apply broadcast with a ground sprayer when grain sorghum is at least 4 inches tall up to layby or last cultivation (atrazine must be applied before grain sorghum reaches 12 inches tall). Use drop nozzles if crop foliage will prevent uniform coverage of the soil surface within the rows.
3. Thoroughly and uniformly incorporate PENDANT 3.3 EC treatments into the soil (1) with a sweep-type or rolling cultivator set to provide thorough incorporation in the top 1 inch of soil OR (2) with adequate irrigation water or rainfall. For best results, incorporate PENDANT 3.3 EC treatments as soon as possible after application. Incorporation must be completed within 7 days after application. If adequate moisture is not received within 7 days after application, incorporate PENDANT 3.3 EC with a sweep-type or rolling cultivator.

Under situations of low rainfall or soil moisture, when deep germinating weeds such as shattercane or field sandbur are anticipated, mechanical incorporation will provide best results.

If cultivation is needed after application and incorporation of PENDANT 3.3 EC, the depth of cut should be no deeper than the depth of cut used to incorporate.

DO NOT APPLY PENDANT 3.3 EC in grain sorghum more than once per crop season.

Broadcast Rate per Acre of PENDANT 3.3 EC CULTI-SPRAY (Postemergence Incorporated) in Grain Sorghum (Pints per Acre)

Soil Texture	Southern States ¹	Northern States ¹
Coarse	1.2 to 1.8 pts	1.8 to 2.4 pts
Medium	1.8 to 2.4 pts	2.4 to 3.6 pts
Fine	1.8 to 3.6 pts	2.4 to 3.6 pts

¹ See map at end of this label for specific states.

For PENDANT 3.3 EC plus atrazine tank mixtures, apply 1.0 lb a.i. per acre of atrazine. DO NOT apply tank mixture on coarse textured soils.

NOTE: Livestock can graze or be fed forage from treated grain sorghum after 21 days following application. Observe all precautions, limitations, and follow crop restrictions on atrazine labels.

EARLY POSTEMERGENCE – For use in states east of the Mississippi River, plus Arkansas, East Texas, Louisiana, and the Missouri 'boothel' only.

Application Instructions

The seedbed should be firm and free of clods and trash. Use only where adequate tillage is practiced to provide good seed coverage.

Plant grain sorghum AT LEAST 1 ½ inches deep to ensure good seed coverage.

Uniformly apply PENDANT 3.3 EC plus atrazine tank mix treatment in water by ground equipment or by aircraft. Apply PENDANT 3.3

EC plus atrazine tank mixture only after grain sorghum has reached the 2-leaf stage and when weeds are no more than 1 inch tall.

DO NOT APPLY PENDANT 3.3 EC in grain sorghum preplant incorporated or preemergence as serious crop injury can result.

DO NOT apply in liquid fertilizer.

PENDANT 3.3 EC plus atrazine treatments are most effective in controlling weeds when adequate rainfall or overhead irrigations is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction, or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain grain sorghum seeds are below the tilled area. Wait 7 to 10 days after application before cultivating.

Broadcast Rate per Acre for Early Postemergence Application of PENDANT 3.3 EC Plus Atrazine in Grain Sorghum

Soil Texture	PENDANT 3.3 EC + Atrazine
Coarse	DO NOT USE
Medium	1.8 to 2.4 pts + 1.0 to 1.2 lbs a.i.
Fine	2.4 pts + 1.0 to 1.2 lbs a.i.

The high rate for each soil texture above should be used if heavy weed populations are anticipated.

Note: DO NOT replant grain sorghum if crop loss occurs due to weather conditions. Observe all precautions, limitations and follow crop restrictions on atrazine labels. Livestock can graze or be fed forage from PENDANT 3.3 EC plus atrazine treated grain sorghum fields after 21 days following application.

NONBEARING FRUIT AND NUT CROPS AND VINEYARDS

General instructions

PENDANT 3.3 EC may be applied for preplant incorporated, preplant surface or preemergence control of most annual grasses and certain broadleaf weeds in several nonbearing fruit and nut crops and vineyards. PENDANT 3.3 EC may be used before or after transplanting the following nonbearing crops.

- Almond
- Apple
- Apricot
- Cherry
- Citrus
- Grape
- Grapefruit
- Lemon
- Nectarine
- Orange
- Peach
- Pear
- Pecan
- Pistachio
- Plum
- Prune
- Tangelo
- Tangerine
- Walnut, English

PENDANT 3.3 EC treatments are most effective in controlling weeds when adequate rainfall or irrigation is received within 21 days after application.

PENDANT 3.3 EC herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Destroy existing weeds before or during application of PENDANT 3.3 EC.

PRECAUTIONS:

Apply the spray directly to the ground beneath the trees or vines. DO NOT apply over the top of trees or vines with leaves or buds. Contact with leaves, shoots or buds by the spray mixture may cause malformed plant tissues.

DO NOT apply to newly transplanted trees or vines until ground has settled and no cracks are present.

DO NOT feed forage or graze livestock in treated fields.

FOR NEWLY TRANSPLANTED AND ONE YEAR OLD GRAPEVINES: Apply only to dormant grapevines. DO NOT apply if buds have started to swell. Application after buds have started to swell may result in leaf distortion.

Use Methods and Timings

Preplant surface – Prior to transplanting, uniformly apply in 10 or more gallons of water per acre (broadcast basis) with ground or aerial equipment. Use PENDANT 3.3 EC use rates listed in the table below.

PENDANT 3.3 EC treatments are most effective in controlling weeds when adequate rainfall or irrigation is received within 21 days after application.

Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Preplant Incorporated – Uniformly apply PENDANT 3.3 EC prior to transplanting but before weeds emerge. Use PENDANT 3.3 EC rates listed in the table below. Incorporate PENDANT 3.3 EC to a depth of 1 to 2 inches. For mechanical incorporation of PENDANT 3.3 EC, use Flex-tyne drag or diamond-toothed harrows operated two times at more than 5 mph with the second pass made at an angle to the first. Application and incorporation must be made prior to transplanting to avoid mechanical injury to the crop.

Avoid root contact with treated soil when placing transplants into the hole or injury may occur.

Chemigation – PENDANT 3.3 EC may be applied through properly equipped chemigation systems.

Preemergence (postplant) – Uniformly apply in 10 or more gallons of water per acre (broadcast basis) with ground equipment. Applications may be band or broadcast.

PENDANT 3.3 EC Use Rates

Recommend use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate Per Acre of PENDANT 3.3 EC Preplant or Preemergence in Nonbearing Fruit and Nut Crops and Vineyards (Quarts per Acre)

Short-term control (4 months)	2.4 qts
Long-term control (6 to 8 months)	4.8 qts

PENDANT 3.3 EC Tank Mixes

PENDANT 3.3 EC may be used in combination with herbicides registered for use in the specific nonbearing crop to remove existing vegetation. Consult contact herbicide label for all directions, precautions and restrictions.

CHEMIGATION

PENDANT 3.3 EC may be applied nonbearing tree and vine crops through solid set, hand move, low volume sprinkler (micro sprinkler) and drip (trickle) irrigation systems. Follow all recommendations, special instructions, and precautions in the general section covering Chemigation at the beginning of this label.

DO NOT apply this product to nonbearing tree and vine crops through any other type of irrigation system.

Low Volume Sprinkler (micro sprinkler) and drip (trickle) Irrigation Instructions.

Low volume sprinkler – 4 to 50 gallons per hour (gph) per emitter, drip – 0.5 to 3 gph per emitter. Point of application should be above ground.

Irrigation system should run a sufficient amount of time prior to PENDANT 3.3 EC injection to have all emitters functioning properly. After system is operation properly, length of injection should be such that at one period of time during the injection, the first and last emitters in the system contain PENDANT 3.3 EC treated water. Add PENDANT 3.3 EC to the supply tank already filled with the volume of water required for the injection period. Maintain proper agitation in PENDANT 3.3 EC injection tank. PENDANT 3.3 EC should be mixed in clean water and injected down-line from filters. Following PENDANT 3.3 EC injection, system should be flushed for a period of time sufficient to clear the line of PENDANT 3.3 EC. (If PENDANT 3.3 EC application is made during a normal irrigation cycle, injection should be made during the last stage.)

Chemigation Calibration (for low volume micro sprinklers)

Calculation of use rate is based on wetted area around emitter – NOT on tree acres. To determine correct amount of PENDANT 3.3 EC, use the following formula:

1. Treated area per each emitter = A

$$A = 3.14 \times (\text{radius} \times \text{radius})$$

Example: If the average distance from emitter to perimeter of wetted area measured one inch below soil surface is 13 inches, then

$$A = 3.14 \times (13'' \times 13'')$$

$$A = 3.14 \times (169'')$$

$$A = 530.7 \text{ square inches}$$

2. The area in square feet wet in each acre = B

$$B = \frac{A \times \text{emitters/acre}}{144}$$

Example: If there are 300 emitters per acre, then

$$B = \frac{530.7 \times 300}{144} = B = 1105.6 \text{ square feet wetted per acre.}$$

3. The total area (in square feet) wet by your system = C

$$C = B \times \text{acres covered by system}$$

Example: If the system covers 20 acres, then

$$C = 1105.6 \text{ square feet per acre} \times 20 \text{ acres}$$

$$C = 22,112 \text{ square feet wetted by system}$$

4. Amount of PENDANT 3.3 EC to inject = S

$$\text{Rate per treated acre of PENDANT 3.3 EC} = R$$

$$S = \frac{C}{43,560} \times R = \text{quarts of PENDANT 3.3 EC}$$

Example: if the desired application rate per treated acre is 2.4 qts of PENDANT 3.3 EC, then

$$S = \frac{22,112}{43,560} \times 2.4 = S = 1.22 \text{ quarts of PENDANT 3.3 EC should be injected into system.}$$

NOTE: Select the proper rate (R) based on length of control required.

PEANUTS

General Instructions

PENDANT 3.3 EC may be applied preplant incorporated in peanuts.

DO NOT use in California

In the event of a crop loss due to weather conditions, peanuts or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, **DO NOT** rework the soil deeper than the treated zone.

Use Methods and Timings

Preplant Incorporated – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply PENDANT 3.3 EC tank mixes as specified under the tank mix section.

PENDANT 3.3 EC Use Rates in Peanuts

Recommended use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate per Acre of PENDANT 3.3 EC in Peanuts (Pints per Acre)

Region	Rate
Texas, Oklahoma and New Mexico	1.2 to 2.4 pts
Other peanut growing states	1.8 to 2.4 pts

For heavy weed infestations, especially of Texas panicum, up to 3.6 pints of PENDANT 3.3 EC can be used in Alabama, Georgia or Florida.

PENDANT 3.3 EC Tank Mixes in Peanuts

PENDANT 3.3 EC may be applied in a tank mix with PURSUIT, Dual or Vernam. Refer to the companion label for weeds controlled in addition to PENDANT 3.3 EC alone.

When using tank mixtures with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

Follow additional use directions in this table for PENDANT 3.3 EC tank mixes.
(Refer to the table above to determine PENDANT 3.3 EC use rate)

PENDANT 3.3 EC plus PURSUIT (4 oz/A) or PURSUIT DG (1.4 oz/A)	Preplant Incorporated – Apply by ground equipment only up to 45 days prior to planting and incorporate within 7 days of application. If crop will be planted on beds, apply and incorporate after bed formation. <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the PURSUIT label. • DO NOT apply tank mix preplant incorporated in West Texas, New Mexico and Arizona.
PENDANT 3.3 EC plus Dual (1.25 to 2.0 pts/A)	Preplant Incorporated – Apply up to 14 days prior to planting and incorporate within 7 days of applications. If crop will be planted on beds, apply and incorporate after bed formation. Read and strictly follow all precautions and instructions on the Dual label.
PENDANT 3.3 EC plus Vernam 7E (2 1/3 to 3.0 pts/A)	Preplant Incorporated – Apply with ground sprayer only up to 10 days prior to planting. Incorporate immediately (within minutes) after application to prevent loss of Vernam herbicide. Whenever possible, application and incorporation should be done in the same operation. Incorporation Instruction – Use power driven cultivation equipment set to cut to a depth of 2 to 3 inches or tandem disks set to cut to a depth of 4 to 6 inches, operated at 4 to 6 mph, followed by a spike-toothed harrow or some other leveling device which extends beyond the ends of the disk. For thorough mixing, disk in two different directions (cross disk). Prior to second disking, raise the disk to prevent cutting deeper than 4 to 6 inches. <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Vernam label. • DO NOT plant seed deeper than 2 inches.

PEAS

(English, Dry, Garden, Dwarf, Green, Pigeon, Edible Pod, and Lentils)

GENERAL INSTRUCTIONS

PENDANT 3.3 EC may be applied preplant incorporated for weed control in peas.

DO NOT use in California.

DO NOT apply PENDANT 3.3 EC preemergence in peas.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

DO NOT USE on peat or muck soils.

NOTE: If loss of pea crop occurs due to adverse weather conditions, peas or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted the same year into PENDANT 3.3 EC treated soils without adverse effects. If replanting is necessary, **DO NOT** rework the soil deeper than the PENDANT 3.3 EC treated zone.

Any crop registered for PENDANT 3.3 EC preplant incorporated use can be double cropped after peas.

DO NOT apply PENDANT 3.3 EC more than once per cropping season.

DO NOT apply to peas, lentils, pea or lentil forage, pea silage, pea hay, or pea straw grown for livestock feed.

When PENDANT 3.3 EC is used in tank-mix or sequential combinations, refer to labels of other herbicides for additional follow crop restrictions. Always follow the most restrictive label.

Use Methods and Timings

Preplant Incorporated – Uniformly apply PENDANT 3.3 EC in 10 or more gallons of water per acre or in 20 or more gallons of liquid fertilizer per acre by ground equipment. Apply in 5 or more gallons of water per acre by aircraft.

Thoroughly mix the previous crop residues into the soil to a depth of 4 to 6 inches by plowing or disking prior to application. PENDANT 3.3 EC may be applied immediately before planting or up to 60 days prior to planting. After application, rotary hoeing, shallow cultivation/tillage, or hand hoeing can be practiced without reducing weed control. Avoid tillage that will bring untreated soil to the surface. A PENDANT 3.3 EC treatment may be followed by any registered herbicide to control weeds not listed on the PENDANT 3.3 EC label. **DO NOT** apply PENDANT 3.3 EC preemergence in peas.

PENDANT 3.3 EC Use Rates in Peas

Recommended use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate per Acre of PENDANT 3.3 EC Preplant Incorporated in Peas

Soil Texture*	PENDANT 3.3 EC
Coarse	1.2 to 1.8 pts
Medium	1.8 to 2.4 pts
Fine	2.4 to 3.6 pts

*Refer to the PENDANT 3.3 EC leaflet label for soil texture classification.

The high rate for each soil texture above, where listed, should be used if heavy weed populations are anticipated or if extensive crop residues are present prior to seedbed preparation. Use the 3.6 pint rate for heavy clay soils.

POTATOES

General Instructions

PENDANT 3.3 EC may be applied preemergence, preemergence incorporated or early postemergence in potatoes. **DO NOT** apply to sweet potatoes or yams. Apply to a seedbed which is firm and free of clods and trash.

DO NOT apply prior to planting crop.

PENDANT 3.3 EC treatments are most effective in controlling weeds when adequate rainfall or irrigation is received within 7 days of application.

DO NOT make more than one application of PENDANT 3.3 EC per season.

In the event of a crop loss due to weather conditions, any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, **DO NOT** rework the soil deeper than the treated zone.

Application of PENDANT 3.3 EC on White Rose variety potatoes during or followed by cool and/or wet weather conditions may result in crop injury.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

Preemergence – Apply PENDANT 3.3 EC after planting but before potatoes and weeds emerge or after drag-off where this operation is practiced. Apply PENDANT 3.3 EC tank mixes as specified under the tank mix section.

Preemergence Incorporated – Apply PENDANT 3.3 EC and incorporate after planting but before potatoes and weed emerge. Where drag-off is practiced, apply PENDANT 3.3 EC and incorporate before, at or after drag-off but before potatoes and weed emerge. Incorporate PENDANT 3.3 EC within 7 dates of application. Apply PENDANT 3.3 EC tank mixes as specified under the tank mix section.

PENDANT 3.3 EC must be thoroughly and uniformly incorporated into the top 1 to 2 inches of soil. Care must be taken so that incorporation equipment does not damage seed pieces or elongating sprouts.

Mechanical incorporation is not required if adequate rainfall for good crop and weed emergence occurs or irrigation is received within 7 days after application

Early Postemergence – Apply PENDANT 3.3 EC from crop emergence to the 6-inch stage of growth. **DO NOT** apply PENDANT 3.3 EC postemergence is potatoes are under stress from cold/wet or hot/dry conditions or crop injury may occur. PENDANT 3.3 EC treatments will not control established weeds. Emerged weeds must be destroyed prior to application. Apply PENDANT 3.3 EC tank mixes as specified under the tank mix section.

Sprinkler Irrigation Systems – Apply PENDANT 3.3 EC alone preemergence after planting, but before potatoes and weeds have emerged, or after drag-off where this operation is practiced, or early postemergence through sprinkler irrigation systems. See **Chemigation** section for complete instructions (see table of contents for page number).

PENDANT 3.3 EC Use Rates in Potatoes

Recommended use rates for PENDANT 3.3 EC alone and in tank mix applications are given in the following table.

Broadcast Rate¹ per Acre of PENDANT 3.3 EC in Potatoes (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 1.8 pts	1.2 to 1.8 pts
Medium	1.8 to 2.4 pts	2.4 to 3.6 pts
Fine	2.4 to 3.6 pts	3.6 pts

¹Use the high rate for PENDANT 3.3 EC alone applications

PENDANT 3.3 EC Tank Mixes in Potatoes

PENDANT 3.3 EC may be applied in a tank mix with Eptam, Lorox, Sencor/Lexone or Eptam plus Sencor/Lexone. Refer to the companion label for weeds controlled in addition to PENDANT 3.3 EC alone. When using tank mixtures with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or

crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

Follow additional use directions in this table for PENDANT 3.3 EC tank mixes.
(Refer to the table above to determine PENDANT 3.3 EC use rate)

<p>PENDANT 3.3 EC Plus Eptam (3.0 to 3.5 pts/A)</p>	<p>Preemergence Incorporated – Apply and incorporate after planting but before potatoes and weeds emerge. In areas where potatoes are normally dragged-off, apply and incorporate following drag-off but before potatoes and weeds emerge. Incorporate immediately after application.</p> <ul style="list-style-type: none"> • <u>Incorporation instructions</u> – Thoroughly and uniformly incorporate into the top 1 or 2 inches of soil. Whenever possible, application and incorporation should be done in the same operation. Care must be taken so that incorporation equipment does not damage seed pieces or elongating sprouts. <p>Early Postemergence – Apply through SPRINKLER IRRIGATION SYSTEMS ONLY from crop emergence to the 6-inch stage of growth. PENDANT 3.3 EC plus Eptam will not control established weeds. EMERGED WEEDS MUST BE DESTROYED PRIOR TO APPLICATION. DO NOT apply if potatoes are under stress from cold/wet or hot/dry conditions or crop injury may result.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Eptam Label. • The Superior variety potato is sensitive to EPTAM and under stress conditions, early season stunting may occur. • For nutsedge control, use 4.5 to 7.0 pints of Eptam per acre.
<p>PENDANT 3.3 EC plus Lorox L (1.5 to 4.0 lbs/A) or Lorox DF (1.5 to 4.0 lbs/A)</p> <p>For use on potatoes grown East of the Rocky Mountains only.</p>	<p>Preemergence – Apply after planting but before potatoes and weeds emerge or after drag-off or hilling where these operations are practiced.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Lorox Label. • Plant seed pieces at least two inches deep. • DO NOT use on sands and loamy sands (except in Wisconsin-Central Sands Area), gravelly soils or exposed subsoils or on soils containing less than 1% organic matter as injury to the treated crop or subsequent crops may result. • Wisconsin-Central Sands Area Only: PENDANT 3.3 EC plus Lorox tank mixture may be used on sands and loamy sands. • Sands – apply 1.2 to 1.8 pints of PENDANT 3.3 EC plus 1 pound of Lorox DF or 1 pint of Lorox L per acre. • Loamy sand – apply 1.2 to 1.8 pints of PENDANT 3.3 EC plus 2 pounds of Lorox DF or 2 pints of Lorox L per acre.
<p>PENDANT 3.3 EC plus Sencor DF/ Lexone DF (0.33 to 0.67 lbs/A) or Sencor F (0.5 to 1.0 pt/A)</p>	<p>Preemergence – Apply after planting but before potatoes and weeds emerge or after drag-off where this operation is practiced.</p> <p>Early Postemergence – Apply from crop emergence to the 6-inch stage of growth. For optimum control of weeds apply before weeds are 1 inch tall. DO NOT apply within 24 hours of application of other pesticides. DO NOT apply within three days after periods of cool, wet or cloudy weather or crop injury may result. DO NOT use on early maturing, smooth-skinned white or red-skinned varieties of potatoes.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Sencor or Lexone label. Observe organic matter restrictions on the labels. • This combination may be applied through sprinkler irrigation systems.
<p>PENDANT 3.3 EC plus Eptam 3.0 to 3.5 pts/A) plus Sencor/ Lexone DF (0.33 to 0.67 lbs/A) or Sencor F (0.5 to 1.0 pt/A) DO NOT use in California.</p>	<p>Early Postemergence – Apply through SPRINKLER IRRIGATION SYSTEMS ONLY from crop emergence to the 6-inch stage of growth. For optimum control of weeds apply before weeds are 1 inch tall.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Eptam and Sencor/Lexone labels. Observe organic matter restrictions on the Sencor/Lexone labels. • DO NOT apply within 24 hours of application of other pesticides. • DO NOT apply within three days after periods of cool, wet or cloudy weather or crop injury may result. • DO NOT use on early maturing, smooth-skinned white or red-skinned varieties of potatoes. • The Superior variety potato is sensitive to EPTAM and under stress conditions, early season stunting may occur. • DO NOT apply within 60 days of potato harvest. • For nutsedge control, use 4.5 to 7.0 pints of Eptam per acre.
<p>PENDANT 3.3 EC plus Eptam (3.0 to 3.5 pts/A) plus Sencor/ Lexone DR (0.33 to 0.67 lbs/A) or Sencor F (0.5 to 1.0 pt/A)</p> <p>For use in Idaho, Oregon and Washington only.</p>	<p>Preemergence – Apply with GROUND EQUIPMENT ONLY after planting but before potatoes and weeds emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Sencor or Lexone label. Observe organic matter restrictions on the labels. • The Superior variety potato is sensitive to EPTAM and under stress conditions, early season stunting may occur. • DO NOT make more than one application per season.

PENDANT 3.3 EC plus Eptam 3.0 to 3.5 pts/A) plus metribuzin (0.33 to 0.67 lbs/A) For use in Idaho, Oregon and Washington only.	Preemergence – SPRINKLER IRRIGATIONS SYSTEMS – after planting but before potatoes and weeds emerge. <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the metribuzin label. Observe organic matter and variety restrictions on the label. • Add metribuzin first and thoroughly mix before adding Eptam. After complete mixing add PENDANT 3.3 EC. Maintain continuous agitation while adding herbicides and until spraying is completed. • The Superior variety potato is sensitive to EPTAM and under stress conditions, early season stunting may occur. • DO NOT make more than one application per season.
PENDANT 3.3 EC plus Matrix (1.0 to 1.5 oz/A)	Preemergence – Apply after planting but before potatoes and weeds emerge or after drag-off where this operation is still practiced. <ul style="list-style-type: none"> • DO NOT apply by air. • DO NOT use on potatoes grown for seed. • Use the most restrictive rotational crop interval. • To avoid injury to desirable crops, follow “Sprayer Tank Cleanout” instructions on the Matrix label. • Read and follow all precautions on the Matrix label.

RICE

General Instructions

PENDANT 3.3 EC plus Facet 75 DF, propanil (or Arrosolo) or propanil plus Londax may be applied as an early postemergence application in dry-seeded rice.

PENDANT 3.3 EC alone or in a tank mix with Facet 75 DF, Bolero 8 EC or glyphosate may be applied as a delayed preemergence application in drilled, dry-seeded rice.

For use in California, refer to Agrilliance’s supplemental labeling entitled “For Weed Control in Drilled, Dry-Seeded Rice in California”.

DO NOT use on water-seeded rice, except as specified in other Agrilliance, LLC labeling.

This pesticide is toxic to fish and aquatic organisms. Fish may be killed at application rates recommended on the label. **DO NOT** contaminate water by cleaning of equipment or disposal of wastes.

DO NOT apply to rice fields if fields are used for fish production, especially catfish or crayfish farming.

DO NOT exceed the maximum labeled rate for any soil type in one season.

DO NOT bale or use rice straw from treated fields for feed or bedding.

DO NOT use water containing **PENDANT 3.3 EC** residues from rice cultivation to irrigate food or feed crops which are not registered for use with **PENDANT 3.3 EC**.

DO NOT plant crops in **PENDANT 3.3 EC** treated fields unless **PENDANT 3.3 EC** is registered for use on those crops.

In case of a crop failure due to weather conditions or disease following treatment with **PENDANT 3.3 EC** alone or in a tank mixture, only drilled dry-seeded rice may be immediately replanted; however, the grower assumes all risks and consequences associated with replanting of rice because there is the potential for stand reduction or stunting. A 10 percent increase in seeding rate is recommended. Replant seed below the herbicide layer because reduced stand or stunting may occur if **PENDANT 3.3 EC** contacts germinating rice seed. **DO NOT** replant with gibberellic acid-treated seed. **DO NOT** reapply **PENDANT 3.3 EC** alone or in a tank mixture.

When using tank mixes with **PENDANT 3.3 EC**, always read the companion product label(s) and follow all precautions and restrictions. Always follow the most restrictive label.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Early Postemergence Applications in Dry Seeded Rice

PENDANT 3.3 EC plus Arrosolo or Propanil

PENDANT 3.3 EC plus Arrosolo or propanil are postemergence treatments that combine the direct contact action of propanil and the residual activity of **PENDANT 3.3 EC**. Since the residual activity of **PENDANT 3.3 EC** provides preemergence control of certain annual grasses which can germinate after this tank-mix treatment is applied, flooding after application can be delayed.

For maximum weed control with this treatment, it is important to carefully follow the directions below for (1) adequate spray coverage of weeds and soil and (2) proper timing of application, when barnyardgrass (watergrass) is in the 1 to 3 leaf stage of growth with an occasional 4 leaf plant (make application when sprangletop is less than ½ inch in height).

The seedbed should be firm and free of clods and trash. The seedbed must be prepared to allow for good seed coverage. Previous crop residues should be thoroughly mixed into the soil to a depth of 4 to 6 inches by plowing or disking before planting rice.

Uniformly apply recommended **PENDANT 3.3 EC** plus Arrosolo or **PENDANT 3.3 EC** plus propanil treatment by aircraft or ground equipment after rice emergence, according to spraying instructions on page 28, when barnyardgrass is in the 1 to 3 leaf growth stage with an occasional 4 leaf plant (make application when sprangletop is less than ½ inch). **THE GROWER SHOULD INSPECT FIELDS FREQUENTLY TO CHECK GROWTH OF BARNYARDGRASS AND/OR SPRANGLETOP TO DETERMINE PROPER APPLICATION TIMING.** Timing of applications should be based on the growth stage of barnyardgrass and/or sprangletop and not on the growth stage of rice. If rice is too small to maintain a flood on the field, the treatment can still be applied since flooding can be delayed because of the residual activity of **PENDANT 3.3 EC**.

While the residual activity of **PENDANT 3.3 EC** allows flooding to be delayed, proper water management practice must be followed for normal rice growth. Flooding should not be delayed if weeds begin to develop after application. Since soil and weeds must be completely exposed to spray coverage, no flood water should be on field at time of application. If necessary, fields may be flushed prior to treatment to produce vigorous rice and weed growth.

Since the residual activity of PENDANT 3.3 EC is activated by moisture, PENDANT 3.3 EC is most effective in controlling emerging weeds when adequate rainfall or irrigation (flush) is received within 7 days after application.

For maximum direct contact activity of propanil or Arrosolo, delay application if there is a chance of rain within 6 to 8 hours

DO NOT apply this tank mixture within 14 days before or after insecticide applications because serious damage to rice may occur.

Broadcast Rate per Acre of PENDANT 3.3 EC plus Arrosolo or Propanil Tank Mixtures in Rice (Pints per Acre)

Soil Texture	Pendant 3.3 EC	Arrosolo or Propanil ¹
Coarse	1.8 pts	6.0 to 8.0 pts
Medium	1.8 to 2.4 pts	6.0 to 8.0 pts
Fine	1.8 to 2.4 pts	6.0 to 8.0 pts

¹Based on 4 lb a.i. per gallon.

Use the high rate of Arrosolo or propanil if the following situations exist at the time of application: (1) unseasonably cool weather, (2) barnyardgrass has progressed to predominantly the 3 to 4 leaf stage, or (3) emerged sprangletop (less than 1/2 inch) is present.

PENDANT 3.3 EC plus propanil plus Londax

Preflood Application

PENDANT 3.3 EC plus propanil (or Arrosolo) may be tank mixed with Londax for early postemergence weed control. Londax will provide improved control of certain broadleaves and sedges when applied as a preflood postemergence application. Refer to the Londax label for specific weeds controlled. Apply Londax at 0.75 to 1.0 oz per acre with PENDANT 3.3 EC plus propanil 1 to 7 days prior to establishment of the permanent flood. Weeds should be actively growing at the time of application.

Preflood Sequential Applications

PENDANT 3.3 EC plus propanil (or Arrosolo) plus Londax may be applied in a sequential application with Londax plus propanil (or Arrosolo). Apply Londax at 0.50 to 0.75 oz per acre with PENDANT 3.3 EC plus propanil when broadleaf weeds are in the cotyledon to 4-leaf stage and the sedges are 3 to 6 inches tall. Apply the second application, if needed, of Londax plus propanil 1 to 7 days prior to establishment of the permanent flood.

PENDANT 3.3 EC plus Facet 75 DF

PENDANT 3.3 EC plus Facet 75 DF herbicides may be tank mixed for early postemergence weed control in dry-seeded rice. This mixture will provide broad spectrum grass and certain broadleaf weed control, especially where sprangletop (*Leptochloa* spp.) is a problem.

Facet 75 DF will control the emerged grasses and broadleaves listed on its label. Refer to the Facet 75 DF label for weed size limitations. PENDANT 3.3 EC will provide residual control of labeled grass weeds which can germinate after this tank mixture is applied and which Facet 75 DF does not control.

For maximum weed control, it is important to carefully follow the directions below for (1) adequate spray coverage of weeds and soil and (2) proper timing of application. PENDANT 3.3 EC will not control emerged weeds. Facet 75 DF does not control sprangletop. Therefore, schedule spraying before sprangletop has emerged.

PENDANT 3.3 EC plus Facet 75 DF tank mixture may be applied early postemergence as follows:

1. Treatments may be applied to conventional, reduced or minimum tillage, and no-till (stale seed-bed) rice. The seedbed should be firm and free of clods. The seedbed must be prepared to allow for complete soil coverage of the rice seed. The use of a planter under conditions which do not allow good soil coverage of the rice seed can result in reduced stand or stunting if PENDANT 3.3 EC contacts germinating rice seed.
2. Apply when soil surface is dry, moist, or wet without standing water. If necessary, fields may be flushed prior to treatment to produce vigorous rice and weed growth. Since soil and weeds must be completely exposed to spray coverage, no flood water should be on the field at time of application. Cloddy soil, standing water (puddles) at the time of application, or cracks in the soil that form after application may result in reduced weed control.
3. Uniformly apply the recommended PENDANT 3.3 EC plus Facet 75 DF treatment after the rice emergence (spiking) and at the correct timing for Facet 75 DF, by aircraft or ground equipment according to spraying instructions on page 28. Refer to timing instructions and illustrations in the Facet 75 DF label. THE GROWER SHOULD INSPECT FIELDS FREQUENTLY TO CHECK GROWTH OF SPRANGLETOP, OTHER LABELED WEEDS AND RICE TO DETERMINE PROPER APPLICATION TIMING.
4. For maximum direct contact activity of Facet 75 DF, delay application of the tank mix if there is a chance of rain within 6 to 8 hours.
5. Because the residual activity of the PENDANT 3.3 EC plus Facet 75 DF allows for delayed flooding, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of PENDANT 3.3 EC and Facet 75 DF. Refer to the Water Management section of the Facet 75 DF label. Flooding should not be delayed if weeds begin to develop after application.

Since the residual activity of PENDANT 3.3 EC and Facet 75 DF is activated by moisture, the tank mix is most effective in controlling emerging weeds when adequate rainfall or irrigation (flush) is received within 3-7 days after application or when new grass/weeds have emerged and are less than one inch tall.

Broadcast Rate per Acre of PENDANT 3.3 EC plus Facet 75 DF Early Postemergence Tank Mixture in Rice

Soil Texture	PENDANT 3.3 EC (pints)	Facet 75 DF (Pounds)
Coarse	1.8	0.50
Medium	1.8 to 2.4	0.50
Fine	1.8 to 2.4	0.50

Use the higher rate of PENDANT 3.3 EC for each soil texture if heavy weed populations are anticipated.

Mixing Instructions

Refer to the Facet 75 DF label for sections on mixing/spraying and on adjuvant for postemergence application. ADDITION OF CROP OIL CONCENTRATE IS REQUIRED WITH FACET 75 DF APPLICATION. DO NOT use liquid fertilizer solution. Add Facet 75 DF to a half-full spray tank of clean water with agitation running. After the Facet 75 DF is thoroughly mixed, add PENDANT 3.3 EC. Mix thoroughly and then add the recommended amount of crop oil concentrate (see Facet 75 DF label) and remaining volume of water. Constantly agitate during application.

Delayed (Late) Preemergence Applications

PENDANT 3.3 EC herbicide alone, or a tank mixture of PENDANT 3.3 EC plus Roundup (glyphosate), Facet 75 DF or Bolero 8 EC herbicides may be applied for delayed (late) preemergence weed control in grain-drilled, dry-seeded rice. The tank mixtures will provide broad spectrum grass and certain broadleaf weed control. Tank mixtures are especially effective where sprangletop (*Leptochloa* spp.) is a problem. PENDANT 3.3 EC alone will not control broadleaf weeds; a postemergence application of propanil, Arrosolo, Bolero, Facet 75 DF or Londax herbicide may be necessary for season-long weed control. Glyphosate or Facet 75 DF will control the emerged grasses and broadleaves listed on their label. PENDANT 3.3 EC will provide residual control of labeled grass weeds which can germinate after the tank mixture is applied and which Facet 75 DF does not control. PENDANT 3.3 EC will not control emerged weeds. Facet 75 DF does not control sprangletop. Therefore, schedule spraying before sprangletop has emerged.

For maximum weed control and rice stand, it is important to carefully follow the directions below for (1) adequate spray coverage of soil and (2) proper timing of application.

1. Treatments may be applied to conventional, reduced or minimum tillage, and no-till (stale seed-bed) rice. The seedbed should be firm and free of clods, and must be prepared to allow for good seed coverage. Plant rice with a grain drill at a depth that provides complete soil coverage of the rice seed. The use of a planter under conditions which do not allow good soil coverage of the rice seed can result in reduced stand or stunting if PENDANT 3.3 EC contacts germinating rice seed.
2. Apply PENDANT 3.3 EC alone or in tank mixture to levees after the levees are pulled and planted. Exposed seeds that come in contact with PENDANT 3.3 EC or Facet 75 DF may be injured.
3. Uniformly apply the recommended PENDANT 3.3 EC plus glyphosate after rice planting (as described below) and before rice emergence (spiking), by aircraft or ground equipment according to spraying instructions on page 28. THE GROWER SHOULD INSPECT FIELDS FREQUENTLY TO CHECK GROWTH OF RICE TO DETERMINE PROPER APPLICATION TIMING. Uniformly apply the recommended rate of PENDANT 3.3 EC alone, PENDANT 3.3 EC plus Facet 75 DF or PENDANT 3.3 EC plus Bolero 8 EC after rice planting (as described below) and before rice and weed emergence (spiking), by aircraft or ground equipment according to spraying instructions on page 28. THE GROWER SHOULD INSPECT FIELDS FREQUENTLY TO CHECK GROWTH OF RICE AND WEEDS TO DETERMINE PROPER APPLICATION TIMING.
4. Apply ONLY when growing conditions favor vigorous rice growth. The seedbed should have adequate moisture for seed germination. If there is insufficient moisture, flushing is recommended before PENDANT 3.3 EC application to supply moisture for root (radicle) initiation and for vigorous rice and weed growth. DO NOT apply PENDANT 3.3 EC and then flush for germination. DO NOT apply to stressed rice. Stress factors include cold or hot temperature extremes, excessive moisture or drought, problem soils, poor field drainage or deep water after application. DO NOT apply early preemergence nor preplant incorporated as severe rice injury is possible.
5. Apply when solid surface is dry, moist, or wet without standing water. Since soil and weeds (only for glyphosate) must be completely exposed to spray coverage, no flood water should be on the field at time of application. Cloddy soil, standing water (puddles) at the time of application, or cracks in the soil that form after application may result in reduced weed control.
6. Apply after the rice seed has absorbed water and germinated and after the soil has been previously sealed over the seed by at least 1 inch of rainfall or by irrigation (flush). If the soil has not been sealed by rain or flush, apply when 80 percent of germinated seeds have a primary root (radicle) or shoot at least 1/2 inch long. If applied to soil prior to these conditions, or to cracked soil, then stand reduction or stunting of rice may occur; Facet 75 DF may cause buggywhipping. Under some conditions, use of gibberellic acid-treated seed. HEAVY RAINFALL AFTER APPLICATION, OR FLUSHING AFTER APPLICATION may result in herbicide injury to rice. Rice can overcome moderate injury with appropriate cultural practices.
7. For maximum direct contact activity of glyphosate, delay application of the tank mix if there is a chance of rain within 6 or 8 hours.
8. Because the residual activity of PENDANT 3.3 EC alone, PENDANT 3.3 EC plus Facet 75 DF or PENDANT 3.3 EC plus Bolero 8 EC tank mix allows for delayed flooding, this treatment may be applied if rice is too small to maintain a flood on the field for weed control. However, proper water management practices must be followed for normal rice growth and activity of PENDANT 3.3 EC, Facet 75 DF or Bolero 8 EC. Refer to the Water Management section of the Facet 75 DF label. Flooding should not be delayed if weeds begin to develop after application. Residual activity of PENDANT 3.3 EC, Facet 75 DF and Bolero 8 EC are most effective when applied to moist soil. Soil should be kept moist after application and not be allowed to crust or crack.

Broadcast Rate per Acre of PENDANT 3.3 EC Alone, PENDANT 3.3 EC plus Facet 75 DF, PENDANT 3.3 EC plus Bolero 8 EC or PENDANT 3.3 EC plus Glyphosate Tank Mixture for Delayed Preemergence Application in Rice.

Soil Texture	PENDANT 3.3 EC (pints)	Facet 75 DF (Pounds)	Bolero 8 EC (pints)
Sands, loamy sands	DO NOT USE	DO NOT USE	DO NOT USE
Sandy loams	1.8	0.33 to 0.44	2 to 3
Loams, silt loams, silts, sandy clay loams	1.8 to 2.4	0.44 to 0.5	3 to 4
Silty clay loams, clay loams, sandy clays, silty clays, clays	1.8 to 2.4	0.5 to 0.67	3 to 4

Use the higher rate of **PENDANT 3.3 EC** for each soil texture if heavy weed populations are anticipated. See the glyphosate label for recommended rates.

Mixing Instructions

PENDANT 3.3 EC plus glyphosate tank mix. Refer to Mixing Instructions in the **PENDANT 3.3 EC** leaflet label and to Mixing, Additives and Application Instructions in the glyphosate label. ADDITION OF NONIONIC SURFACTANT MAY BE REQUIRED WITH GLYPHOSATE APPLICATION (see product label) and use of ammonium sulfate may increase performance.

PENDANT 3.3 EC plus Facet 75 DF tank mix. Refer to the **PENDANT 3.3 EC** and **Facet 75 DF** labels for sections on mixing and spraying. **DO NOT** use liquid fertilizer solution. Add **Facet 75 DF** to a half-full spray tank of clean water with agitation running. After the **Facet 75 DF** is thoroughly mixed, add **PENDANT 3.3 EC**. Mix thoroughly and then add the remaining volume of water. Constantly agitate during application.

PENDANT 3.3 EC plus Bolero 8 EC tank mix. Add **PENDANT 3.3 EC** to the half-full spray tank of clean water with agitator running. After **PENDANT 3.3 EC** is thoroughly mixed, add **Bolero 8 EC** and mix thoroughly. Then add the remaining volume of water. Constantly agitate during application.

Spraying Instructions

Spray drift can cause injury to sensitive crops. See the propanil or Arrosolo, **Facet 75 DF**, **Bolero 8 EC** and glyphosate labels for sensitive crops: follow all recommendations to minimize drift.

DO NOT apply **PENDANT 3.3 EC** through any type of irrigation system.

DO NOT apply in liquid fertilizer.

Aerial Applications

For aerial application, apply the recommended rate in 5 to 10 (10 to 12 for propanil and Arrosolo) gallons of water per acre to ensure adequate coverage. To minimize drift, **DO NOT** apply during periods of gusty winds or when wind conditions favor drifting. It is recommended that a flagman or an automatic mechanical flagging unit on the aircraft be used to avoid overlapping and possible crop injury.

Ground Applications

For ground equipment, apply the recommended rate in 10 to 20 (15 to 25 for propanil and Arrosolo) gallons of water per acre to ensure adequate coverage. Use a properly calibrated low-pressure (20 to 40 psi) sprayer equipped with 8002 or larger size Tee-jet or comparable nozzles to achieve uniform spray distribution and minimize drift. Keep the bypass line on or near the bottom of the tank to minimize foaming. Nozzle screens must be no finer than 50 mesh. **DO NOT** apply **PENDANT 3.3 EC** during periods of gusty winds or when wind velocity is greater than 20 mph.

SOYBEANS

General Instructions

PENDANT 3.3 EC may be applied in conventional, minimum or no-till as a fall surface, fall incorporated, preplant surface, preplant incorporated or preemergence application in soybeans, including Round-up Ready⁵ Soybeans.

DO NOT APPLY POSTEMERGENCE or serious crop injury can result.

DO NOT make applications of **PENDANT 3.3 EC** preemergence north of Interstate 80, except as specified in other supplemental Agriliance, LLC labeling.

DO NOT use **PENDANT 3.3 EC** in soybeans in California.

Preplant surface and preemergence treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If moisture is insufficient to activate **PENDANT 3.3 EC**, a shallow cultivation (preferably with a rotary hoe) should be made after emergence of soybeans, but while weeds are small enough to be controlled by mechanical cultivation. Otherwise, the use of a postemergence herbicide treatment may be required to control weed escapes at planting or following soybean emergence.

In the event of a crop loss due to weather conditions, soybeans or any crop registered for preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, **DO NOT** rework the soil deeper than the treated zone.

Livestock can graze or be fed forage from treated soybean fields. **PENDANT 3.3 EC** is not recommended for use on peat or muck soils.

Use Methods and Timing

Fall Applied - PENDANT 3.3 EC may be surface applied or incorporated in the fall, October 1 to December 31, in states north of I-80 and the entire states of Iowa, Oklahoma and Texas. Fall applications of PENDANT 3.3 EC will not provide season long weed control. For season long weed control, fall applications of PENDANT 3.3 EC should be followed by a postemergence program using PURSUIT, RAPTOR, or other registered postemergence herbicide.

Preplant Surface - Apply PENDANT 3.3 EC up to 15 days prior to planting. PENDANT 3.3 EC may be applied up to 45 days prior to planting when used in tank mix or applied sequentially with PURSUIT or SCEPTER herbicides. Preplant surface applications of PENDANT 3.3 EC alone should be followed by a postemergence program using RAPTOR or PURSUIT herbicide. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

Preplant Incorporated - Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

Preemergence - Apply PENDANT 3.3 EC at planting or up to 2 days after planting. Apply to a seedbed which is firm and free of clods. **DO NOT** make applications of PENDANT 3.3 EC Preemergence north of Interstate 80, except as specified in other supplemental Agrilience, LLC labeling. Apply PENDANT 3.3 EC tank mixes and sequential programs as specified under the tank mix section.

DO NOT apply PENDANT 3.3 EC postemergence or serious crop injury may result.

PENDANT 3.3 EC Use Rates in Soybeans

Recommended use rates for PENDANT 3.3 EC alone and in tank mix combinations or sequential applications with other herbicides are given in the following tables.

Broadcast Rate Per Acre of PENDANT 3.3 EC Fall Surface, Fall Incorporated, Preplant Surface Or Preplant Incorporated in Soybeans (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 1.8 pts	2.4 pts
Medium	1.8 to 3.0 ¹ pts	3.0 to 3.6 pts
Fine	2.4 to 3.6 pts	3.6 pts

¹Do not exceed 2.4 pts/A for Southern states. See map at the end of this label for specific states. For heavy clay soils, apply PENDANT 3.3 EC at the broadcast rate of 3.6 pints per acre. The high rates for each soil texture above should be used if heavy weed populations are anticipated, extensive crop residues were present prior to seedbed preparation or in no-till.

Broadcast Rate Per Acre of PENDANT 3.3 EC Preemergence in Soybeans (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 1.8 pts	1.8 pts
Medium	1.8 to 2.4 pts	1.8 to 2.4 pts
Fine	1.8 to 2.4 pts	2.4 to 3.0 pts

The high rates for each soil texture above should be used if heavy weed populations are anticipated, expensive crop residues were present prior to seedbed preparation or in no-till

PENDANT 3.3 EC Tank Mixes and Sequential programs in Soybeans

PENDANT 3.3 EC may be applied in a tank mix with PURSUIT, SCEPTER, Command, Canopy, Dual, Lasso, Lorox, Lorox Plus, Preview, and Sencor/Lexone. PENDANT 3.3 EC may be applied in a sequential application with PURSUIT, RAPTOR, SCEPTER, SCEPTER O.T., Lorox and Sencor/Lexone. Refer to the companion label for weeds controlled in addition to PENDANT 3.3 EC alone.

When using tank mixtures or sequential applications with PENDANT 3.3 EC, always read the companion product label(s) to determine the specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow all precautions and restrictions including state and local use restrictions that may apply to specific products. Always follow the most restrictive label.

For no-till, PENDANT 3.3 EC alone or PENDANT 3.3 EC tank mixes may be used with Gramoxone Extra, Roundup, 2-4, D or Butyrac 200 (2,4-DB) to kill existing vegetation, if present prior to planting. Refer to these labels for specific use recommendations, restrictions, rates and weeds controlled. PENDANT 3.3 EC alone will control weeds as they germinate, but it will not control emerged weeds.

Follow additional use directions in this table for PENDANT 3.3 EC tank mixes. (Refer to the table above to determine PENDANT 3.3 EC use rate)

<p>PENDANT 3.3 EC plus PURSUIT (4 oz/A) or PURSUIT DG (1.4 oz/A)</p>	<p>Preplant Surface – Apply up to 45 days prior to planting. Preplant Incorporated – Apply up to 45 days prior to planting and incorporate within 7 days of application. Preemergence – Apply at planting or up to 2 days after planting before weeds and crops emerge. Preplant Surface, preplant incorporated or preemergence followed by Early Postemergence – Apply PENDANT 3.3 EC as described above. Follow with a early postemergence treatment of PURSUIT as directed on the PURSUIT label.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the PURSUIT label. • Preplant surface applications of PENDANT 3.3 EC alone should be followed by a postemergence program using PURSUIT herbicide. • DO NOT graze or feed treated soybean forage, hay or straw to livestock.
<p>PENDANT 3.3 EC followed by RAPTOR (4 oz/A)</p>	<p>Preplant Surface followed by Early Postemergence – Apply PENDANT 3.3 EC up to 15 days prior to planting. Follow with a early postemergence treatment of RAPTOR as directed on the RAPTOR label. Preplant Incorporated followed by Early Postemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Follow with an early postemergence treatment of RAPTOR as directed on the RAPTOR label. Preemergence followed by Early Postemergence – Apply PENDANT 3.3 EC at planting or up to 2 days after planting. Follow with a postemergence treatment of RAPTOR as directed on the RAPTOR label.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the RAPTOR label.
<p>PENDANT 3.3 EC plus SCEPTER (0.67 pt/A) or SCEPTER 70 DG 2.8 oz/A)</p>	<p>Preplant Surface – Apply up to 45 days prior to planting. Preplant Incorporated – Apply up to 45 days prior to planting (30 days in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee and Texas) and incorporate within 7 days of application. Preemergence - Apply at planting or up to 2 days after planting before weeds and crops emerge. Preplant Surface, Preplant Incorporated or preemergence followed by Early Postemergence – Apply PENDANT 3.3 EC as described above. Follow with a early postemergence treatment of SCEPTER as directed on the SCEPTER label.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the SCEPTER label. • DO NOT graze or feed treated soybean forage, hay or straw to livestock.
<p>PENDANT 3.3 EC followed by SCEPTER O.T. (1.0 pt/A)</p>	<p>Preplant Surface Followed by Early Postemergence – Apply PENDANT 3.3 EC up to 15 days prior to planting. Follow with a early postemergence treatment of SCEPTER O.T. as directed on the SCEPTER O.T. label. Preplant Incorporated followed by Early Postemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Follow with an early postemergence treatment of SCEPTER O.T. as directed on the SCEPTER O.T. label. Preemergence followed by Early Postemergence – Apply PENDANT 3.3 EC at planting or up to 2 days after planting. Follow with a postemergence treatment of SCEPTER O.T. as directed on the SCEPTER O.T. label.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the SCEPTER O.T. label.
<p>PENDANT 3.3 EC plus Command 4EC (0.75 to 1.5 pts/A)</p>	<p>Preplant Incorporated – Apply to the soil surface and uniformly incorporate. See the Command label for incorporation requirement and instructions.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Command label. • Apply with calibrated ground equipment in 10 to 40 gallons of water per acre. The use of an agriculturally approved drift reduction agent is required at finished spray volumes of 10 to 15 gallons per acre. • Do not apply this tank mix to overly moist or wet soils. • Command is a volatile compound. Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plant parts. • DO NOT allow livestock to graze on soybean vines or use the vines for feed. Cover crops may be planted anytime but stand reductions may occur. DO NOT graze or use the cover crop for feed.
<p>PENDANT 3.3 EC plus Canopy (6 to 12 oz/A)</p>	<p>Preplant Surface – Apply up to 30 days before planting. Preplant Incorporated – Apply up to 14 days prior to planting and incorporate within 7 days of application. Preemergence – Apply at planting or up to 2 days after planting before weeds and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Canopy label. Observe all soil type, soil pH and soybean variety restrictions. • DO NOT apply to soils with less than 0.5% organic matter. • Continuous agitation in the spray tank is required to keep the material in suspension. <p>Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, or crop injury may result.</p>
<p>PENDANT 3.3 EC plus Dual (1.5 to 3.0 pts/A)</p>	<p>Preplant Surface - Apply up to 15 days before planting. Preplant Incorporated – Apply up to 14 days prior to planting and incorporate within 7 days of application. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If soybeans are planted on beds, apply and incorporate after bed formation. Preemergence – Apply at planting or up to 2 days after planting before weeds and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Dual label.

<p>PENDANT 3.3 EC plus Lasso (2.5 to 4.0 qts/A)</p>	<p>Preplant Surface – Apply up to 15 days before planting. Preplant Incorporated – Apply up to 7 days prior to planting and incorporate within 7 days of application. Preemergence – Apply at planting or up to 2 days after planting before weeds and crops emerge. This treatment must be applied within 5 days of the last preplant tillage.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Lasso label. • Apply with ground equipment. • Incorporation on coarse soils may reduce length of control due to leaching of lasso with rainfall or irrigation. • Incorporation is not recommended on coarse soils in the Southeastern States. • High intensity or excessive rainfall or excessive irrigation after preemergence application may reduce weed control.
<p>PENDANT 3.3 EC plus Lorox DF (1.0 to 2.5 lbs/A) or Lorox L (1.0 to 2.5 pts/A)</p>	<p>Preplant Surface – Apply up to 15 days before planting. Preplant Incorporated followed by Preemergence– Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Follow with a preemergence treatment of Lorox as directed on the Lorox label. Preemergence – Apply at planting or up to 2 days after planting before weed and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Lorox label. • DO not use on sands, loamy sands, gravelly soils or soils containing less than 1/2% organic matter.
<p>PENDANT 3.3 EC plus Lorox Plus (12 to 18 oz/A)</p>	<p>Preplant Surface – Apply up to 30 days before planting. Preplant Incorporated – Apply up to 30 days prior to planting and incorporate within 7 days of application. Preemergence – Apply at planting or up to 2 days after planting before weed and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Lorox Plus label. Observe all soil type, soil pH and soybean variety restrictions. • DO NOT apply to soils with less than 0.5% organic matter. • Continuous agitation in the spray tank is required to keep the material in suspension. • Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, or crop injury may result.
<p>PENDANT 3.3 EC plus Preview (6 to 10 oz/A)</p>	<p>Preplant Surface – Apply up to 30 days before planting. Preplant Incorporated – Apply up to 14 days prior to planting and incorporate within 7 days of application. Preemergence – Apply at planting or up to 2 days after planting before weed and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Preview Label. Observe all soil type, soil pH and soybean variety restrictions. • DO NOT apply to soils with less than 0.5% organic matter • Continuous agitation in the spray tank is required to keep the material in suspension. • Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, or crop injury may result.
<p>PENDANT 3.3 EC plus Sencor/ Lexone DF (0.33 to 0.67 lbs/A) or Sencor F (0.5 to 1.0 pt/A)</p>	<p>Preplant Surface – Apply up to 15 days before planting. Preplant Incorporated – Apply up to 7 days prior to planting and incorporate within 7 days of application. Preplant Incorporated followed by Preemergence – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application. Follow with a preemergence treatment of Sencor/Lexone as directed on the Sencor or Lexone label. Preemergence – Apply at planting or up to 2 days after planting before weed and crops emerge.</p> <ul style="list-style-type: none"> • Read and strictly follow all precautions and instructions on the Sencor/Lexone label. Observe all soil type, soil pH and soybean variety and incorporation restrictions. • DO NOT use on sands. DO NOT use on loamy sands or on sandy loams that contain less than 0.5% organic matter as crop injury may result. This tank mixture is not recommended for use on soils with less than 2 % organic matter in the Coastal Plain of New Jersey or the Delmarva Peninsula. • Soybean seed should be planted at least 1.5 inches below the soil surface. • If replanting is necessary, DO NOT rework the soil.

Special Weeds

1. Shattercane/Woolly Cupgrass

Preplant incorporated treatments of **PENDANT 3.3 EC** alone, **PENDANT 3.3 EC** plus **SCEPTER** or **PENDANT 3.3 EC** plus **Sencor/Lexone** preplant incorporated tank mixtures, or preplant incorporated **PENDANT 3.3 EC** applications followed by registered sequential preemergence or postemergence herbicides will control shattercane (wild cane) (*Sorghum bicolor*).

Preplant incorporated treatments of **PENDANT 3.3 EC** plus **PURSUIT**® or preplant incorporated **PENDANT** applications followed by an early postemergence application of **PURSUIT** herbicide or other registered sequential preemergence or postemergence herbicides will control woolly Cupgrass (*Eriochola villosa*).

Shattercane and woolly Cupgrass are difficult to control in many soybean growing areas because of their ability to germinate from a greater depth in the soil than most other weeds. Shattercane and woolly Cupgrass can emerge throughout the growing season depending on soil temperature, moisture, and seed dormancy.

Thoroughly mix previous crop residues into the soil and destroy any existing vegetation prior to herbicide application. Apply **PENDANT 3.3 EC** or **PENDANT 3.3 EC** combinations at the recommended rates listed in table following. See **Incorporation Instructions** section (see table of contents for page number) for incorporation directions. Thoroughly and uniformly mechanically

incorporate PENDANT 3.3 EC into TOP 2 INCHES of soil. Acceptable results CANNOT be expected if PENDANT 3.3 EC is applied preemergence. Cultivation may be necessary to control escapes and late germinating shattercane or wooly cupgrass during the crop season.

Broadcast Rate Per Acre of PENDANT 3.3 EC Preplant Incorporated for Control of Shattercane or Wooly Cupgrass in Soybeans (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	2.4 pts	2.4 pts
Medium	3.0 pts	3.6 pts
Fine	3.6 pts	3.6 pts

Broadcast Rate Per Acre of PENDANT 3.3 EC Plus Sencor/Lexone Preplant Incorporated Tank Mixture for Control of Shattercane in Soybeans¹

Soil Texture ¹	Pendant 3.3 EC + Sencor/Lexone DF 0.5% to 3% Organic Matter	PENDANT 3.3 EC + Sencor/Lexone ² DF More than 3% Organic Matter
Coarse	1.8 pts + 0.5 lb.	1.8 pts to 0.5 lb.
Medium	2.4 pts + 0.5 lb.	2.4 pts + 0.5 to 0.67 lb.
Fine	2.4 pts + 0.5 to 0.67 lb.	3.0 pts + 0.67 lb.

¹ DO NOT USE on sands. DO NOT USE on loamy sands or sandy loams that contain less than 1% organic matter, or on other soils with less than 1/2% organic matter as crop injury may result.

² Read and strictly follow all precautions and instruction on the Sencor/Lexone Label. Observe all soil type, soil pH and soybean variety restrictions.

2. Red Rice, Itchgrass

PENDANT 3.3 EC applied preplant incorporated will provide control of red rice (*Oryza sativa*) and aid in the control of and reduce competition from itchgrass (*Rottboellia exaltata*) at the rates listed in the following table.

Broadcast Rate Per Acre of PENDANT 3.3 EC Preplant Incorporated for Control of Red Rice and Suppression of Itchgrass in Soybeans (Pints per Acre)

Soil Texture	Up to 3% Organic Matter ¹
Coarse	2.4 to 3.6 ² pts
Medium	3.6 pts
Fine	4.8 pts

¹ This use is not recommended for soils with more than 3% organic matter.

² The higher rate should be used if heavy red rice or itchgrass populations are anticipated.

Note: Livestock can graze or be fed forage from treated soybean fields. If soybean crop loss occurs due to weather conditions, cotton or soybeans can be replanted the same year into treated soil without adverse effects. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.

3. Rhizome Johnsongrass

PENDANT 3.3 EC applied preplant incorporated for two consecutive years will provide control of rhizome Johnsongrass (*Sorghum halepense*) in soybeans at the rates recommended for soil textures listed in the following table. This use is not recommended for Arizona, New Mexico and California. Rhizome Johnsongrass will be suppressed after the first year and controlled after the second year.

Before application, use a chisel plow or similar implement to bring johnsongrass rhizomes to the surface. Chop rhizomes into small pieces with a disk harrow set to cut 4 to 6 inches deep and operated in two different directions at 4 to 6 mph

PENDANT 3.3 EC is to be incorporated into the soil within 7 days after application prior to planting. For maximum control of rhizome johnsongrass, incorporate PENDANT 3.3 EC as soon as possible after application. Deep and thorough incorporation of PENDANT 3.3 EC is necessary for control of rhizome johnsongrass. Mechanical incorporation can be achieved by the following methods:

(a) Disk harrow set to cut 4 to 6 inches deep and operated in two different directions at 4 to 6 mph.

(b) PTO-driven equipment (tiller, cultivators, hoes) set to cut 3 to 4 inches deep and operated one time at 4 mph or less.

For johnsongrass escapes during the crop season, cultivation and/or application of registered postemergence herbicide are recommended. Follow the directions for use on the labels of the respective herbicides.

**Broadcast Rate per Acre of PENDANT 3.3 EC Preplant Incorporated for Control of Rhizome Johnsongrass
APPLY FOR TWO CONSECUTIVE YEARS (Pints per Acre)**

Soil Texture	Up to 3% Organic Matter ¹
Coarse	2.4 pts
Medium	3.6 pts
Fine	4.8 pts

¹This use is not recommended for soils with more than 3% organic matter.

SUGARCANE (except Hawaii)

General Instructions

PENDANT 3.3 EC may be applied to newly planted or ratoon sugarcane preemergence through layby and again in the late summer or early fall to the newly planted sugarcane.

DO NOT apply through irrigation systems on sugarcane.

DO NOT make aerial applications at close-in because complete and uniform coverage cannot be obtained.

DO NOT apply more than 14.4 pints per acre of PENDANT 3.3 EC during one growing season.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

DO NOT apply within 90 days of harvest.

DO NOT graze treated fields or feed treated forage or fodder to livestock.

Use Methods and Timings

PENDANT 3.3 EC may be applied preemergence through layby, to newly planted or ratoon sugarcane, and again in late summer or early fall to the newly planted sugarcane. Applications may be made band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

PENDANT 3.3 EC Use Rates

Broadcast Rate of PENDANT 3.3 EC for use in Sugarcane (except Hawaii)

Apply 4.8 to 7.2 pints of PENDANT 3.3 EC but DO NOT exceed 14.4 pints per acre in one growing season. See spraying instructions for calculation of band treatment rate.

Use the 7.2 pint rate:

- a. heavy clay soils;
- b. if no mechanical incorporation is planned;
- c. if heavy weed populations are anticipated;
- d. if itchgrass infestation is anticipated;
- e. or if no shaving is planned.

INCORPORATION INSTRUCTIONS

PENDANT 3.3 EC must be thoroughly and uniformly incorporated into the soil with either (a) mechanical incorporation equipment as outlined below, or (b) with rainfall or irrigation, if rainfall or irrigation is adequate for good crop and weed emergence and received within 7 days after application. If rainfall or irrigation is not obtained, PENDANT 3.3 EC should be mechanically incorporated.

Mechanical Incorporation

PENDANT 3.3 EC herbicide should be applied to loosened beds and incorporated into the top 1 to 2 inches of soil within 7 days after application. Ratoon sugarcane must be lightly shaved in early spring to remove the old stubble before incorporation over the line of sugarcane is possible. Carefully adjust equipment to incorporate without causing excessive damage to emerging shoots. Mechanical incorporation can be achieved by the following:

- a. Rolling cultivator (Lilliston type – Lely Roterra) set to cut 2 or 3 inches deep and operated two times at 6 to 8 mph. This technique may be used with all application timings.
- b. Rolling disc cultivator (Hipper) set to cut 2 to 3 inches deep and operated two times at 6 to 8 mph. This technique may be used to incorporate between sugarcane lines (rows) at layby only.

PENDANT 3.3 EC Tank Mixes

PENDANT 3.3 EC may be used in combination with any registered herbicide. Consult labels for all directions, precautions and restrictions. Always check compatibility when applying in a tank mix and follow the most restrictive label.

SUGARCANE for use in Hawaii

General Instructions

PENDANT 3.3 EC may be applied preemergence through layby in plant or ratoon sugarcane.

DO NOT apply through irrigation systems on sugarcane.

DO NOT make aerial applications at close-in because complete and uniform coverage cannot be obtained.

DO NOT apply more than 14.4 pints per acre of PENDANT 3.3 EC during one growing season.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

DO NOT apply within 90 days of harvest.

DO NOT graze treated fields or feed treated forage or fodder to livestock.

Use Methods and Timings

PENDANT 3.3 EC may be applied twice per season, preemergence through layby, in plant or ratoon sugarcane. Applications may be made band or broadcast. Although there may be adequate crop tolerance for postemergence applications at layby, the spray must be directed under the sugarcane canopy in order to obtain effective weed control.

PENDANT 3.3 EC Use Rates

Broadcast Rate of PENDANT 3.3 EC for Sugarcane Grown in Hawaii

Apply 4.8 to 9.7 pints per acre of PENDANT 3.3 EC but DO NOT exceed 14.4 pints per acre in one growing season. See spraying instructions for calculation of band treatment rate.

Use the higher rates:

- a. for dark clay soils;
- b. if no mechanical incorporation is planned;
- c. if heavy weed populations are anticipated;
- d. or if no shaving is planned.

PENDANT 3.3 EC Tank Mixes

PENDANT 3.3 EC may be used in combination with any registered herbicide. Consult labels for all directions, precautions and restrictions. Always check compatibility when applying in a tank mix and follow the most restrictive label.

SUNFLOWERS

General Instructions

PENDANT 3.3 EC may be applied preplant incorporated in the spring in sunflowers in all states.

PENDANT 3.3 EC alone may be applied preplant incorporated in the fall in sunflowers in the states of North Dakota, South Dakota and Minnesota only).

PENDANT 3.3 EC plus Eptam tank mixture may be applied preplant incorporated in the fall in sunflowers in the states of North Dakota and Minnesota only.

PENDANT 3.3 EC may be applied as preplant surface or preemergence application in no-till sunflowers in states of Colorado, Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas.

PENDANT 3.3 EC may be applied as a preemergence application in conventional tillage sunflowers in the states of Colorado, Kansas, Minnesota, Nebraska, North Dakota and South Dakota.

In the event of a crop loss due to weather conditions, sunflowers or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, DO NOT rework the soil deeper than the treated zone. DO NOT feed forage or graze livestock in treated sunflower fields. PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

Preplant Incorporated (Spring) – Apply PENDANT 3.3 EC up to 60 days prior to planting and incorporate within 7 days of application.

Preplant Incorporated (fall) – Apply PENDANT 3.3 EC or PENDANT 3.3 EC plus Eptam and immediately incorporate in late fall prior to planting sunflowers the following spring. Refer to **Incorporation Instructions** section (see table of contents for page number) for incorporation directions and the Eptam label for incorporation instructions. Destroy existing weeds before applying PENDANT 3.3 EC or PENDANT 3.3 EC tank mixture. The soil should be dry enough to permit good incorporation.

DO NOT apply PENDANT 3.3 EC plus Eptam tank mixture by air.

Apply PENDANT 3.3 EC or PENDANT 3.3 EC plus EPTAM tank mixture in the late fall when soil temperatures are 45°F or below but before the ground freezes. DO NOT apply when the air temperature is below 45°F.

Prior to sunflower planting in the spring, fields treated with PENDANT 3.3 EC or PENDANT 3.3 EC plus Eptam 7E should be received at least one shallow additional incorporation. Spring incorporation should be at an angle to the last tillage operation.

Preemergence (Spring) – PENDANT 3.3 EC may be applied immediately after planting. DO NOT apply PENDANT 3.3 EC postemergence since sunflowers exposed at the time of application will be killed or injured. PENDANT 3.3 EC is most effective in controlling weeds when adequate rainfall or irrigation is received within 7 days after application. Otherwise, a registered postemergence grass herbicide treatment may be required.

Existing living vegetation must be controlled at or before the application of PENDANT 3.3 EC.

A registered contact herbicide for use in sunflowers may be applied sequentially or in a tank mix with PENDANT 3.3 EC. Consult the contact herbicide label for all directions, precautions and restrictions.

Preemergence applications of PENDANT on conventional tillage sunflowers may increase the likelihood of crop injury and decrease herbicide performance compared to Preplant Incorporated applications. If dry conditions with limited precipitation exist or unseasonably cool temperatures following planting are forecasted, apply PENDANT 3.3 EC prior to planting and mechanically incorporate with tillage.

PENDANT 3.3 EC Use Rates in Sunflowers

Recommended use rates for PENDANT 3.3 EC are given in the following tables.

Broadcast Rate per Acre of PENDANT 3.3 EC for Spring Preplant Incorporated or Preemergence in Sunflowers (Pints per Acre)

Soil Texture	Southern States ¹	Northern States ¹	
		<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 1.8 pts	1.2 to 2.4 pts	2.4 pts
Medium	1.8 to 2.4 pts	1.8 to 3.0 pts	3.0 to 3.6 pts
Fine	1.8 to 3.6 pts	2.4 to 3.6 pts	3.6 pts

¹ See map at the end of this label for specific states.
Use the 3.6 pint rate for heavy clay soils.

Broadcast Rate per Acre of Pendant 3.3 EC for Fall Preplant Incorporated Application in Sunflowers¹ (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.8 to 3.0 pts	3.0 pts
Medium	2.4 to 3.6 pts	3.6 to 4.2 pts
Fine	3.0 to 4.2 pts	4.2 pts

¹ For use in North Dakota, South Dakota and Minnesota only
Use the high rate for each soil texture if heavy weed pressure is anticipated.

Broadcast Rate per Acre of PENDANT 3.3 EC Plus Eptam 7E For Fall Preplant Incorporated Tank Mixture Application in Sunflowers¹ (Pints per Acre)

Soil Texture	PENDANT 3.3 EC plus Eptam 7E	
	<3% Organic Matter	>3% Organic Matter
Coarse	1.2 to 2.4 + 2.25	2.4 + 2.25
Medium	2.8 to 3.0 + 2.25	3.0 to 3.6 + 2.25
Fine	2.4 to 3.6 + 2.25	3.6 + 2.25

¹ For use in North Dakota and Minnesota only.
Use the high rate for each soil texture if Heavy weed pressure is anticipated.
Read and strictly follow all precautions and instruction on the Eptam label.

No-Till Sunflowers for use in Colorado, Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota, and Texas

PENDANT 3.3 EC may be applied immediately after planting, or up to 30 days before planting. DO NOT apply PENDANT 3.3 EC postemergence since sunflowers exposed at the time of application will be killed. PENDANT 3.3 EC is most effective in controlling weeds when adequate rainfall or irrigations is received within 7 days after application. Otherwise, a registered postemergence grass herbicide treatment may be required.

Uniformly apply PENDANT 3.3 EC alone or in combination with other herbicides registered for gallonage for fields with history of heavy weed infestations or if the field has excessive crop residues.

Existing living vegetation must be controlled at or before the application of PENDANT 3.3 EC.

A registered contact herbicide for use in sunflowers may be applied sequentially or in a tank mix with PENDANT 3.3 EC. Consult the contact herbicide label for all directions, precautions and restrictions.

Broadcast Rate per Acre of PENDANT 3.3 EC Applied Preplant Surface or Preemergence in No-Till Sunflowers (Pints per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	3.0 to 3.6 pts	3.6 pts
Medium	3.6 pts	3.6 pts
Fine	3.6 pts	DO NOT USE

TOBACCO

General Instructions

PENDANT 3.3 EC may be applied preplant incorporated or as a layby application in transplanted tobacco.

In the event of a crop loss due to weather conditions, transplanted tobacco or any crop registered for PENDANT 3.3 EC preplant incorporated use can be replanted without adverse effects the same year. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.

PENDANT 3.3 EC is not recommended for use on peat or muck soils.

Use Methods and Timings

Preplant Incorporated - Apply PENDANT 3.3 EC with ground sprayer up to 60 days prior to transplanting tobacco and incorporate within 7 days of application.

Applied according to directions and under normal growing conditions, PENDANT 3.3 EC will not harm transplanted tobacco. Under stress conditions for plant growth such as cold/wet or hot/dry weather, PENDANT 3.3 EC can produce a temporary retardation of tobacco development.

Layby - PENDANT 3.3 EC may be applied as a directed spray following the last normal cultivation (layby), usually 4 to 6 weeks after transplanting tobacco. Apply PENDANT 3.3 EC in a 16 to 24-inch band in the middle of the row between the crop rows. The spray

should not contact tobacco plants. If the spray nozzles on the ends of the spray boom pass over the same row middle twice, use nozzles which apply one-half (1/2) the normal number of gallons per acre to prevent over-application. Layby applications can be applied in tobacco previously treated with herbicides registered for use in tobacco. Consult the labels of those herbicides for suggested treatments, rates to be used, and precautions or restrictions for use in tobacco and for follow crop restrictions.

PENDANT 3.3 EC is most effective in controlling weeds when adequate rainfall or irrigation is received within 7 days after application. **PENDANT 3.3 EC** will not control established weeds. **DESTROY EMERGED WEEDS PRIOR TO APPLICATION.** **DO NOT APPLY AS A BROADCAST SPRAY OVER THE TOP OF TOBACCO LEAF. CONTACT MAY CAUSE MALFORMED LEAVES.**

PENDANT 3.3 EC Use Rates in Transplanted Tobacco

Recommended use rates for **PENDANT 3.3 EC** alone is given in the following table.

Broadcast Rate¹ per Acre of PENDANT 3.3 EC Preplant Incorporated in Transplanted Tobacco (Pints per Acre)

Region	Soil Texture	PENDANT 3.3 EC (pints)
Maryland, Virginia, North Carolina, South Carolina, Georgia, Florida	COARSE	1.8 TO 2.4
	MEDIUM sandy clay loams, loams	1.8 TO 2.4
	silt loams, silts	2.4 to 3.0
	FINE	2.4 to 3.0
Other tobacco-growing states	COARSE	1.8 to 2.4
	MEDIUM	3.0 TO 3.6
	FINE	3.0 TO 3.6


Broadcast Rate¹ Per Acre of PENDANT 3.3 EC Layby Application in Transplant Tobacco (Pints per Acre)

Soil Texture	Rate ²
Coarse	1.2 to 1.8 pts.
Medium	1.8 to 2.4 pts.
Fine	1.8 to 2.4 pts.

¹For calculation of band treatment rate, see Spraying Instructions section (see table of contents for page number).

²The high rate for each soil texture above should be used if a registered herbicide treatment was not applied prior to layby.

REGIONAL MAP FOR RATE DETERMINATION



LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER THE MANUFACTURER OR SELLER SHALL BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS

WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW.

WARRANTY DISCLAIMER

THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER MANUFACTURER OR SELLER SHALL BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

¹ Accent SP, Basis Gold, Bladex, Extrazine II, Karmex, Londax, Lorox, Lorox Plus, Lexone, Matrix, Canopy and Preview are trademarks of E.I. du Pont de Nemours and Company.

² Beacon, Bicep, Bicep II, Dual, Dual II, Cotoran, Exceed, Caparol, Zorial are trademarks of Novartis Crop Protection.

³ Balance, Buctril, and Butyrac are trademarks of Rhone-Poulenc Ag Company.

⁴ Harness, Harness Xtra, Lariat, Bullet, Lasso, Micro-Tech, Partner WDG, Permit, Roundup, and Roundup Ready are trademarks of Monsanto Agricultural Products Company.

⁵ Surpass, Surpass 100, Eptam, Vernam, Gramoxone Extra and Arrosolo are trademarks of Zeneca Inc.

⁶ Sencor is a trademark of Bayer Corporation.

⁷ Banvel, Clarity, Facet 75 DF, Frontier, Guardsman, Laddok S-12, Marksman, Poast Plus, and SR Sethoxydim-resistant Field corn are trademarks of BASF Corporation.

⁸ Command is a trademark of FMC Corporation.

⁹ Bolero 8EC is a trademark of Valent.

¹⁰ Hornet, Python, WDG, Treflan, Scorpion III and Sonolan are trademarks of Dow Agrosciences

¹¹ LELY-ROTERA is a trademark of Lely.

¹¹ Registered Trademarks of BASF

¹² Pendant is a trademark of Agriliance, LLC