



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (H7505C)
 401 "M" St., S.W.
 Washington, D.C. 20460

EPA Reg. Number:
 1381-225

Date of Issuance:
 NOV 2 2007

Term of Issuance: Conditional

Name of Pesticide Product:
 Imidacloprid 1.0% G - Orn
 Insecticide

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Gary Halvorson
 Agrilience LLC
 P.O. Box 64089
 St. Paul, MN 55164

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.
 Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

1. Revise the EPA Registration Number to read, EPA Reg. No. "1381-225".

Signature of Approving Official:

 Dani Daniel
 Insecticide-Rodenticide Branch
 Registration Division (7505P)

Date:

 NOV - 2 2007

2 78

Page Two
1381-225

2. On page 2 under the headings Personal Protective Equipment and Agricultural Use Requirements replace waterproof gloves with the Agency's glove statement, which reads as follows: **Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.**

3. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(a). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact me at (703) 305-5409.

Enclosure

3 8 8

Imidacloprid 1.0% G – Orn Insecticide

A systemic insecticide for use on ornamental plants in greenhouses, nurseries and interior landscapes.

ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 1.0%

OTHER INGREDIENTS: 99.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
No specific antidote is available. Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

EPA Reg. No. 1381-

Distributed By:
Agrilience, LLC
P.O. Box 64089
St. Paul, MN 55164-0089

**ACCEPTED
with COMMENTS
In EPA Letter Dated
NOV - 2 2007**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

1381-225

EPA Est. No.

NET CONTENTS

PROP01/15/07
05/24/07

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment wash waters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

[Paper Bags]: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Plastic Container]: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Imidacloprid 1.0% G – Orn Insecticide will control insects on ornamental plants grown in nurseries and greenhouses. Imidacloprid 1.0% G – Orn Insecticide is a systemic insecticide that is translocated upward within the plant. For best results, Imidacloprid 1.0% G – Orn Insecticide must be applied where the roots of the target plants can absorb the active ingredient. After application, irrigate moderately but thoroughly being sure to avoid leaching and runoff from the container for at least three irrigations or 10 days whichever is longer.

Use Precautions:

- Do NOT graze treated areas or use clippings from treated areas for feed or forage.
- Do NOT apply to soils which are waterlogged, or saturated. This condition will not allow penetration of water into the root zone of the plants.
- Do NOT over-irrigate or allow excessive runoff to occur following an application of Imidacloprid 1.0% G – Orn Insecticide.
- Do NOT apply more than 40 lbs. of Imidacloprid 1.0% G – Orn Insecticide per acre per year.
- On plants with a production cycle of less than one year, application is not to exceed a frequency of more than once each 16 weeks for a particular plant. On stock plants and woody crops with a production cycle of greater than one year, application may not exceed once a year.

Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for which the active ingredient have been established, a 12-month plant-back interval should be observed.

Insect Resistance: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

For resistance management purposes, a foliar application of any chloronicotinyl insecticide following Imidacloprid 1.0% G – Orn Insecticide soil application in the same crop is not recommended.

Woody Perennials: Protection in woody perennials is slower than in herbaceous species and a delay of 2 or more weeks should be expected, with longer delays for larger plants. Because of this, applications to woody perennials should be made well in advance of expected insect activity.

Incorporation: Incorporation of Imidacloprid 1.0% G – Orn Insecticide can be achieved by cultivation, irrigation, rainfall, mechanical placement, or by using mechanical soil or media mixing equipment.

Bark Media: Imidacloprid 1.0% G – Orn Insecticide treatments to media with 30 - 50% or more bark content may confer a shorter period of protection.

APPLICATION TO GRASSY AREAS IN NURSERIES

NOTE: Imidacloprid 1.0% G – Orn Insecticide may NOT be used on commercial sod farms.

Imidacloprid 1.0% G – Orn Insecticide will control soil-inhabiting pests in grassy areas in or around nurseries such as under or around field or container grown plants, on roadways or other grassy areas. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made preceding or during the egg laying period. For best results, make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Pests Controlled:

- | | | |
|--|--|---|
| Annual bluegrass weevil (<i>Hyperodes</i> spp.) | Asiatic garden beetle (<i>Maladera castanea</i>) | Billbugs (<i>Spherophorus</i> spp.) |
| European chafer (<i>Rhizotroqus majalis</i>) | Green June beetle (<i>Cotinis nitida</i>) | Japanese beetle (<i>Popillia japonica</i>) |
| May or June beetle (<i>Phyllophaga</i> spp.) | Mole Crickets (<i>Scapteriscus</i> spp.) | Oriental beetle (<i>Anomala orientalis</i>) |
| Black turfgrass ataenius (<i>Ataenius spretulus</i> and <i>Aphodius</i> spp.) | | |
| Northern and Southern masked chafers (<i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i>) | | |

Pests Suppressed:

- | | |
|----------|------------------|
| Cutworms | Hairy Chinchbugs |
|----------|------------------|

Grassy Area Use Precautions:

- Applications must NOT exceed a total of 40 lbs. (0.4 lb of active ingredient) per acre per year.
- Applications should NOT be made when grassy areas are waterlogged or the soil is saturated with water because adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

APPLICATION TO ORNAMENTALS GROWN IN FLATS, BENCHES, BEDS AND CONTAINERS

Imidacloprid 1.0% G – Orn Insecticide will suppress or control the following insect pests when applied to ornamentals grown in flats, benches, beds and containers in greenhouses and nurseries.

- | | | |
|--|--|-----------------------------------|
| Adelgids | Aphids | Armored scale (suppression) |
| Fungus gnats (larvae only) ¹ | Japanese beetles (adults) | Lacebugs |
| Leafminers | Mealybugs | Psyllids |
| Root Weevil | Soft scale | Thrips (suppression) ³ |
| Whiteflies | Leaf beetles (including Elm and Viburnam leaf beetles) | |
| Leafhoppers (including glassy-winged sharpshooter) | Complex (such as Black Vine Weevil, Apopka Weevil, Citrus Root Weevil ²) | |

¹ Fungus gnat larvae in the soil will be controlled by incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of Imidacloprid 1.0% G – Orn Insecticide from a healthy root system translocating the active ingredient up into the plant.

² Citrus Root Weevil: For use on non-bearing citrus nursery stock.

³ Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.

HERBACEOUS AND WOODY PERENNIAL SPECIES						
USE PATTERN	Container Size (inches)	APPLICATION RATE				APPLICATION INSTRUCTIONS
		Bulk Application		Topical Application		
		Pots/cu. yd. of mix	Bulk Rate lbs./cu. Yard	Topical Rate (level teaspoons)	Grams / Pot	
Herbaceous Species (one or two plants per pot)	2.0 to 3.0	4500 to 3200	7.0 to 3.5	1/8 to 1/4	0.5 to 1.0	Bulk Rate Application: Mix the required amount of Imidacloprid 1.0% G – Orn Insecticide per cubic yard of medium. Mix thoroughly before filling pots and
	3.5 to 5.0	1600 to 1300	3.9 to 3.0	1/4 to 1/3	1.0 to 1.4	
	5.5 to 7.0	900 to 400	1.8 to 1.2	1/3 to 1/2	1.4 to 2.0	
	8.0 to 10.0	200 to 160	1.0 to 0.8	1/2 to 2/3	2.0 to 2.7	
	11.0 to 12.0	80 to 60	0.8 to 0.5	2/3 to 1	2.7 to 4.1	

Woody Perennial Species or Herbaceous Species (Three or more plants per pot)	2.0 to 3.0	4500 to 3200	9.0 to 7.0	¼ to 1/3	1 to 1.3	transplanting. Topical Application: May be made at transplanting, repotting, or on established containerized plants. Irrigate moderately after application into the root zone. Do not allow significant leaching and runoff for at least 3 irrigations or 10 days, whichever is longer.
	3.5 to 5.0	1600 to 1300	6.0 to 4.0	1/3 to ½	1.3 to 2.0	
	5.5 to 7.0	900 to 400	2.5 to 2.0	½ to 2/3	2.0 to 2.7	
	8.0 to 10.0	200 to 160	1.5 to 1.0	2/3 to 1	2.7 to 4.1	
	11.0 to 12.0	80 to 60	1.0 to 0.8	to 1 ½	4.1 to 6.1	

CONTAINERIZED PLANTS					
Container Size (Gallons)	APPLICATION RATE				APPLICATION INSTRUCTIONS
	Bulk Application		Topical Application		
	Pots/cu. yd. of mix	Bulk Rate lbs./cu. Yard	Topical Rate (level teaspoons)	Grams / Pot	
1	260 to 230	5.0 to 2.5	1 ¼ to 2 ½	5 to 10	For optimum control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.
2	125 to 100	4.4 to 2.2	2 ½ to 5.0	10 to 20	
3	70 to 60	3.3 to 1.7	3 2/3 to 7 1/3	15 to 30	
5	50 to 30	2.0 to 1.0	4 ½ to 9.0	18.5 to 37	

ORNAMENTAL PLANTS GROWN IN FLATS, BENCHES OR BEDS	
Application Rate	Application Instructions
Apply 15 ounces per 1000 sq. ft.	Apply as a broadcast treatment and incorporate into the soil or media before planting. On established plants, irrigate moderately after application to move the active ingredient into the root zone. Minimize leaching and runoff for at least 3 irrigations or 10 days, whichever is longer.

1 level tsp. of Imidacloprid 1.0% G – Orn Insecticide = 4.1 grams 3 teaspoons = 1 tablespoon

FOR CONTROL OF WHITE GRUB LARVAE [†] ON ORNAMENTALS IN FIELD AND FOREST NURSERIES	
APPLICATION RATE	APPLICATION INSTRUCTIONS
Use 45 oz. per 1000 sq. ft. of row or 3000 sq. ft. For grub control in areas of turf, apply as a broadcast application using 45 oz. per 3000 sq. ft.	Vegetation in the area to be treated should be mowed to a height of 3 inches or less prior to application. Mowing to the lowest possible height will ensure greater consistency of control. Apply May through July. For optimum control, treatment should be followed by rainfall or irrigation. Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap.
[†] White grub larvae include Japanese beetle, Masked Chafers, European Chafer, Oriental beetle, and Asiatic Garden beetle	

LIMITATION OF LIABILITY

IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, TO THE EXTENT ALLOWED BY LAW.

WARRANTY DISCLAIMER

THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD

TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. UNINTENDED CONSEQUENCES, INCLUDING, BUT NOT LIMITED TO, CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER. IN NO CASE SHALL MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.