



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (H7505C)  
 401 "M" St., S.W.  
 Washington, D.C. 20460

EPA Reg.

Number:

1381-227

Date of Issuance:

NOV 2 2007

Term of Issuance: Conditional

Name of Pesticide Product:

Imidacloprid 75% Turf WSP  
Insecticide

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Mr. Gary Halvorson  
 Agriliance LLC  
 P.O. Box 64089  
 St. Paul, MN 55164

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

1. Revise the EPA Registration Number to read, EPA Reg. No. "1381-227".

Signature of Approving Official:

Dani Daniel  
 Insecticide-Rodenticide Branch  
 Registration Division (7505P)

Date:

NOV - 2 2007

2 7 18

Page Two  
1381-227

2. As this label as written is not a 100 percent repack. In order to come into compliance, delete everything from the label starting with page 6. A one hundred percent repack can only contain the language of the parent label.

3. On page five of this label, add the following statement to the warranty statement **Limitation of Liability section** at the beginning of first sentence. Also add the same statement at the beginning of the second sentence of the **Warranty Disclaimer** section: **"To the extent consistent with applicable law..."**

4. Submit two copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(a). Your release for shipment of the product constitute acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact me at (703) 305-5409.

Enclosure

3718

# Imidacloprid 75% WSP Turf Insecticide

Systemic and foliar insect control in grapes, turfgrass (including sod farms), and on fruit and nut trees, landscape ornamentals, and interior plantscapes.

### ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine .....	75.0%
<b>OTHER INGREDIENTS:</b> .....	<b>25.0%</b>
<b>TOTAL:</b> .....	<b>100.0%</b>

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not do anything by mouth an unconscious person</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
No specific antidote is available. Treat the patient symptomatically.	

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

May be fatal if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse. Keep children and pets away from treated area until spray is dry.

EPA Reg. No. 1381-

**Distributed By:**  
Agrilience, LLC  
P.O. Box 64089  
St. Paul, MN 55164-0089

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**NOV - 2 2007**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

1381-227

EPA Est. No. ....

NET CONTENTS .....

PROP 6145/07

05/29/07

4 8 18

### PERSONAL PROTECTIVE EQUIPMENT (PPE):

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on the foliage of blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This product is toxic to wildlife.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**NOT FOR USE IN COMMERCIAL GREENHOUSES, NURSERIES, ON GRASSES GROWN FOR SEED, OR ON COMMERCIAL FRUIT AND NUT TREES**

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticide. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to the uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

- Keep children and pets off treated areas until dry.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE [For product packaged in plastic containers]:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**PESTICIDE STORAGE [For product packaged in Water Soluble Packaging]:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Exposure to moisture or excessive handling of water soluble packets may cause breakage.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL [For product packaged in plastic container]:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL [For product packaged in Water Soluble Packaging]:** Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### SPRAY DRIFT MANAGEMENT

**Aerial applications are permitted ONLY on agricultural use grapes for Grapeleaf skeletonizer suppression.**

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

#### No-Spray Zone Requirements for Foliar Applications

Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams, marshes or natural ponds; estuaries and commercial fish farm ponds.

#### Mixing and Loading Requirements

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

**Aerial Applications**

The spray boom should be mounted on the aircraft so as to minimize the drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

**Importance of Droplet Size**

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

**Wind Speed Restrictions**

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any give wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

**Restrictions During Temperature Inversions**

*Because the potential for spray drift is high during temperature inversions, do NOT make aerial or ground applications during temperature inversions.* Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

**Airblast (Air Assist) Specific Recommendations for Vineyards**

Airblast sprayers carry droplets into the canopy of vines via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e. turn off sprayer when turning at end rows);
- Only spray inward, toward the vineyard, for applications to the outside rows.

**RUNOFF MANAGEMENT**

Do not cultivate within 10 feet of the aquatic areas to allow growth of vegetative filter strip. When used on erodible soils, best management practices for minimizing runoff should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

**ENDANGERED SPECIES NOTICE**

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

**RESISTANCE MANAGEMENT**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

Imidacloprid 75% WSP Turf Insecticide contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by Imidacloprid 75% WSP Turf Insecticide and to other Group 4A products.

The active ingredient in Imidacloprid 75% WSP Turf Insecticide is a member of the neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of Imidacloprid 75% WSP Turf Insecticide and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Etigra LLC strongly

encourages the rotation to a block of applications with effective products of a different mode before using additional applications of neonicotinoid products. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Foliar applications of Imidacloprid 75% WSP Turf Insecticide or other Group 4A products from the neonicotinoid chemical class should not be used on crops previously treated with a long-residual, soil-applied products from the neonicotinoid chemical class.

Other Group 4A, neonicotinoid products used as foliar treatments include: Actara<sup>®</sup>, Assail<sup>®</sup>, Calypso<sup>®</sup>, Centric<sup>®</sup>, Intruder<sup>™</sup>, Leverage<sup>®</sup>, Provado<sup>®</sup>, and Trimax<sup>™</sup>. Other 4A Group, neonicotinoid products used as soil treatment include: Admire<sup>®</sup> and Platinum<sup>®</sup>.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

#### Use Precautions:

- Keep children and pets off treated areas until dry.
- Do NOT apply through any type of irrigation system.
- Do NOT apply by air. [Note to Reviewer: This statement will not appear on final sub-label for agricultural use grapes]
- Do NOT graze treated areas or use clippings from treated areas for feed or forage.
- Do NOT apply Imidacloprid 75% WSP Turf Insecticide to soils that are waterlogged or saturated and avoid runoff or puddling of irrigation water following application.
- Do NOT allow lechate to run out for the first 10 days after application or reduced efficacy may result.
- Do NOT exceed a total of 8.6 oz. / Acre per year (0.4 lb. AI/A).

#### LIMITATION OF LIABILITY

IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT, TO THE EXTENT ALLOWED BY LAW.

#### WARRANTY DISCLAIMER

THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. UNINTENDED CONSEQUENCES, INCLUDING, BUT NOT LIMITED TO, CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER. IN NO CASE SHALL MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS

8 7 18

ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.



9 8 18

# FOR PRODUCT PACKAGED IN WSP

## MIXING AND APPLICATION INSTRUCTIONS

Inside each foil pouch is a clear, water soluble inner packet containing the Imidacloprid 75% WSP Turf Insecticide. To prepare a solution, remove the outer foil pouch and drop the required number of unopened clear water soluble packets into the spray tank while filling with water to the desired level. Be sure to agitate while mixing and depending on the amount of agitation and the water temperature, the packets should completely dissolve within a few minutes of being added to the water. Note that cooler water temperatures increase the time needed for the inner packet to completely dissolve.

### Mixing Precautions:

- Do NOT allow packets to become wet prior to adding to the tank.
- Do NOT handle the clean inner packets with wet hands or wet gloves.
- Do NOT use this product in a tank-mix with products that contain Boron or release free chlorine. Combining these products will result in a plastic that is not soluble in water or solvents (such as diesel oils, kerosene, gasoline or alcohol). NOTE: Chlorinated water may be used.
- Because the water soluble packets are not soluble in petroleum-based liquids, do NOT attempt to use Imidacloprid 75% WSP Turf Insecticide water soluble packets directly in diesel oils or summer spray type oils such as those used in ULV or LV applications.
- Rough handling of the packets may cause breakage.
- Reseal outer carton to protect remaining packets.

**Tank Mixes:** Imidacloprid 75% WSP Turf Insecticide has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. If Imidacloprid 75% WSP Turf Insecticide is not known to be compatible with your particular tank mix partners, compatibility should be checked using the correct proportion of products in the following small jar test:

- 1) Add proportionate amount of each ingredient in the appropriate order to a pint or a quart jar;
- 2) Cap and shake for 5 minutes;
- 3) Let set for 5 minutes.

Poor mixing or formation of precipitates that do not readily re-disperse indicates an incompatible mixture that should not be used. For further information, contact your local Etigra, LLC representative.

**Mixing Instructions:** The enclosed packets containing Imidacloprid 75% WSP Turf Insecticide are water-soluble and will completely dissolve in water. The proper mixing procedure for Imidacloprid 75% WSP Turf Insecticide alone or in tank mix combinations with other pesticides is:

1. Fill the spray tank ¼ to 1/3 full with clean water.
2. While recirculating and with the agitator running, add the required number of unopened Imidacloprid 75% WSP Turf Insecticide packets.
3. The packets should completely dissolve in 5 to 10 minutes; allow sufficient time for thorough mixing.
4. Continue to fill spray tank with water until ½ full.
5. If applicable, add remaining tank mix components in the following order: wettable powders, flowables, and emulsifiable concentrates. Ensure good agitation as each component is added. Do not add a tank mix component until the previous component is thoroughly mixed.
6. Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

## TURFGRASS

Imidacloprid 75% WSP Turf Insecticide will control soil-inhabiting pests in grassy areas such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields and sod farms. The need for an application can be based on historical monitoring of the

site, previous records or experiences, current season adult trapping or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made proceeding or during the egg laying period. For best results, make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

**Use Precautions:**

- Applications must NOT exceed a total of 8.6 oz. (0.4 lb of active ingredient) per acre per year.
- Applications should NOT be made when grassy areas are waterlogged or the soil is saturated with water because adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- Avoid mowing treated areas until after sufficient rainfall or irrigation has occurred in order to maintain the uniformity of the application.

**Application Instructions:**

Apply Imidacloprid 75% WSP Turf Insecticide in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

Pest	Application Rate	Specific Instructions
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass atanius Cutworms† European Chafer Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	1.6 oz. (1 packet) per 11,000 sq. ft.	<b>Grubs, European Crane Fly, billbugs and annual bluegrass weevil:</b> For best results make applications prior to egg hatch of the target pest. <b>Cinchbugs:</b> Make applications prior to the hatching of the first instar nymphs. <b>Mole Crickets:</b> Make applications prior to or during the peak egg hatching period. When adults or large nymphs are present and actively tunneling, Imidacloprid 75% WSP Turf Insecticide should be accompanied by a curative insecticide. <b>NOTE:</b> For best results, the active ingredient must be moved through the thatch by irrigation or rainfall occurring within 24 hours after application.
Cinchbugs† Mole Crickets	8.6 oz. per Acre or 4 level Tbs. per 1,000 sq. ft.	
† Suppression only.		

**ORNAMENTALS, GROUNDCOVERS AND INTERIOR PLANTSCAPES**

Imidacloprid 75% WSP Turf Insecticide is a systemic insecticide that may be applied to ornamentals, groundcovers and interior plantscapes in and around industrial and commercial buildings and residential areas. The insecticide is translocated upward into the plant system and for best results must be placed where the growing portions of the target plant can absorb the active ingredient. When applicable, adding a fertilizer containing nitrogen into the spray solution may enhance plant uptake of Imidacloprid 75% WSP Turf Insecticide.

**Rotational Crops:**

As soon as practical following the last application, treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed. **NOTE:** Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed.

**Ant Management Programs:**

Imidacloprid 75% WSP Turf Insecticide may be used to limit the honeydew available as a food source for ant populations when controlling aphids, scale insects, mealy bugs and other sucking pests on ornamentals. Imidacloprid 75% WSP Turf Insecticide applications may be supplemented with bait traps, residual sprays and other methods to further reduce the unwanted ant population.

**Insect Resistance:**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

**Woody Perennials:**

Protection in woody perennials is slower than in herbaceous species and a delay of 2 or more weeks should be expected, with longer delays for larger plants. Because of this, applications to woody perennials should be made well in advance of expected insect activity.

**Bark Media:**

Imidacloprid 75% WSP Turf Insecticide treatments to media with 30 - 50% or more bark content may confer a shorter period of protection.

**FOLIAR AND BROADCAST APPLICATIONS**

Imidacloprid 75% WSP Turf Insecticide may be applied as a broadcast or foliar application to trees (including non-bearing fruit and nut trees), shrubs, evergreens, flowers, foliage plants, ground covers, interior landscapes and vegetable plants intended for resale.

**Application Instructions:**

Apply Imidacloprid 75% WSP Turf Insecticide in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

NOTE: When making foliar applications to plants with hard-to-wet foliage such as holly, pine or ivy, use of a spreader / sticker is recommended.

Pest	Application Method	Application Rate	Specific Instructions
Adelgids Aphids Japanese beetle (adult) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips† Whiteflies	Foliar	1.6 oz. (1 packet) per 300 gal. of water	Make applications prior to establishment of large pest populations and retreat as necessary. NOTE: Applying Imidacloprid 75% WSP Turf Insecticide foliarly after a soil application in the same crop is not recommended for resistance management purposes.
White grub larvae (such as Japanese beetle larvae, chafers, Phyllophaga spp., Asiatic garden beetle and Oriental beetle)	Broadcast	1.6 oz. (1 packet) per 11,000 sq. ft.	Mix the recommended amount of Imidacloprid 75% WSP Turf Insecticide in sufficient water to uniformly cover the area being treated using at least 2 gallons of water per 1000 sq. ft. For best results, incorporate the Imidacloprid 75% WSP Turf Insecticide into the upper soil profile by irrigating after the application is made.
† Suppression only.			

**SOIL INJECTION AND DRENCH APPLICATIONS**

Application Site	Recommended Rate	Application Instructions	Pests Controlled
<p>Trees</p> <p>NOTE: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p>	<p>1.6 oz. (1 packet) per 24 – 48 inches of cumulative trunk diameter (DBH)</p>	<p><b>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York.</b></p> <p><b>GRID SYSTEM:</b> Holes should be spaced on 2.5 ft centers, in a grid pattern, extending to the drip line of the tree.</p> <p><b>CIRCLE SYSTEM:</b> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) extending in from the drip line of the tree.</p> <p><b>BASAL SYSTEM:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.</p> <p><b>SOIL DRENCH:</b> Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.</p>	<p>Adelgids Aphids Armored Scale<sup>†</sup> Black vine weevil larvae Eucalyptus Longhorned Borers Flatheaded Borers (including bronze birch and alder borers) Japanese Beetles (adults) Lacebugs Leaf Beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter)</p>
<p>Shrubs</p>	<p>0.1 – 0.2 fl. oz. per foot of shrub height</p>	<p><b>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York.</b></p> <p>Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Using a minimum of 4 holes per shrub, apply to individual plants maintaining a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days.</p> <p><b>SOIL DRENCH:</b> Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.</p>	<p>Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips<sup>†</sup> White grub larvae Whiteflies</p>
<p>Flowers and Ground Cover</p>	<p>0.46 - 0.60 fl. oz. per 1000 sq. ft.</p>	<p>Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.</p>	
<p><sup>†</sup> Suppression only of these species.</p>			

**POME FRUIT IN AND AROUND RESIDENTIAL AREAS**

Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear) Quince

- Pre-Harvest Interval (PHI): 7 days
- Reapplication Interval: At least 10 days
- Maximum Applications per Year: 5

Pest	Ounces per 300 gal. of Water	Ounces per Acre <sup>‡</sup>	Specific Instructions
Aphids (except Woolly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs <sup>†</sup> San Jose Scale <sup>†</sup>	1.6 (1 packet)	2.1	Apply as a foliar spray as needed after petal-fall is complete. <b>Rosy Apple Aphid:</b> Apply prior to leaf rolling caused by the pest. <b>Leafhopper:</b> For late season (preharvest) control, apply while most leafhoppers are in the nymphal stage. <b>Leafminer:</b> Make first application as soon as petal-fall is complete for control of first generation, with best results occurring when the application is made at the earliest possible time. For succeeding generations, best results occur when applications are made early in the adult flight against egg and early instar larvae. If generations are overlapping or severe pressure continues, a second application may be necessary after 10 days. A single application may result in suppression only. <b>NOTE:</b> Imidacloprid 75% WSP Turf Insecticide will not control late stage larvae. <b>Mealybug:</b> For best results be sure to have good spray coverage of the trunk and scaffolding limbs or other nesting sites. <b>San Jose Scale:</b> Time applications to the crawler stage and treat each generation.

<sup>†</sup> Not permitted for control on pears in California.  
<sup>‡</sup> The amount of Imidacloprid 75% WSP Turf Insecticide required per acre depends on tree size and volume of foliage. The listed rate per acre is based on a standard of 400 gallons of dilute spray per acre for large trees.

**PECANS IN AND AROUND RESIDENTIAL AREAS**

**NOT PERMITTED IN CALIFORNIA unless otherwise directed by supplemental labeling**

- Reapplication Interval: At least 10 days
- Maximum Applications per Year: 3
- Maximum Imidacloprid 75% WSP Turf Insecticide allowed per Year: 6.3 oz. / Acre

Pest	Ounces per 300 gal. of Water	Ounces per Acre <sup>‡</sup>	Specific Instructions
Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1.6 (1 packet)	2.1	Apply as a foliar spray as pest pressure builds but before infestation is extremely heavy. Two applications at a 10 – 14 day interval may be required to achieve control. For best results, thorough and uniform coverage is necessary. Coverage may be improved through the use of an organosilicone-based spray adjuvant.

<sup>‡</sup> The amount of Imidacloprid 75% WSP Turf Insecticide required per acre depends on tree size and volume of foliage. The listed rate per acre is based on a standard of 400 gallons of dilute spray per acre for large trees.

**GRAPES  
ORNAMENTAL AND AGRICULTURAL USE**

- Reapplication Interval: At least 14 days
- Maximum Imidacloprid 75% WSP Turf Insecticide allowed per Year: 2.0 oz. / Acre

<b>Ornamental Grapes In and Around Industrial and Commercial Buildings and Residential Areas</b>			
<b>Pest</b>	<b>Ounces per 300 gal. of Water</b>	<b>Ounces per Acre<sup>†</sup></b>	<b>Specific Instructions</b>
Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.6 (1 packet)	1.0	Apply as a foliar spray using 200 gallons of water per acre.

<b>Agricultural Use Grapes including American bunch grape, Muscadine grape and Vinefera grape</b>		
<b>Pest</b>	<b>Application Rate</b>	<b>Specific Instructions</b>
Leafhoppers/Sharpshooters Mealybugs	0.8-1.0 oz. per Acre	<b>Pre-Harvest Interval (PHI): 0 day</b> <b>Minimum interval between applications: 14 days</b> <b>Maximum Imidacloprid 75% WSP Turf Insecticide allowed per crop season: 2.0 oz. per Acre (0.1 lb. A/A)</b>
Grapeleaf skeletonizer <sup>1</sup>	1.0 oz. per Acre	

<sup>†</sup> Control of Grapeleaf skeletonizer can be expected from ground applications that provide thorough coverage of foliage. Aerial applications may provide suppression.

## FOR PRODUCT PACKAGED IN NON-WSP

### TURFGRASS

Imidacloprid 75% WSP Turf Insecticide will control soil-inhabiting pests in grassy areas such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields and sod farms. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made proceeding or during the egg laying period. For best results, make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

#### Use Precautions:

- Applications must NOT exceed a total of 8.6 oz. (0.4 lb of active ingredient) per acre per year.
- Applications should NOT be made when grassy areas are waterlogged or the soil is saturated with water because adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.
- Avoid mowing treated areas until after sufficient rainfall or irrigation has occurred in order to maintain the uniformity of the application.

#### Application Instructions:

Apply Imidacloprid 75% WSP Turf Insecticide in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

Pest	Level Teaspoons per 1000 sq. ft.	Ounces per Acre	Specific Instructions
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms <sup>†</sup> European Chafer Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	2.5 – 4.0	5.4 – 8.6	<p><b>Grubs, European Crane Fly, billbugs and annual bluegrass weevil:</b> For best results make applications prior to egg hatch of the target pest.</p> <p><b>Cinchbugs:</b> Make applications prior to the hatching of the first instar nymphs.</p> <p><b>Mole Crickets:</b> Make applications prior to or during the peak egg hatching period. When adults or large nymphs are present and actively tunneling, Imidacloprid 75% WSP Turf Insecticide should be accompanied by a curative insecticide.</p> <p><b>NOTE:</b> For best results, the active ingredient must be moved through the thatch by irrigation or rainfall occurring within 24 hours after application.</p>
Cinchbugs <sup>†</sup> Mole Crickets	4.0	8.6	
<sup>†</sup> Suppression only.			

1 level teaspoon = 1.4 grams of Imidacloprid 75% WSP Turf Insecticide  
3 level teaspoons = 1 level tablespoon

**ORNAMENTALS, GROUNDCOVERS AND INTERIOR PLANTSCAPES**

Imidacloprid 75% WSP Turf Insecticide is a systemic insecticide that may be applied to ornamentals, groundcovers and interior plantscapes in and around industrial and commercial buildings and residential areas. The insecticide is translocated upward into the plant system and for best results must be placed where the growing portions of the target plant can absorb the active ingredient. When applicable, adding a fertilizer containing nitrogen into the spray solution may enhance plant uptake of Imidacloprid 75% WSP Turf Insecticide.

**Rotational Crops:**

As soon as practical following the last application, treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed. NOTE: Cover crops for soil building or erosion control may be planted at any time, but do not graze or harvest for food or feed.

**Ant Management Programs:**

Imidacloprid 75% WSP Turf Insecticide may be used to limit the honeydew available as a food source for ant populations when controlling aphids, scale insects, mealy bugs and other sucking pests on ornamentals. Imidacloprid 75% WSP Turf Insecticide applications may be supplemented with bait traps, residual sprays and other methods to further reduce the unwanted ant population.

**Insect Resistance:**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

**Woody Perennials:**

Protection in woody perennials is slower than in herbaceous species and a delay of 2 or more weeks should be expected, with longer delays for larger plants. Because of this, applications to woody perennials should be made well in advance of expected insect activity.

**Bark Media:**

Imidacloprid 75% WSP Turf Insecticide treatments to media with 30 - 50% or more bark content may confer a shorter period of protection.

**FOLIAR AND BROADCAST APPLICATIONS**

Imidacloprid 75% WSP Turf Insecticide may be applied as a broadcast or foliar application to trees (including non-bearing fruit and nut trees), shrubs, evergreens, flowers, foliage plants, ground covers, interior plantscapes and vegetable plants intended for resale.

**Application Instructions:**

Apply Imidacloprid 75% WSP Turf Insecticide in sufficient water to provide adequate distribution in the treated area. Use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

NOTE: When making foliar applications to plants with hard-to-wet foliage such as holly, pine or ivy, use of a spreader / sticker is recommended.

Pest	Application Method	Application Rate		Specific Instructions
Adelgids Aphids Japanese beetle (adult)	Foliar	Imidacloprid 75% WSP Turf Insecticide	Water	Make applications prior to establishment of large pest populations and retreat as necessary.



Pest	Application Method	Application Rate		Specific Instructions
Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips <sup>†</sup> Whiteflies		0.25 tsp. 0.50 tsp. 1.0 tsp. 2.5 tsp. 5.0 tsp.	2.5 gal. 5.0 gal. 10.0 gal. 25.0 gal. 50.0 gal.	NOTE: Applying Imidacloprid 75% WSP Turf Insecticide foliarly after a soil application in the same crop is not recommended for resistance management purposes.
White grub larvae (such as Japanese beetle larvae, chafers, Phyllophaga spp., Asiatic garden beetle and Oriental beetle)	Broadcast	3 Tbs. + 1 tsp.	100.0 gal.	Mix the recommended amount of Imidacloprid 75% WSP Turf Insecticide in sufficient water to uniformly cover the area being treated using at least 2 gallons of water per 1000 sq. ft. For best results, incorporate the Imidacloprid 75% WSP Turf Insecticide into the upper soil profile by irrigating after the application is made.
† Suppression only.				

1 level teaspoon = 1.4 grams of Imidacloprid 75% WSP Turf Insecticide

3 level teaspoons = 1 level tablespoon

### SOIL INJECTION AND DRENCH APPLICATIONS

Application Site	Recommended Rate	Application Instructions	Pests Controlled
Trees  NOTE: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.	0.7 – 1.4 level teaspoons per inch of trunk diameter (DBH)  or  1 – 2 oz. per 30 cumulative inches of trunk diameter (DBH)	<b>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York.</b>  <b>GRID SYSTEM:</b> Holes should be spaced on 2.5 ft centers, in a grid pattern, extending to the drip line of the tree.  <b>CIRCLE SYSTEM:</b> Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) extending in from the drip line of the tree.  <b>BASAL SYSTEM:</b> Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.  Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.	Adelgids Aphids Armored Scale <sup>†</sup> Black Vine Weevil larvae Emerald Ash Borer (adult) <sup>†</sup> Eucalyptus Longhorned Borers Flatheaded Borers (including bronze birch and alder borers) Japanese Beetles (adults) Lacebugs Leaf Beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter)
		<b>SOIL DRENCH:</b> Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.	Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips <sup>†</sup> White Grub larvae Whiteflies
Shrubs	0.7 – 1.4 level teaspoons per foot of shrub height  or  1 – 2 oz. per 30 cumulative feet of shrub height	<b>SOIL INJECTION – No Soil Injection Application allowed in Nassau or Suffolk counties of New York.</b>  Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Using a minimum of 4 holes per shrub, apply to individual plants maintaining a low pressure and use sufficient solution for distribution of the liquid into the treatment zone.  Keep the treated area moist for 7 to 10 days.	Leafminers Mealybugs Pine Tip Moth larvae Psyllids Royal Palm Bugs Sawfly larvae Soft Scales Thrips <sup>†</sup> White Grub larvae Whiteflies
		<b>SOIL DRENCH:</b> Remove plastic or any other barrier that will stop solution from reaching the root zone. Uniformly apply around the base of the tree, direct to the root zone as a drench in no less than 10 gallons of water per 1000 square feet.	

Application Site	Recommended Rate	Application Instructions	Pests Controlled
Flowers and Ground Cover	3 - 4 level teaspoons per 1000 sq. ft.	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application.	

† Suppression only of these species.  
 ‡ Make a minimum of two applications per season with the first application in late May / early June or when adult beetles begin to fly. Make the second application approximately one month after the first application.

1 level teaspoon = 1.4 grams of Imidacloprid 75% WSP Turf Insecticide  
 3 level teaspoons = 1 level tablespoon