

1381-231

02/15/2008

1/3



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 15 2008

Ms. Glenda Haage
Winfield Solutions
P. O. Box 64589 -- MS 5705
Shoreview, MN 55126-2910

Subject: Additional Crops to Supplemental Label
Nitro Shield
EPA Reg. No. 1381-231
Your Submission date, January 22, 2008

Dear Ms. Haage:


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable provided you incorporate this supplemental label into the main label at the next printing or within 18 months, whichever comes first.

A stamped copy is enclosed for your records. Submit two (2) copies of your final printed labeling before you release the product for shipment. If there are questions call me at 703 305-5409.

Sincerely,

151

Dani Daniel
Insecticide-Rodenticide Branch
Registration Division 7505C



SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

When using this product as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

NITRO SHIELD

EPA Reg. No. 1381-231

KEEP OUT OF REACH OF CHILDREN.

CAUTION

In case of an emergency involving this product, Call CHEMTREC toll free at 1-800-424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

EARLY SEASON PROTECTION AGAINST SUCKING INSECTS

CROP	PEST	RATE	COMMENT
Flax, crambe, and borage	wireworms, seed corn maggots and flea beetles	25.6 fluid ounces per hundredweight seed	
Safflower	wireworms	0.25 to 0.50 mg a.i./seed (One fluid ounce of this product contains 17.7g imidacloprid.)	
Sunflower	wireworms, seed corn maggots and flea beetles	0.25 to 0.50 mg a.i./seed. (One fluid ounce of this product contains 17.7g imidacloprid.)	Sunflower seed treated in California must be destined for planting in states other than California and is not to be planted in California.

Flax, sunflower, crambe, and borage (For end use application at agricultural establishments): To provide early season protection of seedlings against wireworms, seed corn maggots and flea beetles, apply using an HCBT or a batch treater at 12.8 fluid ounces per hundredweight of seed.

Safflower (For end use application at agricultural establishments): To provide early season protection of seedlings against wireworms, apply using HCBT or a batch treater at 12.8 fluid ounces per hundredweight of seed.