



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

EPA Reg. Number:

1381-258

Date of Issuance:

03/31/2016

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
 (under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Kochiavore

Name and Address of Registrant (include ZIP Code):

Diana Williams  
 Winfield Solutions, LLC  
 P.O. Box 64589  
 St. Paul, MN 55164-0589

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Kathryn V. Montague, Product Manager 23  
 Herbicides Branch, Registration Division (7505P)  
 Office of Pesticide Programs

Date:

03/31/2016

2. You are required to comply with the data requirements described in the DCI Order and/or Registration Review Workplans identified below:
  - a. 2-ethylhexyl ester of 2,4-D GDCI-030063-1362;
  - b. Bromoxynil Final Workplan, Registration Review Case #2070 issued 05/13/2014;
  - c. Fluroxypyr Preliminary Workplan, Registration Review Case #7248 issued 09/11/2014.

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 1381-258."
4. Submit one copy of the final printed label for the record before you release the product for shipment.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF and Alternate CSF 1 dated 10/13/2015.

If you have any questions, please contact Terri Stowe at (703) 305-6117 or by email at [stowe.terri@epa.gov](mailto:stowe.terri@epa.gov).

Enclosure



GROUP	<b>4 6</b>	HERBICIDE
-------	------------	-----------

# Kochiavore™

FOR SELECTIVE POSTEMERGENCE CONTROL OF BROADLEAF WEEDS  
 IN WHEAT, BARLEY, CORN, SORGHUM, NON-CROPLAND AREAS,  
 CONSERVATION RESERVE PROGRAM LAND AND FALLOWLAND

DO NOT APPLY TO RESIDENTIAL AREAS

**ACTIVE INGREDIENTS:**

2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid <sup>1</sup> .....	25.93%
Octanoic acid ester of bromoxynil (3,5-dibromo-4-hydroxybenzotrile) <sup>2</sup> .....	25.13%
Fluroxypyr 1-methylheptyl ester ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic acid, 1-methylheptyl ester <sup>3</sup> .....	10.00%

<b>OTHER INGREDIENTS:</b> .....	<u>38.94%</u>
<b>TOTAL</b> .....	<b>100.00%</b>

Contains petroleum distillates

Equivalent to: <sup>1</sup> 2,4-dichlorophenoxyacetic acid 17.26%, 1.67 lb/gal

<sup>2</sup> Bromoxynil 17.26%, 1.67 lb/gal

<sup>3</sup> Fluroxypyr acid 6.94%, 0.67 lb/gal

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

FIRST AID	
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of health emergency, call toll-free 1-877-424-7452.	
<b>NOTE TO PHYSICIAN:</b> May pose an aspiration pneumonia hazard. Contains petroleum distillate.	

See booklet for additional PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY  
 DISCLAIMER AND LIMITATION OF LIABILITY.

EPA Reg. No. 1381- \_\_\_\_\_

EPA Est. No. \_\_\_\_\_

Distributed By  
 Winfield Solutions, LLC  
 P.O. Box 64589, St. Paul, MN 55164-0589

NET CONTENTS \_\_\_\_\_ GALLONS

1/0324/6

**WINFIELD™**

## **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are listed below.

#### **All mixers, loaders, applicators, flaggers, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton
- Chemical-resistant apron when mixing or loading, cleaning equipment, cleaning up spills, or otherwise exposed to the concentrate
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

To reduce exposure to residues, wash the spray rig, tractor, and all other equipment used to handle or apply this product with water daily or before using the equipment for any other purpose.

**ENGINEERING CONTROLS:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE may be reduced or modified as specified in the WPS. Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)]

### **USER SAFETY RECOMMENDATIONS**

#### **Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### **PHYSICAL AND CHEMICAL HAZARDS**

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, or viton, shoes plus socks, and protective eyewear.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

### **USE INFORMATION**

Kochiavore is an emulsifiable concentrate that provides selective postemergence control of broadleaf weeds in wheat, barley, corn and sorghum and non-cropland areas, including Conservation Reserve Program land and fallowland.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

Kochiavore is a premix of two Group 4 and one Group 6 herbicides. Any weed population may contain or develop plants naturally resistant to Group 4 and/or Group 6 herbicides. Weed species with acquired resistance to Group 4 and/or Group 6 herbicides may eventually dominate the weed population if Group 4 and/or Group 6 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Kochiavore or other Group 4 and/or Group 6 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of Kochiavore or other target site of action Group 4 and/or Group 6 herbicides that have a similar target site of action, on the same weed species.
- Using tank-mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or Winfield Solutions, LLC representative for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

### **USE RESTRICTIONS**

- For Crop Uses: Do NOT apply more than 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) of this product per acre per growing season
- For Non-Cropland Uses: Unless otherwise noted for a specific non-cropland use, Do NOT apply more than 2.5 pints (0.52 lb. AE 2,4-D, 0.52 lb. AE bromoxynil and 0.21 lb. AE fluroxypyr) of this product per acre per growing season.
- When applying this product, do not contaminate water used for domestic purposes or irrigation ditches.
- Do not allow spray drift to come in contact with or apply this product directly to susceptible broadleaf plants or broadleaf crops, including but not limited to the following: alfalfa, canola, cotton, edible beans,

grapes, lentils, lettuce, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco or tomatoes.

- Do not apply this product through any type of irrigation system (i.e., chemigation).
- A 30 day Plant Back Interval (PBI) is required for all crops listed on this label or Federally approved supplemental labeling. A 120 day PBI is required for all other crops not listed on the label.
- Aerial application is prohibited within 300 feet of residential areas (e.g., homes, schools, playgrounds, shopping areas, hospitals, etc.).
- Do not apply with backpack or hand-held application equipment.
- Do not apply to playgrounds, schoolyards, or other residential turfgrass areas, including golf courses, putting greens, or tees.
- Do not use this product on sod farms.

### **SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

#### **Droplet Size**

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASABE S572.1) or a volume median diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASABE S572.1) or a volume median diameter of 300 microns or greater for spinning atomizer nozzles.

#### **Wind Speed**

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

#### **Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if:

- a) conditions of temperature inversion exist, or
- b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### **Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

#### **Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### **Equipment**

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

### **Additional requirements for aerial applications:**

**Restriction: In non-cropland areas (including rights-of-way), this product may be applied aerially only by helicopter.** Do NOT apply this product to non-cropland areas using fixed-wing aircraft.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

### **Additional requirements for ground boom application:**

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### **MIXING INSTRUCTIONS**

This product may be tank mixed with other products at specified rates as long as tank mixing is not prohibited by the label(s) of the tank mix partner products, or as otherwise noted within the specific crop use directions for this product, and the tank mix partner products are labeled for the timing and method of application for the use site to be treated. If compatibility with another product is not known, perform a (jar) test to determine compatibility (see below).

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (see below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

### **Tank Mixing Precautions**

- Follow all applicable directions, restrictions and precautions that appear on the respective tank mix product labels.
- Do not exceed specified application rates. Do NOT tank mix with other pesticide products that contain the same active ingredient as this product unless the label of either mix partner specifies the maximum dosages that may be applied.
- Prior to final use, perform a (jar) test to verify the compatibility of tank mix partner products (see below).

### **Tank Mix Compatibility (Jar) Test**

The following jar test is recommended prior to tank mixing to ensure the compatibility of this product with other tank mix partner products:

- 1) Mix the desired tank mix ingredients in their relative proportions in a clear glass quart jar with lid.
- 2) Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour.
- 3) If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combinations should not be used.

### **Tank Mixing Instructions**

Note: When adding ingredients to the mixture, allow time for each ingredient to be thoroughly mixed before adding the next.

1. Fill spray tank with water equal to 1/4 to 1/3 of the required spray volume and start agitation.
2. Maintain agitation continuously during mixing, final filling and while applying.
3. Add different formulation types in the following order, being certain to allow sufficient time for each product to completely mix and disperse after addition. Kochiavore is an emulsifiable concentrate (EC) formulation.
  - a) dry flowables
  - b) wettable powders
  - c) aqueous suspensions
  - d) flowables or liquids

4. Maintain agitation and fill spray tank to 3/4 of total spray volume and then add this product, other emulsifiable concentrates and any solutions.
5. Finish filling the spray tank.

While spraying, the tank mix ingredients may settle out of suspension if agitation is stopped before the spray tank is empty. The settled materials must be resuspended before any spraying is resumed. Note that settled material may be more difficult to resuspend than when originally mixed.

### **APPLICATION TIMING**

Only weeds that have emerged at the time of application will be controlled, therefore, be certain to apply to actively growing weeds. Weed control may be reduced and the risk of crop injury (at all stages of growth) may increase if extreme growing conditions (such as drought or near-freezing temperatures) occur prior to, at, or following application. Control may be decreased if target plant foliage is wet at the time of application. Applications of this product are rainfast within 1 hour after application.

### **Effect of Temperature on Herbicidal Activity**

The herbicidal activity of this product is influenced by weather conditions. Optimum herbicidal activity requires active plant growth and temperatures between 55°F to 75°F. Reduced efficacy will occur when temperatures are below 45°F or above 85°F. Weed control and crop tolerance may be reduced if frost occurs before or shortly after application (3 days).

### **SPRAY COVERAGE**

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. For best results (and to minimize spray drift), apply in a spray volume of 8 gallons or more per acre by ground and 3 or more gallons of total spray volume per acre by air. Spray volume should be increased as weed density and vegetative canopy increase in order to obtain equivalent weed control; however, do not exceed 40 gallons per acre total spray volume. Rather than increasing boom pressure, decrease spraying speed or use larger nozzle tips to increase spray volume.

Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, be certain to follow the precautions under the heading, "Spray Drift Management".

### **Adjuvants**

Unless otherwise noted within the specific crop and non-cropland use directions, a high-quality adjuvant may be used to improve weed control. An adjuvant can optimize herbicidal activity when applications are made at lower carrier volumes, under conditions of cool temperature, low relative humidity or drought, or to small, heavily pubescent kochia. When an adjuvant is to be used with this product, Winfield Solutions, LLC recommends the use of a Council of Producers & Distributors of Agrotechnology (CPDA) certified adjuvant.

### **Spot Treatments**

Only apply using a calibrated boom sprayer using the directions below. Application rates in the table below are based on an area of 1,000 square feet.

Do not apply with backpack or hand-held application equipment.

Mix the amount of this product (fluid ounces or ml) corresponding to the desired broadcast rate in one or more gallons of spray. To calculate the amount of this product required for larger areas, multiply the table value (fluid ounces or ml) by the area to be treated in "thousands" of square feet. An area of 1,000 square feet is approximately 10.5 X 10.5 yards (strides) in size.

For example: If the area to be treated is 3,500 square feet, multiply the table value by 3.5 (calc.  $3,500 \div 1,000 = 3.5$ ).



<b>Broadcast Rate Conversion Table for Spot Treatments</b>	
Broadcast Rate (Pints per Acre)	Kochiavore per Gallon (Fluid Ounces (ml))
1 (0.21 lb. AE 2,4-D; 0.21 lb. AE bromoxynil; 0.08 AE fluroxypyr)	0.37 (10.9)
1-1/2 (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr)	0.55 (16.3)
2-1/2 (Non-Cropland Use Only) (0.52 lb. AE 2,4-D, 0.52 lb. AE bromoxynil and 0.21 lb. AE fluroxypyr)	0.92 (27.2)

### APPLICATION RATES

In general, the application rates at the lower end of the specified rate range will be efficacious when applied to susceptible weed species with young, succulent growth. Use the higher rates within the rate range when applying to less sensitive species, perennials, and under conditions where control is more difficult (e.g., when plants are stressed due to drought or extreme temperatures, in dense weed stands and/or the weeds are larger). The higher rates will also be needed to control or suppress weeds in areas where competition from crops is not present (e.g., fallowland).

### SPRAYER CLEANUP

To avoid injury to desirable plants, before applying other chemicals with the equipment used to apply this product, all equipment must be thoroughly cleaned.

1. After applying this product, flush and rinse application equipment with water thoroughly, disposing of the water according to the disposal instructions in this label. All rinse water must be disposed of in compliance with local, state and federal guidelines.
2. Hose down the interior surfaces of the tank, flushing the tank, hoses, boom and nozzles with clean water for 10 minutes.
3. Fill the tank with water and recirculate for 15 minutes.
4. Spray part of the mixture through the hoses, boom and nozzles and drain the tank.
5. Remove the nozzles and screens and clean separately.
6. If the spray equipment will be used on crops other than those labeled for this product, repeat steps 1 and 2 and thoroughly wash the outside of spray tank and the boom.

### APPLICATION INFORMATION

<b>Broadleaf Weeds Controlled or Suppressed</b>		
Arrowhead	Goatsbeard	Nightshade spp <sup>†</sup> .
Bedstraw (cleavers)	Gumweed	Pennycress, field
Bindweed, field <sup>†</sup>	Hemp dogbane	Pigweed spp.
Bindweed, hedge	Horsetail, field <sup>†</sup>	Plantain
Buckwheat, wild	Horseweed (marestail)	Potato, volunteer <sup>†</sup>
Canola, volunteer	Jimsonweed	Prickly lettuce
Chickweed	Knotweed	Prickly pear cactus
Clover, white	Kochia <sup>1</sup>	Puncturevine
Cocklebur	Lambsquarter	Purslane, common
Cockle, cow	Mallow, common <sup>†</sup>	Ragweed, common
Coffeeweed	Mallow, venice	Smartweed, annual
Devilsclaw	Marshelder	Sunflower
Fiddleneck	Morningglory	Thistle, Russian
Flax, volunteer	Mustard spp.	Velvetleaf
Frenchweed		

<sup>†</sup>Indicates Suppression Only - Suppression is a reduction in weed competition (reduction is population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

<sup>1</sup>Includes biotypes that are herbicide-resistant or tolerant. Weeds germinating after spraying will not be controlled.

### Management of Kochia Biotypes

Research indicates many biotypes of kochia may occur within a single field and while kochia biotypes can vary in their susceptibility to this product, in general all biotypes will be suppressed or controlled at the labeled rate of 1 to 1-1/2 pints per acre. A shift to more tolerant biotypes within a field may occur if this product is applied at rates lower than specified.

### Best Practices for Resistance Management

Extensive populations of dicamba -tolerant kochia have been identified in certain small grain and corn production regions (such as Chouteau, Fergus, Liberty, Toole, and Treasure counties in the state of Montana). For optimal control of dicamba-tolerant kochia in these counties, apply this product at the rate of 1-1/2 pints per acre.

To minimize selection pressure and preserve the utility of this product for control of dicamba-tolerant kochia biotypes, this product should be rotated with products that do not contain dicamba.

### CROP USES

CROP	KOCHIAVORE APPLICATION RATE	DIRECTIONS
<b>BARLEY WHEAT</b>	1 to 1-1/2 pints per acre	POSTEMERGENT APPLICATION: Apply from the 4-leaf (Zadoks 1 <sub>4</sub> ) to emergence of the flag leaf (Zadoks 3 <sub>7</sub> ). Use the higher rates when weeds are less susceptible or for heavy weed populations. To avoid injury, do not use with atrazine, crop oils or other adjuvants, such as non-ionic surfactants.
Restrictions and limitations for barley and wheat: <ul style="list-style-type: none"> <li>• Do NOT harvest treated forage or allow livestock to graze treated areas within 45 days of application.</li> <li>• Do NOT feed treated straw to livestock.</li> <li>• The risk of crop injury at all stages of growth and poor weed control is increased if the application is made and extreme growing conditions (such as drought or near freezing temperatures) occur prior to, at, and following the application. Reduced weed control may also occur during these conditions.</li> <li>• Do NOT apply when crop canopy covers the weeds as poor control will result.</li> <li>• Preharvest Interval (PHI): Do NOT apply within 40 days prior to harvesting grain and straw or within 14 days prior to cutting hay.</li> <li>• Make no more than one postemergence application per growing season at a maximum of 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) per acre per application, up to flag leaf emergence stage.</li> <li>• Do NOT spray plants beyond the flag leaf emergence stage.</li> <li>• Do NOT apply more than 1.5 pints of this product per acre per growing season.</li> </ul>		
<b>FIELD CORN</b>	1 to 1-1/2 pints per acre	PREPLANT APPLICATION: Apply prior to planting. Delay planting for a minimum of 7 days after application. Planting sooner may result in unacceptable crop injury. Do not perform tillage for at least 7 days after application. Do not use on sandy soils or unacceptable crop injury may result.
	1 to 1-1/2 pints per acre	PREEMERGENCE and REDUCED TILLAGE APPLICATION: Make applications after corn is planted but before emergence to control emerged broadleaf weeds. Use the higher rates when weeds are less susceptible or for heavy weed populations. The seed furrow must be completely closed at application or severe crop injury may result. Do not use on sandy soils or unacceptable crop injury may result.
	1 to 1-1/2 pints per acre	POSTEMERGENT APPLICATION: Apply to field corn as a broadcast or band treatment from the V3 to the V5 growth

		<p>stages. Application prior to the V3 growth stage may result in increased crop leaf burn. Apply with drop nozzles and directed spray only from the V6 growth stage. DO NOT apply from 2 weeks before tasseling to dough stage.</p> <p>To avoid crop injury, do not use with atrazine, crop oils or other adjuvants, such as non-ionic surfactants. Application during high moisture and temperature conditions may cause crop injury or brittleness. DO NOT cultivate for a week to 10 days after treatment or stalk breakage may occur. Do not use on sandy soils or unacceptable crop injury may result.</p>
<p>Restrictions and limitations for field corn:</p> <ul style="list-style-type: none"> <li>• Do not cut treated crop for feed or fodder or graze for 47 for days following application.</li> <li>• Do not harvest within 90 days of application.</li> <li>• Seed corn producers should consult the respective seed corn company regarding tolerance of certain seed production inbred lines to bromoxynil octanoate.</li> <li>• Maximum application of 3.0 pints (0.62 lb. AE 2,4-D; 0.62 lb. AE bromoxynil; 0.26 lb. AE fluroxypyr) per acre per growing season.</li> </ul> <p>Preplant or preemergence:</p> <ul style="list-style-type: none"> <li>• Limited to one preplant or preemergence application per growing season.</li> <li>• Maximum rate of application: 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) per acre.</li> </ul> <p>Postemergence:</p> <ul style="list-style-type: none"> <li>• Limited to one postemergence application per growing season.</li> <li>• Maximum rate of application: 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) per acre.</li> </ul>		
<p><b>SORGHUM, GRAIN AND FORAGE</b></p>	<p>1 to 1-1/2 pints per acre</p>	<p>Apply to sorghum from the V4 but prior to the pre-boot stage. Use drop nozzles and directed spray only from the V8 to the pre-boot stage. To reduce the potential for crop injury, drop nozzles should direct the spray toward the soil surface, to avoid spray contact with grain sorghum foliage. Do not apply during the boot, flowering or early dough stages.</p> <p>The higher rate may be needed for some weeds, such as perennials or for heavy weed populations, but chances of crop injury may increase. To avoid injury, do not use with atrazine, crop oils or other adjuvants, such as non-ionic surfactants.</p> <p>Some sorghum varieties and hybrids are 2,4-D sensitive. Crop injury may also be increased by high moisture and temperature conditions. Contact your seed company or Extension Service for advice.</p>
<p>Restrictions and limitations for sorghum:</p> <ul style="list-style-type: none"> <li>• Do not harvest within 70 days of application.</li> <li>• Do not permit meat or dairy animals to consume treated crop as fodder or forage or graze for 45 for days following application.</li> <li>• Limited to 1 application per growing season.</li> <li>• Maximum of 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) per acre per application.</li> </ul>		

**NON CROP USES  
(Non-Residential Sites Only)**

NON-CROP AREA	KOCHIAVORE APPLICATION RATE	DIRECTIONS
<b>RIGHTS-OF-WAY, ROADSIDES, INDUSTRIAL SITES, FENCE ROWS, NON-IRRIGATION DITCHBANKS, NON-RESIDENTIAL TURFGRASS AREAS</b>	1-1/4 to 2-1/2 Pints per Acre	<p>Mix with an adequate amount of water to thoroughly saturate all weeds with spray mixture. This may require a spray volume of 50 to 300 gallons of water per acre. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.</p> <p>Use a high surfactant crop oil concentrate such as Superb® HC, a high surfactant methylated seed oil such as Destiny® HC or another crop oil for improved performance.</p>
<p>Restrictions and limitations:</p> <ul style="list-style-type: none"> <li>• Maximum of 2.5 pints (0.52 lb. AE 2,4-D, 0.52 lb. AE bromoxynil and 0.21 lb. AE fluroxypyr) of product per acre per year.</li> <li>• Limited to 1 application per year.</li> <li>• Do not allow livestock to graze in treated areas or feed treated plant material to livestock.</li> <li>• Do not use this product on sod farms</li> <li>• Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.</li> </ul>		
<b>CONSERVATION RESERVE PROGRAM (CRP) LAND</b>	1-1/4 to 2-1/2 Pints per Acre	<p>This product may be used for Conservation Reserve Programs (CRP), weed and brush control or in State Recognized Noxious Weed area (noncropland areas). Refer to "Weeds Controlled" section for list of susceptible species. <b>For areas underseeded with alfalfa, do not apply more than 1.5 pints per acre per year.</b></p> <p>Some weed species require tank mixes for adequate control. Use a high surfactant crop oil concentrate such as Superb HC, a high surfactant methylated seed oil such as Destiny HC or another crop oil for improved performance.</p>
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> <li>• Maximum of 2.5 pints (0.52 lb. AE 2,4-D, 0.52 lb. AE bromoxynil and 0.21 lb. AE fluroxypyr) of product per acre per year. Limited to 1 application per year.</li> <li>• Do not apply more than 1.5 pints (0.31 lb. AE 2,4-D; 0.31 lb. AE bromoxynil; 0.13 lb. AE fluroxypyr) of product per acre per year to CRP areas that are underseeded with alfalfa.</li> <li>• Do not add spray adjuvants or fluid fertilizers when applying to CRP areas planted with alfalfa or other legumes.</li> <li>• Consult the Conservation Reserve Program rules for CRP uses. The more restrictive requirements of the program rules or this label must be followed.</li> <li>• Do not allow livestock to graze in treated areas or feed treated plant material to livestock.</li> </ul>		

<b>FALLOWLAND</b>	1-1/4 to 2-1/2 Pints per Acre	<p>Unless otherwise specified below, apply when broadleaf weeds are up to the 8-leaf stage, 4 inches in height or 2 inches in diameter, whichever comes first. Apply to grasses from the 3-leaf stage. Use the higher application rate for less susceptible weeds.</p> <p>Use a high surfactant crop oil concentrate such as Superb HC, a high surfactant methylated seed oil such as Destiny HC or another crop oil for improved performance.</p> <p>Kochia seedlings less than 4 inches tall (including ALS resistant biotypes) will be controlled using the 1.25 pint per acre rate. However, when conditions for control are less favorable, such as under drought or cool temperature, a rate of 1.5 pints per acre will provide more consistent control of kochia seedlings 1 to 4 inches tall. For more consistent control of small kochia, apply when the plants are at least 1 inch tall. A rate of 1.5 pints per acre should be used for optimal control of dicamba tolerant kochia populations (refer to the "Management of Kochia Biotypes" in the "Application Information" section above).</p>
<p>Restrictions and Limitations:</p> <ul style="list-style-type: none"> <li>• Do not allow livestock to graze in treated areas or feed treated grass to livestock.</li> <li>• Limited to two applications per year.</li> <li>• Minimum of 30 days between applications.</li> <li>• Do not apply more than 2-1/2 pints (0.52 lb. AE 2,4-D, 0.52 lb. AE bromoxynil and 0.21 lb. AE fluroxypyr) of this product per acre per application.</li> </ul>		

## STORAGE AND DISPOSAL

### Prohibitions

Do not contaminate water, food, or feed by storage and disposal.

### Pesticide Handling

Store in original containers only. Keep container closed when not in use. Do not store near food or feed.

In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent.

Remove to chemical waste area. **DO NOT ALLOW PRODUCT TO FREEZE.**

### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Use label language appropriate for container size and type.

**Nonrefillable containers.** Do not reuse or refill this container. Clean container promptly after emptying.

**Nonrefillable container equal to or less than 5 gallons.** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Nonrefillable container greater than 5 gallons.** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures.

Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call  
CHEMTREC 1-800-424-9300.**

#### **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Kochiavore is a trademark of Winfield Solutions, LLC.

Destiny and Superb are registered trademarks of Winfield Solutions, LLC.