



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

1381-267

Date of Issuance:

12/7/18

NOTICE OF PESTICIDE:

☒ Registration  
☐ Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Tetraban Fungicide

Name and Address of Registrant (include ZIP Code):

Lisa Mathias, Registration Specialist  
Land O' Lakes, LLC  
1080 Country Road F West, MS 5705  
Shoreview, MN 55126-2910

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch  
Registration Division 7505P

Date:

12/7/18

2. Make the following label changes before you release the product for shipment:

- Revise the EPA Registration Number to read, "EPA Reg. No. 1381-267."

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 07/16/2018

If you have any questions, please contact Eleanor Thornton by phone at 703-305-6799, or via email at [thornton.eleanor@epa.gov](mailto:thornton.eleanor@epa.gov).

Enclosure

# Tetraban® Fungicide

[Alternate Brand Names: Tetraban® 2F, Tetraban® 2 SC,  
Tetraban® 2.0 F, Tetraban® 2 F]

Broad spectrum fungicide for control of plant diseases

## Active Ingredient:

Azoxystrobin: methyl (*E*)-2-{2-[6-(2-cyanophenoxy)  
pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate\* .....22.9%

**Other Ingredients:** .....77.1%

**Total:** .....100.0%

Contains 2.08 lb of active ingredient per gallon

\*IUPAC

Reformulation is prohibited. See individual container labels for repackaging limitations.

## KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. Call 1-877-424-7452 for emergency medical treatment information.	

See booklet for additional **PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER, AND LIMITATION OF LIABILITY.**

**EPA Reg. No.** 1381-XXX

**EPA Est.** \_\_\_\_\_

## Manufactured for:

Winfield Solutions, LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589

**Net Contents** \_\_\_\_\_

1/1206/8

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

**IMPORTANT:** When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

### User Safety Recommendations

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer.

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash-water or rinsate.

### Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

## Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Winfield Solutions, LLC immediately if you observe any adverse environmental effects due to use of this product.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**Use of this product through airblast application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard and Springfield.**

This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.

**FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber or butyl rubber
- Shoes plus socks

---

## PRODUCT USE PRECAUTIONS

---

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.**

### ATTENTION

This product is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray this product where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

---

## PRODUCT INFORMATION

---

This product is a broad spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. This product may also improve the yield and/or quality of the crop. These additional benefits are due to positive effects on plant physiology. The effects may vary according to factors such as the crop, crop hybrid, or environment. This product may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

### Restrictions for Resistance Management Purposes

**Greenhouse Use:** To help manage fungicide resistance, do not use for commercial transplant production in the greenhouse except where specified on the label.

---

## PRODUCT USE INSTRUCTIONS

---

**Application:** Thorough coverage is necessary to provide good disease control. Make no more spray solution than is needed for application. Avoid spray overlap, as crop injury may occur.

**Adjuvants:** When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Council of Producers & Distributors of Agrotechnology (CPDA) adjuvant certification is recommended.

**Efficacy:** Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of this product has been used. If resistant isolates to Group 11 fungicides are present, efficacy can be reduced for certain diseases. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, with highly susceptible varieties, or when environmental conditions are conducive to disease.

**Crop Tolerance:** Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank-mix combinations have been tested under all conditions. Test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur as a result of application. See Product Use Precautions for apple phytotoxicity information.

## INTEGRATED PEST (DISEASE) MANAGEMENT

Integrate this product into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development. Cultural practices include selection of varieties with disease tolerance, removal of plant debris in which inoculum overwinters, and proper timing and placement of irrigation. Consult your local agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

## RESISTANCE MANAGEMENT

For resistance management, Tetraban Fungicide contains a Group 11 fungicide (azoxystrobin). The mode of action for Tetraban Fungicide is the inhibition of the QoI (quinone outside) site within the electron transport system. Any fungal population may contain individuals naturally resistant to Tetraban Fungicide and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Because resistance development cannot be predicted, use resistance management strategies established for the crop and use area. Consult your local or State agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include alternating and/or tank-mixing with products having different modes of action or limiting the total number of applications per year. Winfield Solutions, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

Follow the crop specific resistance management instructions in the directions for use.

If no resistance instructions on number of applications is specified in the directions for use, follow the guidelines in the table below.

<b>If planned total number of fungicide applications per crop is:</b>	1	2	3	4	5	6	7	8	9	10	11	12
<b>Solo QoI fungicide sprays</b>	1	1	2	2	2	2	2	3	3	3	3	4
<b>QoI fungicide sprays in mixture (tank-mix or formulated)</b>	1	2	2	2	2	3	3	4	4	5	5	6

In situations requiring multiple sprays, develop spray programs for Group 11 (QoI) fungicides. In crops where two sequential Group 11 fungicide applications are made, alternate the applications with two or more applications of a fungicide that is not in Group 11. If more than 12 applications are made, observe the following guidelines:

- When using a QoI fungicide as a solo product, the number of applications must be no more than 1/3 (33%) of the total number of fungicide applications per year.
- For QoI mixes in programs in which tank mixes or premixes of QoI with mixing partners of a different mode of action are utilized, the number of QoI containing applications must be no more than 1/2 (50%) of the total number of fungicide applications per year.
- In programs in which applications of QoI are made with both solo products and mixtures, the number of QoI containing applications must be no more than 1/2 (50%) of the total number of fungicide applications per year.

If a Group 11 fungicide is applied to the seed or soil, do not make another application with a Group 11 fungicide for at least 3 weeks.

## ROTATIONAL CROP RESTRICTIONS

The following crops may be planted at the specified interval following application of this product.

### Crop Rotational Interval

	Plant back interval
Buckwheat and millet	12 months
All other crops with Azoxystrobin registered uses	0 days

## SOILBORNE/SEEDLING DISEASE CONTROL

For those crops that have specific use directions for soilborne disease control: This product can provide control of many soilborne diseases if applied early in the growing season. Specific applications for soilborne diseases include in-furrow applications and banded applications applied over the row, either shortly after plant emergence or during herbicide applications or cultivation. These applications will provide control of pre- or postemergence damping off and diseases that infect plants at the soil-plant interface.

The use of either type of application depends on the cultural practices in the region. In some locations, one type of application may provide better disease control than the other, depending on the timing of the disease epidemic. Seedling diseases are generally controlled by in-furrow applications while banded applications are more effective against soilborne diseases that develop later in the season. Consult your local expert to get some guidance regarding application type.

Under cool, wet conditions, crop injury from soil directed applications can occur.

### BANDED

- Apply this product prior to infection as a directed spray to the soil, using single or multiple nozzles, adjusted to provide thorough coverage of the lower stems and the soil surface surrounding the plants.
- Limit the band width to 7 inches or less.
- Apply this product at a rate of 0.40-0.80 fl oz product (0.10-0.20 oz ai)/1000 row feet. For banded applications on 22-inch rows, the maximum application rate is 0.70 fl oz/1000 row feet.
- These applications come into contact with the foliage and are counted as foliar applications when considering resistance management.
- They may be applied during cultivation or hilling operations to provide soil incorporation.

### IN-FURROW

- Apply this product as an in-furrow spray in 3-15 gallons of water at planting.
- Mount the spray nozzle so the spray is directed into the furrow just before the seeds are covered.
- Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place.

## IN-FURROW APPLICATION RATES

Rate per 1000 row-feet		Row Spacing (inches)										
fl oz product	oz ai	22	30	32	34	36	38	40	48	60	72	80
		Product per Acre (fl oz)										
0.40	0.10	9.5	7.0	6.5	6.1	5.8	5.5	5.2	4.4	3.5	2.9	2.6
0.60	0.15	14.3	10.5	9.8	9.2	8.7	8.3	7.8	6.5	5.2	4.4	3.9
0.80	0.20		13.9	13.1	12.3	11.6	11.0	10.5	8.7	7.0	5.8	5.2
1.00	0.25					14.5	13.8	13.1	10.9	8.7	7.3	6.5
1.20	0.30								13.1	10.5	8.7	7.8
1.38	0.36								15.0	12.0	10.0	9.0
1.50	0.40									13.1	10.9	9.8
1.72	0.45									15.0	12.5	11.2
2.00	0.50										14.5	13.1
2.07	0.54										15.0	13.5
2.30	0.60											15.0

Do not apply more than 15 fl oz/A.

Row spacing (inches)	Row-Feet Per Acre
22	23,760
30	17,424
32	16,335
34	15,374
36	14,520
38	13,756
40	13,068
48	10,890
60	8,712
72	7,260
80	6,534

## DRIP

Refer to the **Application Instructions Through Irrigation System** section.

---

## SPRAY DRIFT MANAGEMENT

---

To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather-related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

### ATTENTION

This product is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray this product where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.

---

## MIXING AND APPLICATION METHODS

---

**Spray Equipment:** This product may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

### Nozzles

- Equip sprayers with nozzles that provide accurate and uniform application.
- Ensure that nozzles are the same size and uniformly spaced across the boom.
- Calibrate sprayer before use.
- It is suggested that screens be used to protect the pump and to prevent nozzles from clogging.
- Use 16-mesh or coarser screens when they are placed on the suction side of the pump.
- Do not place a screen in the recirculation line.
- Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles.
- Check nozzle manufacturer's specifications.

### Pump

- Use a pump with capacity to:
  - (1) Maintain 35-40 psi at nozzles.
  - (2) Provide sufficient agitation in tank to keep mixture in suspension - this requires recirculation of 10% of tank volume per minute.
- Use a jet agitator or liquid sparge tube for agitation.
- Do not air sparge.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural recommendations.

### **Mixing Instructions**

- This product is a suspension concentrate (SC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

### **Tetraban Fungicide Alone (No Tank Mix)**

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add Tetraban Fungicide to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Tetraban Fungicide has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

**Tetraban Fungicide + Tank Mixtures:** This product is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

This product has demonstrated some phytotoxic effects when mixed with products that are formulated as emulsifiable concentrates (EC). These effects are enhanced if applications are made under cool, cloudy conditions and these conditions remain for several days following application. In addition, adjuvants that contain some form of silicone have also contributed to phytotoxicity.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. In particular, no dosage rate listed in any label may be exceeded and the most restrictive label precautions and limitations must be followed. Do not use any product which prohibits mixing with this product.

### **Mixing in the Spray Tank**

- Add 1/2 - 2/3 of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Tetraban Fungicide to the spray tank.
- Allow Tetraban Fungicide to completely disperse.
- Spray the mixture with the agitator running.

---

## APPLICATION INSTRUCTIONS THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

---

### Application Through Irrigation Systems (Chemigation)

- Use only on crops for which chemigation is specified on this label.
- Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
- Apply in 0.1-0.25 inches/acre. Excessive water may reduce efficacy.
- If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Spray Preparation:** Thoroughly clean chemical tank and injector system. Flush system with clean water.

**Drip Irrigation:** This product may be applied through drip irrigation systems for soilborne disease control. Ensure that the soil has adequate moisture capacity prior to drip application.

Terminate drip irrigation at fungicide depletion from the main feed supply tank or after 6 hours from start, whichever is shorter. For maximum efficacy, delay subsequent irrigation (water only) for at least 24 hours following drip application.

### Sprinkler Irrigation

- Apply this product through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems.
- Do not apply this product through any other type of irrigation system except as specified on this label.
- Apply with center pivot or continuous-move equipment distributing  $\frac{1}{2}$  acre-inch or less during treatment.
- In general, use the least amount of water required for proper distribution and coverage.
- If stationary systems (solid set, handlines or wheel lines other than continuous-move) are used, inject this product into no more than the last 20-30 minutes of the set.
- Do not apply when winds are greater than 10-15 mph to avoid drift or wind skips.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- Plant injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform treated water.
- Thorough coverage of foliage is required for good control.
- Maintain thorough agitation during the entire application period.

If you have questions about calibration contact State Extension Service specialist, equipment manufacturers or other experts.

## **Operating Instructions**

1. Do not apply when wind speed favors drift beyond the area intended for treatment.
2. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
3. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
6. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
7. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
8. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
9. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

## **Center Pivot Irrigation Equipment**

- Determine the size of the area to be treated.
- Determine the time required to apply 1/8 - 1/2 inch of water over the area to be treated when the system and injection equipment are operated at normal pressures as specified by the equipment manufacturer. When applying this product through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution. Run the system at 80-95% of the manufacturer's rated capacity.
- Using water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of this product required to treat the area covered by the irrigation system.
- Add the required amount of this product and sufficient water to meet the injection time requirements to the solution tank.
- Make sure the system is fully charged with water before starting injection of the Tetraban Fungicide solution. Time the injection to last at least as long as it takes to bring the system to full pressure.
- Maintain constant solution tank agitation during the injection period.
- Continue to operate the system until the Tetraban Fungicide solution has cleared the sprinkler head.

**Notes:** (1) Use only with drive systems which provide uniform water distribution. (2) Do not use end guns when chemigating this product through center pivot systems because of non-uniform application.

## **Solid Set, Hand Move, and Moving Wheel Irrigation Equipment**

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying this product through irrigation equipment use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of this product required to treat the area covered by the irrigation system.
- Add the required amount of this product into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Tetraban Fungicide solution has cleared the last sprinkler head.

### **Specific Instructions for Public Water Systems**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## Directions For Use

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Alfalfa</b> (See Nongrass Animal Feeds Forage, Fod- der, Straw and Hay)			
<b>Almonds</b>	Alternaria Leaf and Fruit Spot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichum acutatum</i> ) Leaf Blight ( <i>Seimatosporium lichenicola</i> ) Leaf Rust ( <i>Tranzschelia discolor</i> ) Scab ( <i>Cladosporium carpopophilum</i> ) Shot Hole ( <i>Wilsonomyces carpophilus</i> )	6.0-15.5 (0.10-0.25)	<p>Begin applications prior to disease development and continue following the resistance management guidelines. Applications may be made by ground, air or chemigation. For aerial applications apply in a minimum of 15 GPA. Thorough and uniform coverage is essential for disease control. Reduced efficacy has been observed when uniform coverage cannot be obtained.</p> <p>This product may be applied by air only at growth stages prior to and including 5 weeks after petal fall. An adjuvant may be added at specified rates.</p> <p>Anthracnose, scab and shot hole: Begin applications prior to disease development and continue at 7- to 14-day intervals.</p> <p>Blossom blight: Begin applications at early bloom and continue through petal fall.</p>
	Brown Rot Blossom Blight ( <i>Monilinia laxa</i> , <i>M. fructicola</i> )	12.0-15.5 (0.20-0.25)	Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

### Specific Use Restrictions:

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 28 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Artichoke, Globe</b>	Ramularia Leaf Spot ( <i>Ramularia cynarae</i> )	11.0-15.5 (0.18-0.25)	<p>Begin applications prior to or in the early stages of disease development and continue as needed at a 2-3 week interval, up to and including the day of harvest. Do not apply at less than 7-day intervals. Applications may be made by ground, air or chemigation. For ground applications, apply in 50-200 gallons of water per acre to obtain coverage without excessive runoff. For aerial applications, apply in a minimum of 5 gallons of water per acre. An adjuvant may be added at specified rates.</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 8 applications per year when applying at the lowest rate (11.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

<b>Asparagus</b>	Stemphyllium Purple Spot ( <i>Stemphyllium vesicarium</i> )	6.0-15.5 (0.10-0.25)	<p>Begin applications prior to disease development and continue on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre by ground, and minimum of 3 gallons per acre by air.</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
------------------	--	-------------------------	---

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 100 days

<b>Bananas Plantains</b>	Black Sigatoka ( <i>Mycosphaerella fijiensis</i> ) Yellow Sigatoka ( <i>Mycosphaerella musicola</i> )	5.5-8.5 (0.09-0.135)	<p>Begin applications prior to disease development and continue every 12-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
------------------------------	--	-------------------------	---

**Specific Use Restrictions:**

- 1) Do not apply more than 66.4 fl oz (1.08 lb ai) of this product/A/year.
- 2) Do not apply more than 1.08 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 7 applications of this product per year when applying at the highest rate (8.5 fl oz/A) or 12 applications per year when applying at the lowest rate (5.5 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Cereals  Barley Oats Rye	Kernel Blight or Black Point ( <i>Alternaria</i> spp.) ( <i>Cochliobolus sativus</i> ) Leaf Rust ( <i>Puccinia hordei</i> ) ( <i>P. recondita</i> )	6.0-12.0 (0.10-0.20)	Apply this product prior to disease development. Protecting the flag leaf is important for maximizing disease control. For best results, sufficient water volume must be used to provide thorough coverage.  This product can be applied by ground, air or chemigation.
	Barley Stripe ( <i>Drechslera graminea</i> = <i>Pyrenophora graminea</i> ) Net Blotch ( <i>Pyrenophora teres</i> ) Scald ( <i>Rhynchosporium secalis</i> ) Septoria Leaf and Glume Blotch ( <i>Septoria</i> spp., <i>Stagonospora</i> spp.) Spot Blotch ( <i>Cochliobolus sativus</i> ) Stem Rust ( <i>Puccinia graminis f.sp.</i> <i>tritici</i> ) Stripe Rust ( <i>Puccinia striiformis</i> ) Tan Spot ( <i>Pyrenophora trichostroma</i> )	9.0-12.0 (0.15-0.20)	A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.  For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Powdery Mildew ( <i>Erysiphe graminis f. sp.</i> <i>hordei</i> ) Stagonospora Blotch ( <i>Stagonospora nodorum</i> )	12.0 (0.20)	

- 1) **Specific Use Restrictions:** Do not apply more than 24.6 fl oz (0.40 lb ai) of this product/A/year.
- 2) Do not apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.
- 3) Do exceed 2 applications of this product or other Group 11 fungicide per year.
- 4) Do not apply after Feekes 10.54.
- 5) Pre-Harvest Interval (PHI): 7 days (forage and hay)
- 6) Do not apply within 7 days of grazing forage and hay.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Berries, Bushberry Subgroup 13-07B</b>  Aronia Berry Blueberry, Highbush Blueberry, Lowbush Buffalo Currant Chilean Guava Cranberry, Highbush Currant, Black Currant, Red Elderberry European Barberry Gooseberry Honeysuckle Edible Huckleberry Jostaberry Juneberry (Saskatoon Berry) Lingonberry Native Currant Salal Sea Buckthorn  Including all culti- vars and/or hybrids of these	Alternaria Fruit Rot ( <i>Alternaria</i> spp.) Anthracnose Fruit Rot ( <i>Colletotrichum</i> <i>gloeosporoides</i> ) Botryosphaeria Canker ( <i>Botryosphaeria</i> spp.) Leaf Spot and Blotch ( <i>Mycosphaerella</i> spp., <i>Septoria</i> spp.) Mummyberry ( <i>Monilinia</i> <i>vacciniicorymbosi</i> ) Phomopsis Leaf Spot, Twig Blight and Stem Canker ( <i>Phomopsis vaccinii</i> ) Powdery Mildew ( <i>Sphaerotheca</i> spp.) Septoria Blight ( <i>Septoria</i> spp.) Spur Blight ( <i>Didymella</i> spp., <i>Phoma</i> spp.)	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**Specific Use Restrictions:**

- 1) Do not apply more than 46 fl oz (0.75 lb ai) of this product/A/year.
- 2) Do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 7 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Berries, Caneberry Subgroup 13-07A</b>  Blackberry Bingleberry Boysenberry Dewberry Lowberry Marionberry Olallieberry Youngberry Loganberry Red and Black Raspberry Wild Raspberry  Including all culti- vars and/or hybrids of these	Anthracnose ( <i>Spaceloma necator</i> ) ( <i>Elsinoe veneta</i> ) Botryosphaeria Canker ( <i>Botryosphaeria dothidea</i> ) Colletotrichum Rot ( <i>Colletotrichum gloeosporioides</i> ) Leaf Spot and Blotch ( <i>Mycosphaerella</i> spp.) ( <i>Septoria rubi</i> ) ( <i>Sphaerulina rubi</i> ) Powdery Mildew ( <i>Sphaerotheca macularis</i> ) ( <i>Microphaera</i> spp.) ( <i>Oidium</i> spp.) Rosette or Double Blossom of Blackberries ( <i>Cercospora rubi</i> ) Spur Blight ( <i>Didymella applanata</i> )	6.0-15.5 (0.10-0.25)	Begin applications at onset of disease and continue as required until harvest. Make applications on a 7- to 14-day schedule. Use a minimum water volume of 10 gallons per acre by ground and a minimum of 3 gallons by air.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Blackberry Rust ( <i>Phragmidium</i> spp.)	10.0-15.5 (0.16-0.25)	

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Berries, Low Growing Subgroup 13-07G (except Cranberry)</b>  <b>Strawberry</b>  See additional crops below.	Anthracnose <i>(Colletotrichum fragariae)</i> Leather Rot <i>(Phytophthora cactorum)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i>  Suppression of Botrytis on the Foliage <i>(Botrytis cinerea)</i>	6.0-15.5 (0.10-0.25)	<p>Begin applications prior to disease development and continue on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>For leather rot control apply 2 applications on a 7-day schedule from late bloom through harvest.</p> <p><b>Field Nurseries:</b> Apply to young plants in field nurseries by ground, drip, or overhead chemigation.</p> <p>If applying through drip irrigation, calculate the rate as a band application with a band width equal to the root zone width. Inject this product into the irrigation water.</p> <p><b>For dip applications at transplanting for commercial berry production:</b>            For suppression of root and crown rot caused by <i>Colletotrichum</i> spp., mix 5-8 fl oz of this product per 100 gallons of water. Dip plants for 2-5 minutes. Plant treated plants as quickly as possible. Wash transplants to remove excess soil prior to dipping. For continued anthracnose control, follow with foliar applications beginning 2-3 weeks after transplant.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
	<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Additional Low Growing Berries:** Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry including all cultivars and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 61.5 fl oz (1.0 lb ai) of this product/A/year.
- 2) Do not apply more than 1.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 3 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 10 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Berries, Low Growing Subgroup 13-07H (except Strawberry)</b>  <b>Cranberry</b>  See additional crops below.	Cottonball <i>(Monilinia oxycocci)</i> Fruit Rots <i>(Phylospora vaccinii)</i> <i>(Glomerella cingulata)</i> <i>(Coleophoma empetri)</i> Lophodermium Twig Blight <i>(Lophodermium spp.)</i>	6.0-15.5 (0.10-0.25)	Begin applications at 5-10% bloom for fruit rot, cottonball, and twig blight. Continue applications on a 7- to 14-day schedule if conditions are favorable for disease development. Applications may be made by ground, chemigation or air.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Fairy Ring (suppression) <i>(Psilocybe spp.)</i>	15.5 (0.25)	Make the first application at bud break. Measure the ring diameter and add 10 feet to that diameter. Apply this product at a rate equivalent to 15.5 fl oz/A in 30 – 100 gallons of water to the affected area. Irrigation (1 – 2 hours) following application is advisable to ensure penetration to the base of the plant. If necessary, make another application 2 – 4 weeks later. For ground application ensure adequate water volume for thorough canopy penetration.

**Additional Low Growing Berries:** Bearberry; Bilberry; Blueberry, lowbush; Cloudberry; Lingonberry; Muntries; and Partridgeberry including all cultivars and/or hybrids of these

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Do not treat cranberry fields used for aquaculture of fish and crustacea.
- 5) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Use care in making applications near non-target aquatic habitats.
- 6) Do not apply to flooded crop.
- 7) Do not allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.
- 8) Pre-Harvest Interval (PHI): 3 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Brassica, Head and Stem Subgroup 5A</b>  Broccoli Chinese Broccoli (gai lon) Brussels Sprouts Cabbage Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) Cauliflower Cavalo Broccolo Kohlrabi  Including all culti- vars and/or hy- brids of these	Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Cercospora Leaf Spot ( <i>Cercospora</i> <i>brassicicola</i> ) Downy Mildew ( <i>Peronospora</i> <i>parasitica</i> ) Pin Rot ( <i>Alternaria</i> spp.) Powdery Mildew ( <i>Erysiphe polygoni</i> ) Rhizoctonia Blight ( <i>Rhizoctonia solani</i> ) Ring Spot ( <i>Mycosphaerella</i> <i>brassicicola</i> ) White Leaf Spot ( <i>Pseudocercospora</i> <i>capsellae</i> ) White Rust ( <i>Albugo candida</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a minimum of 10 gallons of water per acre by ground, and minimum of 3 gallons per acre by air.  Do not apply more than two applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0-day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Brassica, Leafy Greens Subgroup 5B</b>  Broccoli Raab Cabbage Chinese Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens  Including all cultivars and/or hybrids of these	Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum</i> spp.) Black Spot ( <i>Alternaria</i> spp.) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Downy Mildew ( <i>Peronospora</i> <i>parasitica</i> ) Powdery Mildew ( <i>Erysiphe polygoni</i> ) Ring Spot ( <i>Mycosphaerella</i> <i>brassicicola</i> ) White Rust ( <i>Albugo candida</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 46 fl oz (0.75 lb ai) of this product/A/year.
- 2) Do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 7 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Bulb Vegetables Crop Group 3-07</b>  Garlic Leek Onion, bulb Daylily, bulb Fritillaria, bulb Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Lily, bulb Onion, bulb Onion, Chinese, bulb Onion, pearl Onion, potato, bulb Shallot, bulb Onion, green Chive, fresh leaves Chive, Chinese, fresh leaves Elegans hosta Fritillaria, leaves Kurrat Lady's leek Leek Leek, wild Onion, beltsville bunching Onion, fresh Onion, green Onion, macrostem Onion, tree, tops Onion, Welsh, tops Shallot, fresh leaves  Including all cultivars and/or hybrids of these	<b>Foliar Diseases</b> Cladosporium Leaf Blotch ( <i>Cladosporium allii</i> ) Powdery Mildew ( <i>Leveillula taurica</i> ) Purple Blotch and Leaf Blight ( <i>Alternaria porri</i> ) ( <i>Stemphylium vesicarium</i> ) Rust ( <i>Puccinia allii</i> )	6.0-12.0 (0.10-0.20)	For downy mildew, make preventative applications on a 5- to 7-day schedule.  For all other diseases, begin applications prior to disease development and continue every 7-14 days, following the resistance management guidelines. Applications may be made by ground, air or chemigation. For aerial application, use the higher rates for adequate control. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Botrytis Leaf Blight ( <i>Botrytis aclada</i> ) Downy Mildew ( <i>Peronospora destructor</i> )	9.0-15.5 (0.15-0.25)	Mixtures of this product with insecticides and silicone adjuvants must be tested for crop safety before application to the crop.
	<b>Soilborne Diseases</b> Rhizoctonia Damping-Off ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.  For in-furrow application, spray just prior to seed placement so that the majority of the chemical is under the seed. This will reduce the potential for phytotoxicity, especially if fertilizer is added to the application.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Canola</b> (see Oilseed Crops for additional information)	Alternaria Blackspot ( <i>Alternaria</i> spp.) Blackleg ( <i>Leptosphaeria maculans</i> ) Sclerotinia Stem Rot ( <i>Sclerotinia sclerotiorum</i> )	6.0-15.5 (0.10-0.25)	<p>In general, apply 7.0 fl oz of this product at early bud followed by 14.0 fl oz at about 45 days before harvest. A third application of 7.0 fl oz may be made 30 days before harvest.</p> <p>Specifically, for blackleg, apply at the 2- to 4-leaf stage.</p> <p>For Alternaria or Sclerotinia, apply 9.0-15.5 fl oz product/A at 10-25% flowering (3-7 days following first flower). Use the higher rate under heavy disease pressure or when conditions are favorable for disease.</p> <p>For control of Alternaria alone, 8.0 fl oz product/A may be applied at pod stage (approximately 95% petal fall).</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternating with a fungicide that is not in Group 11.</p> <p>Applications may be made by ground, air or chemigation. Use a minimum of 10 gallons of water per acre for ground applications.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 27.7 fl oz (0.45 lb ai) of this product/A/year.
- 2) Do not apply more than 0.45 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 1 application of this product per year when applying at the highest rate (15.5 fl oz/A) or 4 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 30 days

<b>Carrots</b>	Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Early Blight ( <i>Cercospora carotae</i> ) Late Blight ( <i>Alternaria dauci</i> ) Powdery Mildew ( <i>Erysiphe</i> spp.) White Mold ( <i>Sclerotium rolfsii</i> )  For additional diseases, see <b>Vegetables, Root, Subgroup.</b>	9.0-20.0 (0.15-0.33)	<p>Begin applications prior to disease development and continue every 7-14 days, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
	<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 6 applications of this product per year when applying at the highest rate (20 fl oz/A) or 13 applications when applying at the lowest rate (9.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Celery	Early Blight ( <i>Cercospora apii</i> ) Late Blight ( <i>Septoria apicola</i> )  For additional diseases, see <b>Leafy Vegetables.</b>	9.0-15.5 (0.15-0.25)	Begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 10 applications when applying at the lowest rate (9.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Christmas Trees	Diplodia Tip Blight ( <i>Diplodia pinea</i> ) Lophodermium Needlecast ( <i>Lophodermium pinastri</i> ) Swiss Needlecast ( <i>Phaeocryptopus gaumannii</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue at 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
-----------------	---	-------------------------	---

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 7 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 20 applications per year when applying at the lowest rate (6.0 fl oz/A).

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Citrus Fruit Crop Group 10-10</b>  Calamondin Citron Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma Mandarin Tangerine  Including all cultivars and/or hybrids of these  See complete list of citrus fruit crops be- low.	Albinism ( <i>Alternaria alternata</i> pv <i>citri</i> ) Alternaria Leaf and Fruit Spot ( <i>Alternaria citri</i> ) Anthracnose ( <i>Colletotrichum acutatum</i> , <i>C. gloeosporioides</i> ) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Diplodia Stem-End Rot ( <i>Diplodia natalensis</i> ) Greasy Spot ( <i>Mycosphaerella citri</i> ) Melanose ( <i>Diaporthe citri</i> ) Penicillium Decays Green Mold, Whisker Mold, Suppression of Blue Mold ( <i>Penicillium</i> spp.) Phomopsis Stem-End Rot ( <i>Phomopsis citrii</i> ) Post Bloom Fruit Drop (PFD) ( <i>Colletotrichum acutatum</i> ) Powdery Mildew ( <i>Erysiphe</i> spp.) Scab ( <i>Elsinoe fawcettii</i> ) Sweet Orange Scab ( <i>Elsinoe australis</i> )	12.0-15.5 (0.20-0.25)	Begin applications prior to disease development and continue at 7- to 21-day intervals following the resistance management guidelines. Under conditions that favor severe disease epidemics, use the higher application rates. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. Use a horticultural spray oil to improve control of greasy spot.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Black Spot ( <i>Guidnardia citricarpa</i> )	9.0-15.5 (0.15-0.25)	
Pummelo Citrus Hybrid (Uniq fruit only)	<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Complete List of Citrus Fruit Crops:** Australian Desert Lime (*Eremocitrus glauca*); Australian Finger Lime (*Microcitrus australasica*); Australian Round Lime (*Microcitrus australis*); Brown River Finger Lime (*Microcitrus papuana*); Calamondin (*Citrofortunella microcarpa*); Citron (*Citrus medica*); Citrus Hybrids, *Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.; Grapefruit (*Citrus paradise*); Japanese Summer Grapefruit (*Citrus natsudaidai*); Kumquat (*Fortunella* spp.); Lemon (*Citrus limon*); Lime (*Citrus aurantiifolia*); Mediterranean Mandarin (*Citrus deliciosa*); Mount White Lime (*Microcitrus garrowayae*); New Guinea Wild Lime (*Microcitrus warburgiana*); Orange, Sour (*Citrus aurantium*); Orange, Sweet (*Citrus sinensis*); Pummelo (*Citrus maxima*); Russell River Lime (*Microcitrus inodora*); Satsuma Mandarin (*Citrus unshiu*); Sweet Lime (*Citrus limetta*); Tachibana Orange (*Citrus tachibana*); Tahiti Lime (*Citrus latifolia*); Tangelo (*Citrus x tangelo*); Tangerine (Mandarin) (*Citrus reticulata*); Tangor (*Citrus nobilis*); Trifoliate Orange (*Poncirus trifoliata*); Uniq Fruit (*Citrus aurantium* Tangelo group); cultivars, varieties and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 4 applications of this product or other Group 11 fungicide per year.
- 4) Do not use this product in citrus plant propagation nurseries.
- 5) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Clover (and stands containing Clover)</b> (See Nongrass Animal Feeds Forage, Fodder, Straw and Hay)			
<b>Corn</b>  Field Pop Sweet (Includes Seed Production)	Rust ( <i>Puccinia sorghi</i> )	6.0-9.0 (0.10-0.15)	For gray leaf spot, apply this product at the onset of disease. A second application may be required 14 days later if disease pressure persists.
	Anthraxnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leaf Spot ( <i>Cercospora sorghi</i> ) Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) Physoderma Brown Spot ( <i>Physoderma maydis</i> ) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> ) Southern Rust ( <i>Puccinia polyspora</i> )	6.0-15.5 (0.10-0.25)	For all other diseases, begin applications prior to disease development. Applications may be continued every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Early Application (V4 – V8)	6.0 (0.10)	This product may be applied early (V4 – V8) for early season disease control and beneficial physiological benefits. If mixing with herbicides, other than solo glyphosate products, Callisto®, Callisto® Xtra, or Halex® GT, consult your local Winfield Solutions, LLC representative.
	<b>Soilborne Diseases</b> Rhizoctonia Root and Stalk Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/ SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 7 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 20 applications per year when applying at the lowest rate (6.0 fl oz/A), except for field corn and field corn grown for seed.
- 4) **For field corn and field corn grown for seed**, do not exceed 2 applications per year.
- 5) Pre-Harvest Interval (PHI): 7 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Cotton	Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Glomerella gossypii</i> ) Areolate Mildew ( <i>Ramularia gossypii</i> ) Ascochyta Blight ( <i>A. gossypii</i> ) Boll Rots ( <i>Ascochyta gossypii</i> , <i>Alternaria</i> spp., <i>Diplodia</i> spp., <i>Phoma</i> spp.) Cotton Rust ( <i>Puccinia schedonnardi</i> ) Diplodia Boll Rot ( <i>Diplodia</i> spp.) Hardlock ( <i>Fusarium verticillioides</i> ) Leaf Spots and Blights ( <i>Alternaria</i> spp., <i>Ascochyta gossypii</i> , <i>Cercospora</i> spp., <i>Stemphyllium</i> spp.) Southwestern Cotton Rust ( <i>Puccinia cacabata</i> ) ( <i>Puccinia</i> spp.) Stemphyllium Leaf Spot ( <i>Stemphyllium</i> spp.) Target spot ( <i>Corynespora cassiicola</i> )	6.0-9.0 (0.1-0.15)	<p>For optimum disease control, begin applications prior to or in the early stages of disease development. Applications may be made by ground, air, or chemigation. An adjuvant may be added at specified rates. Minimum application volumes for air and ground are 5 and 10 gallons per acre, respectively.</p> <p>Target the first application of this product approximately at pinhead square to first bloom to protect the plant from diseases. Subsequent application(s) are specified on a 14- to 21-day schedule. An additional application may be made depending on environmental conditions and the health of the cotton plant.</p> <p>Under poor environmental conditions conducive to seedling disease and poor cotton growth, this product may be applied to early season cotton to suppress damping off and other diseases which result in plant stand loss.</p> <p>Do not apply more than two foliar applications of this product or other Group 11 fungicides before alternating with a fungicide that has a different mode of action.</p>
	Pythium Seedling Blight ( <i>Pythium aphanidermatum</i> ) Rhizoctonia Seedling Blight ( <i>Rhizoctonia solani</i> )	<b>In-Furrow</b> 0.40-0.80 fl oz product per 1000 row feet (0.10-0.20 oz ai per 1000 row feet)	<p>Apply this product as an in-furrow spray in 3-7 gallons of water at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of Pythium problems, or if minimum/low till programs are in place.</p> <p>See the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section for a table illustrating total fluid ounces per acre with various row spacings.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 27 fl oz of this product/A/year as a foliar spray.
- 2) Do not exceed 3 foliar applications of this product or other Group 11 fungicides per acre per year.
- 3) Pre-Harvest Interval (PHI): 45 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Cucurbits, Crop Group 9</b>  Cantaloupe Chayote Chinese Waxgourd Cucumber Gourds Honeydew Melons <i>Momordica</i> spp. (bitter melon, balsam apple) Muskmelon Watermelon Pumpkin Squash Zucchini  Including cultivars and/or hybrids of these	Alternaria Blight ( <i>Alternaria cucumerina</i> ) Anthracnose ( <i>Colletotrichum lagenarium</i> ) Belly Rot ( <i>Rhizoctonia solani</i> ) Cercospora Leaf Spot ( <i>Cercospora citrulina</i> ) Downy Mildew ( <i>Pseudoperonospora cubensis</i> ) Gummy Stem Blight ( <i>Didymella bryoniae</i> ) Leaf Spots ( <i>Alternaria</i> spp., <i>Cercospora</i> spp.) Myrothecium Canker ( <i>Myrothecium roridum</i> ) Plectosporium Blight ( <i>Plectosporium tabacinum</i> ) Powdery Mildew ( <i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i> ) Target Leaf Spot ( <i>Corynespora cassicola</i> ) Ulocladium Leaf Spot ( <i>Ulocladium cucurbitae</i> )	6.0-15.5 (0.10-0.25)	For both downy and powdery mildew, make preventative applications on a 5- to 7-day schedule.  For belly rot control, make the first application at the 1-3 leaf crop stage with a second application just prior to vine tip over or 10-14 days later whichever occurs first.  For all other diseases, make applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not tank mix this product with crop oil concentrates (COC), methylated spray oil (MSO) or silicon adjuvants.  Do not tank mix this product with malathion, methomyl, chlorpyrifos, potassium laurate or dicloran.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 4 foliar applications of this product or other Group 11 fungicides per acre per year.
- 4) Pre-Harvest Interval (PHI): 1-day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Fruiting Vegetables</b> <b>Crop Group 8-10</b>  <b>Pepper</b> Bell Pepper Non-Bell Pepper Sweet Non-Bell Pepper	Anthracnose ( <i>Colletotrichum</i> spp.) Powdery Mildew ( <i>Sphaerotheca</i> spp.)	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 7- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
<b>Eggplant</b> <b>Okra</b> <b>Pepino</b>  Including all cultivars and/or hybrids of these  <b>See specific directions for use for Tomatoes.</b>  See complete list of fruiting vegetables below.	<b>Soilborne Diseases</b> Rhizoctonia Seedling Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Complete List of Fruiting Vegetables:** African Eggplant; Bell Pepper; Eggplant; Martynia; Nonbell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant; cultivars, varieties; and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 61.5 fl oz (1.0 lb ai) of this product/A/year.
- 2) Do not apply more than 1.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 3 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 10 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Grapes and Other Small Fruit Vine Climbing Subgroup 13-07F (except fuzzy kiwifruit)</b>  Amur River Grape Kiwifruit, Hardy Maypop Muscadines Schisandra Berry  Including all cultivars and/or hybrids of these	Black Rot ( <i>Guignardia bidwellii</i> ) Downy Mildew ( <i>Plasmopara viticola</i> ) Phomopsis Cane and Leaf Spot ( <i>Phomopsis viticola</i> ) Powdery Mildew ( <i>Uncinula necator</i> )  <b>Suppression Only:</b> Botrytis Bunch Rot ( <i>Botrytis cinerea</i> )	10.0-15.5 (0.16-0.25)	Begin applications prior to disease development and continue every 10-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential foliar applications of this product or other Group 11 fungicides before alternating with a fungicide that is not in Group 11.  <b>ATTENTION</b> This product is extremely phytotoxic to certain apple varieties.  AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).  DO NOT spray this product where spray drift may reach apple trees.  DO NOT use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.  AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 9 applications per year when applying at the lowest rate (10.0 fl oz/A).  
Pre-Harvest Interval (PHI): 14 days

<b>Grasses</b> (grown for seed)	Ergot Stem Diseases Powdery Mildew ( <i>Erysiphe graminis</i> ) Rust ( <i>Puccinia</i> spp.)	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 10- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternating with a fungicide that is not in Group 11.
------------------------------------	--	-------------------------	--

**Specific Use Restrictions:**

- 1) Do not apply more than 49 fl oz (0.8 lb ai) of this product/A/year.
- 2) Do not apply more than 0.8 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 3 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 8 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Do not feed treated straw, seed, or screenings to livestock.
- 5) Pre-Harvest (swathing) Interval (PHI): 8 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Herbs &amp; Spices (except black pepper) Crop Group 19</b>  Allspice; Angelica; Anise (seed); Anise, star; Annatto; Balm; Basil; Borage; Burnet; Camomile; Caper (buds); Caraway; Caraway, Black; Cardamom; Cassia (buds); Catnip; Celery Seed; Chervil (dried); Chive; Chive, Chinese; Cinnamon; Clary; Clove (buds); Coriander (cilantro or Chinese parsley) (leaf); Coriander (seed); Costmary; Culantro (leaf and seed); Cumin; Curry (leaf); Dill (seed); Dillweed; Fennel, Common; Fennel, Florence (seed); Fenugreek; Grains of Paradise; Horehound; Hyssop; Juniper (berry); Lavender; Lemongrass; Lovage (leaf and seed); Mace; Marigold; Marjoram; Mustard (seed), Nasturtium; Nutmeg; Parsley (dried); Pennyroyal; Pepper, White; Poppy Seed; Rosemary; Rue; Saffron; Sage; Savory, Summer and Winter Sweet Bay; Tansy; Tarragon; Thyme; Vanilla; Wintergreen; Woodruff; Wormwood	Corynespora Blight ( <i>Corynespora cassiicola</i> ) Dill Blight ( <i>Cercosporidium punctum</i> ) Phoma Blight ( <i>Passalora puncta</i> )	6.0-15.5 (0.10-0.25)	<p>Begin applications at the onset of disease development and continue on a 7-day schedule, following the resistance management guidelines. Applications may be made by ground only. An adjuvant may be added at specified rates. Use a minimum of 30 gallons of water per acre.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
Wasabi	Fusarium Rhizome and Root Rot ( <i>Pythium</i> spp.)	6.2-15.4 (0.10-0.25)	<p>Begin applications at the onset of disease development and continue on a 7-day schedule, following the resistance management guidelines. Applications may be made by ground or through the irrigation system (chemigation). An adjuvant may be added at specified rates. Use a minimum of 30 gallons of water per acre.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Leafy Vegetables (except Brassica), Crop Group 4</b>  Amaranth Arugula Cardoon Celery Celtuce Chervil Chrysanthemum Edible Corn Salad Cress Dandelion Dock Endive Fennel Lettuce Head and Leaf Orach Parsley Purslane Radicchio Rhubarb Spinach Swiss Chard  Including cultivars and/or hybrids of these	<b>Foliar Diseases</b> Alternaria Leaf Spot ( <i>Alternaria sonchi</i> , A. spp.) Anthracnose ( <i>Microdochium panattonianum</i> , <i>Colletotrichum dematium</i> ) Ascochyta Leaf Spot ( <i>Ascochyta</i> spp.) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Rust ( <i>Puccinia</i> spp.) ( <i>Uromyces</i> spp.) Septoria Leaf Spot ( <i>Septoria petroselinii</i> ) White Rust ( <i>Albugo occidentalis</i> )	6.0-15.5 (0.10-0.25)	For both downy and powdery mildew, make preven- tative applications on a 5- to 7-day schedule.  For all other diseases, begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemi- gation. An adjuvant may be added at specified rates.  Do not apply more than one application of this prod- uct or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.  <b>ATTENTION:</b> Applications of this product to leafy vegetable foliage have contributed to phytotoxicity under certain circumstances. Proceed with caution with regard to tank mixes and adjuvants when treating all leafy vegetables with this product.  This product must not be tank-mixed on leaf lettuce with permethrin, fosetyl-al, lambda-cyhalothrin, or another product that may increase the penetration of this product into the leaf surface, such as but not limited to, silicone wetters.
	Downy Mildew ( <i>Bremia lactucae</i> ) Powdery Mildew ( <i>Erysiphe cichoracearum</i> )	12.0-15.5 (0.20-0.25)	
	<b>Soilborne Diseases</b> Webb Blight Bottom Rot Crater Rot Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see direc- tions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Legume Vegetables, Dry and Succulent, Crop Group 6 and Legume Vegetables, Foliage of any Cultivar of Bean (<i>Phaseolus</i> spp.) and Field Pea (<i>Pisum</i> spp.), Crop Group 7</b>  Bean ( <i>Lupinus</i> spp.) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin) Bean ( <i>Phaseolus</i> spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean) Bean ( <i>Vigna</i> spp.) (includes adzuki bean, asparagus bean, blackeye pea, cowpea, catjang, Chinese longbean, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean) Bean ( <i>Glycine</i> max) Soybean, Immature Seed (edamame) Broad bean (fava bean) ( <i>Vicia faba</i> ) Chickpea (garbanzo bean) ( <i>Cicer arietinum</i> ) Guar ( <i>Cyamopsis tetragonoloba</i> ) Jackbean ( <i>Canavalia ensiformis</i> ) Lablab Bean (hyacinth bean) ( <i>Lablab purpureus</i> ) Lentil ( <i>Lens esculenta</i> ) Pea ( <i>Pisum</i> spp.) (includes dwarf pea, edible-pod pea, English pea, garden pea, green pea, field pea, snow pea, sugar snap pea) Pigeon Pea ( <i>Cajanus cajan</i> ) Sword Bean ( <i>Canavalia gladiata</i> )	Bean Rust ( <i>Uromyces appendiculatus</i> )	6.0 (0.10)	Begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Use the higher rates under severe disease pressure. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. For best results on rust, use a nonionic surfactant.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Alternaria Blight ( <i>Alternaria</i> spp.) Alternaria Leaf Spot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichum lindemuthianum</i> ) Ascochyta Blight ( <i>Mycosphaerella pinodes</i> ) Ascochyta Leaf and Pod Spot ( <i>Ascochyta</i> spp.) Ascochyta Leaf Spot ( <i>Ascochyta phaseolorum</i> ) Rust ( <i>Phakopsora</i> spp.) Southern Blight ( <i>Sclerotium rolfsii</i> ) Web Blight ( <i>Rhizoctonia solani</i> )	6.0-15.5 (0.10-0.25)	
	<b>Soilborne Diseases</b> Rhizoctonia Root Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/ SEEDLING DISEASE CONTROL</b> section.  This product may be applied to the furrow and covering soil at planting time in a 7-inch band. Avoid a concentrated stream directly on the seed or delayed emergence may occur.  If using a narrow spray as an in-furrow spray, adjust the spray stream to hit the soil next to the seed but not hit the seed.  <b>NOTE:</b> Conduct a seed safety test with your crop before making in-furrow applications.

#### Specific Use Restrictions:

- Do not apply more than 92.3 fl oz 1.5 lb ai of this product/A/year.
- Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- Pre-Harvest Interval (PHI):
  - Dry Legume Vegetables (dry bean and dry pea seeds) - 14 days
  - Succulent Beans and Peas - 0 day
- For use on soybeans, please refer to the soybean crop directions for use.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Mint</b> (Fresh or for processing into mint oil)	Leaf Spot ( <i>Ramularia</i> spp.) ( <i>Alternaria</i> spp.) ( <i>Phoma</i> , spp.) Powdery mildew ( <i>Erysiphe</i> spp.) Rust ( <i>Puccinia menthae</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Seedling Root Rot, Basal Stem Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 46 fl oz (0.75 lb ai) of this product/A/year.
- 2) Do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 7 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI):
  - Processed Mint - 7 days
  - Fresh Mint - 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Nongrass Animal Feeds Forage, Fodder, Straw and Hay, Crop Group 18</b>  For pure/mixed stands of the following or stands mixed with grasses:  Alfalfa ( <i>Medicago sativa</i> subsp. <i>sativa</i> ) Bean, Velvet ( <i>Mucuna pruriens</i> var. <i>utilis</i> ) Clover ( <i>Trifolium</i> spp., <i>Melilotus</i> spp.) Kudzu ( <i>Pueraria lobata</i> ) Lespedeza ( <i>Lespedeza</i> spp.) Lupin ( <i>Lupinus</i> spp.) Sainfoin ( <i>Onobrychis viciifolia</i> ) Trefoil ( <i>Lotus</i> spp.) Vetch ( <i>Vicia</i> spp.) Vetch, Crown ( <i>Coronilla varia</i> ) Vetch, Milk ( <i>Astragalus</i> spp.)	Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum trifolii</i> ) Black Patch ( <i>Rhizoctonia leguminicola</i> ) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Common Leaf Spot ( <i>Pseudopezizza solani</i> ) Downy Mildew ( <i>Peronospora</i> spp.) Leaf Spot ( <i>Leptosphaerulina briosiai</i> ) Powdery Mildew ( <i>Oidium</i> spp., <i>Erysiphe</i> spp.) Rhizoctonia and Stem Blight ( <i>Rhizoctonia solani</i> ) Rust ( <i>Phakopsora</i> spp.) ( <i>Uromyces</i> spp.) Spring Black Stem and Leaf Spot ( <i>Phoma medicaginis</i> ) Stagonospora Leaf Spot ( <i>Stagonospora meliloti</i> ) Stemphyllium Leaf Spot ( <i>Stemphyllium</i> spp.) Summer Black Stem and Leaf Spot ( <i>Cercospora medicaginis</i> ) Yellow Leaf Blotch ( <i>Leptotrichilia medicaginis</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue, following the resistance management guidelines. Use the higher rates under severe disease pressure. Applications may be made by ground, air or chemigation. For best results, use an adjuvant such as a non-ionic surfactant or crop oil concentrate at specified labeled rates.  For management of outbreaks of Asian soybean rust and other Puccinia species on alternate host species such as kudzu, lespedeza, trefoil and vetch, apply this product to forages grown in the vicinity of soybeans and other legume crops (beans and peas) as a part of an Asian rust disease management strategy. Consult with local experts and university extension agents for the latest advice.  Do not apply more than three sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Sclerotinia Crown Rot and Wilt on Clover ( <i>Sclerotinia trifoliorum</i> )	10.0 (0.17)	

**Specific Use Restrictions:**

- 1) Do not apply more than 15.5 fl oz/A (0.25 lb ai/A) of this product per cutting.
- 2) Do not apply more than 46.0 fl oz/A (0.75 lb ai/A) of this product/A/year.
- 3) Do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 4) Do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 7 applications per year when applying at the lowest rate (6 fl oz/A).
- 5) Pre-Harvest Interval (PHI): 14 days (forage and hay)
- 6) Do not apply within 14 days of grazing for forage and hay.
- 7) Not for use on rangeland.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Oilseed Crops Crop Group 20</b>  Crambe Flax Mustard, Indian Mustard, Field Mustard, Black Rapeseed Rapeseed, Indian Safflower Sunflower  Including all cultivars and/or hybrids of these  See complete list of oilseed crops below.	Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Downy Mildew ( <i>Plasmopora halstedii</i> , <i>Plasmopora helianthi</i> ) Pasma ( <i>Septoria linicola grass</i> ) Sunflower Rust ( <i>Puccinia helianthi</i> )	6.0-15.5 (0.1-0.25)	Apply 6.0 fl oz at early bud followed by 14.0 fl oz at about 45 days before harvest. A third application of 7.0 fl oz may be made 30 days before harvest. Applications may be made by ground, air or chemigation. Use a minimum of 10 gallons of water per acre for ground applications.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**Complete List of Oilseed Crops:** Borage; Calendula; Castor Oil Plant; Chinese Tallowtree; Cottonseed; Crambe; Cuphea; Echium; Euphorbia; Evening Primrose; Flax Seed; Gold of Pleasure; Hare's Ear Mustard; Jojoba; Lesquerella; Lunaria; Meadowfoam; Milkweed; Mustard Seed; Niger Seed; Oil Radish; Poppy Seed; Rapeseed; Rose Hip; Safflower; Sesame; Stokes Aster; Sunflower; Sweet Rocket; Tallowwood; Tea Oil Plant; Vernonia; cultivars, varieties, and/or hybrids of these.

#### Specific Use Restrictions:

- 1) Do not apply more than 27.7 fl oz (0.45 lb ai) of this product/A/year.
- 2) Do not apply more than 0.45 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 1 application of this product per year when applying at the highest rate (15.5 fl oz/A) or 4 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 30 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Peanuts	<b>Soilborne Diseases – early season (in-furrow application)</b>  Aspergillus Crown Rot <i>(Aspergillus niger)</i> Pythium Damping-Off <i>(Pythium spp.)</i> Stem Rot/White Mold Suppression <i>(Sclerotium rolfsii)</i>	0.40-0.80 fl oz/1000 row feet	Apply this product in-furrow at planting for control of various seed/seedling diseases including early season suppression of stem rot. See directions and rates under <b>PRODUCT INFORMATION</b> section.
	<b>Soilborne Diseases – midlate season</b> Rhizoctonia Peg and Pod Rot <i>(Rhizoctonia solani)</i> Stem Rot/White Mold <i>(Sclerotium rolfsii)</i>  <b>Suppression Only:</b> Cylindrocladium Black Rot <i>(Cylindrocladium crotalariae)</i> Pythium Pod Rot <i>(Pythium myriotylum)</i>	12.0-24.5 (0.20-0.40)	<p>Apply product at approximately 60 and 90 days after planting as a foliar application. This application regime may be applied earlier in the season if environmental conditions favor disease development. These two applications will provide protection against the soilborne diseases and will also provide control of the foliar diseases listed for a 10- to 14-day period after each spray.</p> <p>Under heavy disease pressure and/or where there is high rainfall and/or irrigation, use 18.5-24.5 fl oz/A. For light disease pressure and dry environmental conditions (non-irrigated, low rainfall), use 12.0-24.5 fl oz/A.</p> <p>For control of Pythium, a rate of 24.5 fl oz/A is required. Additional applications of other fungicides on a leaf spot application schedule will be required to provide season long disease control of the leaf spot diseases. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p>
	<b>Foliar Diseases</b> Early Leaf Spot <i>(Cercospora arachidicola)</i> Late Leaf Spot <i>(Cercosporidium personatum)</i> Rust <i>(Puccinia arachidis)</i> Web Blotch <i>(Phoma arachidicola)</i>	6.0-18.5 (0.10-0.30)	<p>For foliar disease control only, a lower rate of this product may be applied on a 10- to 14 day interval.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 49 fl oz (0.8 lb ai) of this product/A/year.
- 2) Do not apply more than 0.8 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year when applying at the highest rate (24.5 fl oz/A) or 8 applications per year when applying at the lowest rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 14 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Pistachios</b>	Alternaria Late Blight ( <i>Alternaria alternata</i> ) Botryosphaeria Panicle and Shoot Blight ( <i>Botryosphaeria dothidea</i> ) Septoria Leaf Spot ( <i>Septoria pistaciarum</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue at 7- to 21-day interval following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 7 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Potatoes	Black Dot ( <i>Colletotrichum coccodes</i> ) Early Blight ( <i>Alternaria solani</i> ) Late Blight ( <i>Phytophthora infestans</i> ) Powdery Mildew ( <i>Erysiphe cichoracearum</i> )	6.0-20.0 (0.10-0.33)	<p><b>Early blight</b> - For a 7-day application schedule, use 6.2 fl oz product/A. For a 14-day application schedule, use the 12.0 fl oz product/A rate.</p> <p><b>Late blight</b> - Apply this product at 12.0 fl oz product/A on a 7-day schedule. Initiate late blight applications in a preventative schedule prior to disease development according to local practices. If late blight symptoms develop or conditions favor disease, switch immediately to a non-Group 11 fungicide, using a 5-day schedule. Addition of a spreader/sticker may improve coverage.</p> <p>For all other diseases, begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Use the high rate and the shorter interval if disease epidemics are severe. Applications may be made by ground, air or chemigation.</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
	<b>Soilborne Diseases</b> Black Dot ( <i>Colletotrichum coccodes</i> ) Black Scurf ( <i>Rhizoctonia solani</i> ) Silver Scurf ( <i>Helminthosporium solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 6 applications of this product per year when applying at the highest rate (20.0 fl oz/A) or 20 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 14 days

Quinoa	Leaf Spot ( <i>Ascochyta hyalospora</i> ) Stalk Rot ( <i>Phoma exigua</i> )	12 (0.20)	Apply prior to disease development. This product can be applied by either ground, chemigation, or aerial application. An adjuvant may be added at specified rates.
--------	--	--------------	--

**Specific Use Restrictions:**

- 1) Do not apply more than 24.6 fl oz (0.40 lb ai) of this product/A/year.
- 2) Do not apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year.
- 4) Do not apply within 14 days of grazing.
- 5) Pre-Harvest Interval (PHI):
  - 30 days
  - Forage and Hay: 7 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Rice	<b>Sheath/Stem Diseases</b> Sheath Blight ( <i>Rhizoctonia solani</i> )	6.0-18.5 (0.10-0.30)	Apply this product prior to disease development. Applications may be made by ground, air or chemigation. For aerial application, volumes of 5-10 GPA. An adjuvant may be added at specified rates.
	Aggregate Sheath Spot ( <i>Ceratobasidium oryzae-sativae</i> = <i>Rhizoctonia oryzae-sativae</i> ) Black Sheath Rot ( <i>Gaeumannomyces graminis</i> var. <i>graminis</i> ) Sheath Spot ( <i>Rhizoctonia oryzae</i> ) Stem Rot ( <i>Magnaporthe salvinii</i> = <i>Sclerotium oryzae</i> = <i>Nakateae sigmoidea</i> )	9.0-18.5 (0.15-0.30)	For sheath blight control, application rates may vary from 9.0 to 12.0 fl oz/A depending on the growth stage of the rice and the severity of the disease.  For other stem/sheath diseases including stem rot, black sheath rot, aggregate sheath spot and sheath spot, apply when disease is less than 4 inches above water line usually between panicle differentiation (PD) +5 days to PD +10 days or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied.
	<b>Foliar Diseases</b> Brown Leaf Spot ( <i>Cochliobolus miyabeanus</i> ) Leaf Smut ( <i>Entyloma oryzae</i> ) Narrow Brown Leaf Spot ( <i>Cercospora janseana</i> = <i>Cercospora oryzae</i> )		For foliar and panicle diseases, apply this product prior to disease development. For blast control, apply as a preventative treatment prior to favorable conditions for blast development. For panicle blast, make an application at mid-boot to boot-split but prior to full head emergence. Make a second application when panicles are approximately 60-90% emerged from the boot (7-14 days later).
	<b>Panicle Diseases</b> Kernel Smut ( <i>Tilletia barclayana</i> = <i>Neovossia barclayana</i> ) Panicle Blast ( <i>Pyricularia grisea</i> )		When applied for panicle blast on continuous rice acreage (no rotation to other crops), do not make more than two sequential foliar applications of this product or other Group 11 fungicides over multiple years before alternating with a fungicide with a different mode of action.

**Specific Use Restrictions:**

- 1) Do not apply more than 43 fl oz (0.70 lb ai) of this product/A/year.
- 2) Do not apply more than 0.70 lb ai/A/year of azoxystrobin-containing products
- 3) Do not exceed 2 foliar applications of this product or other Group 11 fungicides per acre per year.
- 4) Do not treat rice fields used for aquaculture of fish and crustaceans.
- 5) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat.  
Use care in making applications near non-target aquatic habitats
- 6) Do not allow release of irrigation or flood water for at least 14 days after the last application.
- 7) Pre-Harvest Interval (PHI): 28 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Sorghum	Anthracnose ( <i>Colletotrichum graminicola</i> ) Gray Leaf Spot ( <i>Cercospora sorghi</i> )	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development. Use the high rates under conditions favorable for severe disease pressure, dense plant canopies, or when susceptible varieties are planted. Contact extension personnel for local economic thresholds and timings for specific diseases in your area. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Damping-Off ( <i>Rhizoctonia solani</i> , <i>Pythium aphanadermatum</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) For grain and stover, do not apply more than 46 fl oz (0.75 lb ai) of this product/A/year.
- 2) For grain and stover, do not apply more than 0.75 lb ai/A/year of azoxystrobin-containing products.
- 3) For grain and stover, do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 7 applications per year when applying at the lowest application rate (6.0 fl oz/A).
- 4) For forage, do not apply more than 31 fl oz (0.5 lb ai) of this product/A/year.
- 5) For forage, do not apply more than 0.5 lb ai/A/year of azoxystrobin-containing products.
- 6) For forage, do not exceed 2 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 5 applications per year when applying at the lowest rate (5.0 fl oz/A).
- 7) Pre-Harvest Interval (PHI): 14 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Soybeans Soybean, Immature Seed (edamame)	Aerial Blight ( <i>Rhizoctonia solani</i> ) Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum truncatum</i> ) Brown Spot ( <i>Septoria glycines</i> ) Cercospora Blight and Leaf Spot ( <i>Cercospora kikuchii</i> ) Frogeye Leaf Spot ( <i>Cercospora sojina</i> ) Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> ) Rust ( <i>Phakopsora</i> spp.)	6.0-15.5 (0.10-0.25)	<p>Begin applications prior to disease development. Use the high rates under conditions favorable for severe disease pressure, dense plant canopies, or when susceptible varieties are planted. Contact Extension personnel for local economic thresholds and timings for specific diseases in your area. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates. For best results, when making applications at the lower rate use an adjuvant such as a non-ionic surfactant or crop oil concentrate at specified labeled rates.</p> <p>Soybean rust: This product may be used at 4 fl oz/A when tank mixed with a triazole registered for use on soybean rust.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>
	<b>Soilborne Diseases</b> Rhizoctonia solani ( <i>Rhizoctonia solani</i> ) Southern blight ( <i>Sclerotium rolfsii</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications when applying at the lowest rate (6.0 fl oz/A), except for soybean forage and hay.
- 4) **For soybean forage and hay**, do not exceed 1 application per year at the 15.5 fl oz (0.25 lb ai)/A rate.
- 5) Pre-Harvest Interval (PHI):
  - Soybeans (beans) - 14 days.
  - Soybeans (forage and hay) - 0 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Stone Fruits, Crop Group 12-12</b>  Apricot Cherry, Sweet Cherry, Tart Nectarine Peach Plum Plumcot Prune	Brown Rot Blossom Blight and Fruit Rot <i>(Monilinia fructicola, M. laxa)</i>	12.0-15.5 (0.20-0.25)	For brown rot blossom blight, begin applications at early bloom and continue through petal fall. For brown rot on fruit, this product may be applied to fruit up to the day of harvest.  For scab, begin applications at petal fall and continue at 7- to 14-day intervals.
	Scab <i>(Cladosporium carpophilum)</i> Alternaria Spot and Fruit Rot ( <i>Alternaria alternata</i> ) Anthracnose <i>(Colletotrichum prunicola, C. gloeosporioides)</i> Leaf Rust <i>(Tranzschelia discolor)</i> Powdery Mildew <i>(Sphaerotheca pannosa, Podosphaera clandestina)</i> Shot Hole <i>(Wilsonomyces carpophilus)</i>	6.0-15.5 (0.10-0.25)	For all other diseases, begin application at the onset of disease as a protectant fungicide and continue on a 7- to 14-day schedule.  For peaches only, 9.0-15.5 fl oz of this product may be used for scab control.  Applications may be made by ground, air or chemigation.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.

**Complete List of Stone Fruit Crops:** Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Sugarcane	Brown Rust ( <i>Puccinia melanocephala</i> ) Orange Rust ( <i>Puccinia kuehnii</i> )	9.0-12.0 (0.15-0.20)	<p>Begin applications prior to rust development and continue every 14-28 days following resistance management guidelines. Scout fields and begin applications at the earliest sign of rust. An adjuvant may be used at specified rates. For ground applications, apply in sufficient water volume for adequate coverage and canopy penetration. Applications may be made by ground, air or chemigation.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicide, before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 49 fl oz (0.80 lb ai) of this product/A/year.
- 2) Do not apply more than 0.80 lb ai/A per year of azoxystrobin-containing products.
- 3) Do not exceed 4 foliar applications of this product or other Group 11 fungicide per acre per year.
- 4) Pre-Harvest Interval (PHI): 30 days
- 5) When applying by air, use no less than 5 gallons spray solution per acre.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Ti Palm, Leaves and Roots</b>	<b>Foliar Diseases</b> Alternaria Leaf Spot ( <i>Alternaria</i> spp., <i>A. alternata</i> ) Ascochyta Leaf Spot ( <i>Ascochyta cynarae</i> ) Phyllostica Leaf Spot ( <i>Phyllostica</i> spp.) Rust ( <i>Uromyces betae</i> , <i>Puccinia helianthi</i> ) White Rust ( <i>Albugo</i> <i>tragopogonis</i> )	6.0-20.0 (0.10-0.33)	For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Cercospora Leaf Spot ( <i>Cercospora betae</i> , <i>C. pastinaceae</i> ) Powdery Mildew ( <i>Erysiphe polygoni</i> , <i>Leveillula taurica</i> )	9.0-15.5 (0.15-0.25)	
	<b>Soilborne Diseases</b> Circular Spot, Southern Blight ( <i>Sclerotium rolfsii</i> ) Pythium Root Rot ( <i>Pythium</i> <i>aphanidermatum</i> ) Rhizoctonia Stem Canker, Crown Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do exceed 6 applications of this product per year for *Phyllostica* spp.
- 4) Do exceed 8 applications of this product per year for *Cercospora* spp.
- 5) For all other disease species, do not exceed 6 applications of this product per year when applying at the highest rate (20 fl oz/A) or 20 applications when applying at the lowest rate (6.0 fl oz/A).
- 6) Apply as an in-furrow spray in a minimum of 10 gallons per acre.
- 7) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Tobacco</b>	Blue Mold ( <i>Peronospora tabacina</i> ) Frogeye Leaf Spot ( <i>Cercospora nicotianae</i> ) Target Spot ( <i>Rhizoctonia solani</i> )	6.0-12.0 (0.10-0.20)	<p>Begin applications prior to disease development or at first indication that blue mold is in the area. Do not apply this product as a curative application. If blue mold is present in the field, initiate applications with Acrobat MZ® prior to an application with this product. Apply on a 7- to 14-day interval with shorter intervals under conditions conducive to disease development.</p> <p>Applications may be made by ground, air or chemigation.</p> <p>For ground applications, apply in sufficient water volume for adequate coverage and canopy penetration.</p> <p>For aerial application, apply in 10-15 GPA.</p> <p>Do not apply this product on greenhouse seedlings.</p> <p>Tank mixing this product with insecticides formulated as emulsifiable concentrates (EC) or containing high amounts of solvents, may cause some crop injury.</p> <p>Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>NOTE: This product may enhance weather flecking on the leaves of certain tobacco types. This does not affect yield and quality.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 32 fl oz (0.52 lb ai) of this product/A/year.
- 2) Do not apply more than 0.52 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product per year when applying at the highest rate (12 fl oz/A) or 5 applications when applying at the lowest rate 6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 21 days

<b>Tobacco Transplants in greenhouse</b>	Target Spot ( <i>Rhizoctonia solani</i> )	6.0 (0.10)	Apply 6 oz/A or 0.14 oz (4ml)/1000 sq ft in enough water for thorough coverage. Make only one application prior to transplanting.
<b>GA, KY, IN, MD, MO, NC, OH, PA, SC, TN and VA only</b>			

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Tomatoes</b> <b>Tomatillos</b> <b>Subgroup 8-10A</b>  Including all cultivars and/or hybrids of these  See complete list of tomato crops below.	Anthracnose <i>(Colletotrichum</i> <i>coccodes)</i> Black Mold <i>(Alternaria</i> <i>alternata)</i> Buckeye Rot <i>(Phytophthora</i> <i>spp.)</i> Early Blight <i>(Alternaria solani)</i> Powdery Mildew <i>(Oidiopsis sicula)</i> Septoria Leaf Spot <i>(Septoria</i> <i>lycopersici)</i> Target Spot <i>(Corynespora</i> <i>cassiicola)</i>	5.0-6.2 (0.08-0.10)	Begin applications prior to disease development and continue following the resistance management guidelines. For late blight, apply this product at 5- to 7-day intervals. For all other tomato diseases, apply on 7- to 21-day intervals. Applications may be made by ground, air or chemigation.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.  Under certain weather conditions (particularly high temperatures) this product in combination with high rates of silicone based or oil containing (petroleum or crop) additives or adjuvants may cause injury. Do not exceed 0.125% adjuvant (v/v).  A tank mixture with dimethoate may cause crop injury.
	Late Blight <i>(Phytophthora</i> <i>infestans)</i>	6.2 (0.10)	For fresh market tomatoes, do not use adjuvants or tank mix this product with any emulsifiable concentrate (EC) product.

**Complete List of Tomato Crops:** Bush Tomato; Cocona; Currant Tomato; Garden Huckleberry; Goji Berry; Groundcherry; Naranjilla; Sunberry; Tomatillo; Tomato; Tree Tomato; cultivars, varieties, and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 37 fl oz (0.6 lb ai) of this product/A/year.
- 2) Do not apply more than 0.6 lb ai/A/year of azoxystrobin-containing products.
- 3) Do exceed 5 applications per year of this product when applying at the highest application rate (6.2 fl oz/A) or 7 applications per year when applying at the lowest application rate (5.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Tree Nuts, Crop Group 14-12 (except Pistachios)</b>  Beechnut Brazil Nut Butternut Cashew Chestnut Chinquapin Filbert (hazelnut) Hickory Macadamia Pecan Walnut  <b>Almonds, Pistachios (see specific use instructions)</b>	Alternaria Leaf and Fruit Spot ( <i>Alternaria alternata</i> ) Anthracnose ( <i>Colletotrichum acutatum</i> , <i>Glomerella cingulata</i> ) Eastern Filbert Blight ( <i>Anisogramma anomale</i> ) Late Blight ( <i>Alternaria alternata</i> ) Scab ( <i>Cladosporium carpophilum</i> ) Septoria Leaf Spot ( <i>Septoria pistaciarum</i> ) Shot Hole ( <i>Wilsonomyces carpophilus</i> )  Blossom Blight ( <i>Monilinia laxa</i> , <i>M. fructicola</i> )	6.0-12.0 (0.10-0.20)	Begin applications prior to disease development and continue at 7- to 21-day intervals following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.  For blossom blight, begin applications at early bloom and continue through petal fall.

**Complete List of Tree Nut Crops:** African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not apply more than 73.8 fl oz (1.2 lb ai) of this product/A/year.
- 2) Do not apply more than 1.2 lb ai/A/year of azoxystrobin-containing products.
- 3) Do exceed 6 applications per year of this product when applying at the highest application rate (12.0 fl oz/A) or 12 applications per year when applying at the lowest application rate (6.0 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 45 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Tropical Fruit</b> Acerola Atemoya Avocado Biriba Canistel Cherimoya Custard Apple Dragon Fruit Feijoa Guava Llama Jaboticaba Jackfruit Longan Loquat Lychee Mango Papaya Passionfruit Pawpaw Persimmon Pulasan Rambutan Sapodilla Sapote, Black Sapote, Mamey Sapote, White Soursop Star Apple Starfruit Sugar Apple Spanish Lime Tamarind	Anthracnose ( <i>Colletotrichum</i> spp.) Cercospora Leaf Spot ( <i>Cercospora</i> spp.) Powdery Mildew ( <i>Erysiphe</i> spp.) Rust ( <i>Puccinia</i> spp.)	6.0-15.5 (0.10-0.25)	Begin applications prior to disease development and continue on a 10- to 14-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Follow the resistance management guidelines in the Resistance Management Section. Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	<b>Soilborne Diseases</b> Seedling Root Rot Basal Stem Rot ( <i>Rhizoctonia solani</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 92.3 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Vegetables, Leaves of Root and Tuber Group and Root Subgroup</b>  Beet, Garden and Sugar <sup>1,2</sup> Burdock <sup>1,2</sup> Carrot <sup>1,2</sup> Cassava, Bitter and Sweet <sup>1</sup> Celeriac (celery root) <sup>1,2</sup> Chervil, Turnip- Rooted <sup>1,2</sup> Chicory <sup>1,2</sup> Dasheen (taro) <sup>1</sup> Ginseng <sup>2</sup> Horseradish <sup>2</sup> Parsley, Turnip- Rooted <sup>2</sup> Parsnip <sup>1,2</sup> Radish <sup>1,2</sup>  Radish, Oriental (daikon) <sup>1,2</sup> Rutabaga <sup>1,2</sup> Salsify <sup>2</sup> Salsify, Black <sup>1,2</sup> Salsify, Spanish <sup>2</sup> Skirret <sup>2</sup> Sweet Potato <sup>1</sup> Tanier <sup>1</sup> Turnip <sup>1,2</sup> Yam, True <sup>1</sup>	<b>Foliar Diseases</b> Alternaria Leaf Spot <i>(Alternaria spp., A. alternata)</i> Ascochyta Leaf Spot <i>(Ascochyta cynarae)</i> Rust <i>(Uromyces betae, Puccinia helianthi)</i> White Rust <i>(Albugo tragopogonis)</i>	6.0-20.0 (0.10-0.33)	For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines.  Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Cercospora Leaf Spot <i>(Cercospora betae, C. pastinaceae)</i> Powdery Mildew <i>(Erysiphe polygoni, Leveillula taurica)</i>	9.0-15.5 (0.15-0.25)	
	<b>Soilborne Diseases</b> Circular Spot, Southern Blight <i>(Sclerotium rolfsii)</i> Pythium Root Rot <i>(Pythium aphanidermatum)</i> Rhizoctonia Stem Canker, Crown Rot <i>(Rhizoctonia solani)</i>	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.  <b>For sugar beets</b> apply 3-7 inch banded applications in a minimum of 10 gallons per acre at the 2- to 8-leaf stage. Do not apply as a dribble application over the seed row. Tank mixtures of this product with crop oil concentrates (COC) or methylated spray oil (MSO) may result in crop injury. If cool soil conditions are expected after planting which could result in an extended period of plant emergence, this product must not be applied in-furrow. If using this product at the time of planting, do not use a starter fertilizer with it.

<sup>1</sup> Leaves of Root and Tuber Vegetables, Crop Group 2

<sup>2</sup> Root Vegetable, Crop Subgroup 1A

#### Specific Use Restrictions:

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 6 applications of this product per year when applying at the highest rate (20.0 fl oz/A) or 20 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Apply as an in-furrow spray in a minimum of 10 gallons per acre.
- 5) Pre-Harvest Interval (PHI): 0 day

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Vegetables, Tuberous and Corm Subgroup 1C</b>  Arracacha Arrowroot Artichoke, Chinese and Jerusalem Canna, Edible Cassava, Edible, Bitter and Sweet Chayote (root) Chufa Dasheen (Taro) Ginger Leren Potato Sweet Potato Tanier Turmeric Yam, Bean Yam, True	<b>Foliar Diseases</b> Alternaria Leaf Spot ( <i>Alternaria</i> spp., <i>A. Alternata</i> ) Ascochyta Leaf Spot ( <i>Ascochyta cynarae</i> ) Rust ( <i>Uromyces betae</i> , <i>Puccinia helianthi</i> ) White Rust ( <i>Albugo tragopogonis</i> )	6.0-20.0 (0.10-0.33)	For powdery mildew, make preventative applications on a 5- to 7-day schedule. For all other diseases, begin applications prior to disease development and continue every 7-14 days following the resistance management guidelines.  Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.  Do not apply more than one application of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.
	Cercospora Leaf Spot ( <i>Cercospora betae</i> , <i>C. pastinaceae</i> ) Powdery Mildew ( <i>Erysiphe polygoni</i> , <i>Leveillula taurica</i> )	9.0-15.5 (0.15-0.25)	
	<b>Soilborne Diseases</b> Circular Spot, Southern Blight ( <i>Sclerotium rolfsii</i> ) Rhizoctonia Stem Canker, Crown Rot ( <i>Rhizoctonia solani</i> ) Pythium Root Rot ( <i>Pythium aphanidermatum</i> )	0.40-0.80 fl oz/1000 row feet	For soilborne/seedling disease control, see directions and rates under the <b>SOILBORNE/SEEDLING DISEASE CONTROL</b> section.

**Specific Use Restrictions:**

- 1) Do not apply more than 123 fl oz (2.0 lb ai) of this product/A/year.
- 2) Do not apply more than 2.0 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 6 applications of this product per year when applying at the highest rate (20.0 fl oz/A) or 20 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 14 days

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
<b>Watercress</b>	Cercospora Leaf Spot ( <i>Cercospora</i> spp.)	6.0-15.5 (0.10-0.25)	<p>Begin applications prior to disease development and continue on a 7- to 10-day schedule, following the resistance management guidelines. Applications may be made by ground, air or chemigation. An adjuvant may be added at specified rates.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 93.2 fl oz (1.5 lb ai) of this product/A/year.
- 2) Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 5 applications of this product per year when applying at the highest rate (15.5 fl oz/A) or 15 applications per year when applying at the lowest rate (6 fl oz/A).
- 4) Pre-Harvest Interval (PHI): 7 days

<b>Cereals</b>  <b>Wheat</b> <b>Triticale</b>	Leaf Rust ( <i>Puccinia triticina</i> = <i>Puccinia recondita</i> f.sp. <i>tritici</i> ) Septoria Leaf and Glume Blotch ( <i>Septoria tritici</i> , <i>Septoria nodorum</i> ) Stem Rust ( <i>Puccinia graminis</i> ) Stripe Rust ( <i>Puccinia striiformis</i> ) Tan Spot ( <i>Pyrenophora</i> <i>tritici-repentis</i> )	4.0-12.0 (0.07-0.20)	<p>Apply this product prior to disease development. Applications may be made by ground, air or chemigation. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicide before alternation with a fungicide that is not in Group 11.</p>
--	--	-------------------------	---

**Specific Use Restrictions:**

- 1) Do not apply more than 24.6 fl oz (0.40 lb ai) of this product/A/year.
- 2) Do not apply more than 0.40 lb ai/A/year of azoxystrobin-containing products.
- 3) Do not exceed 2 applications of this product or other Group 11 fungicide per year.
- 4) Do not apply after Feekes 10.54.
- 5) Pre-Harvest Interval (PHI): 7 days (forage and hay)
- 6) Do not apply within 14 days of grazing.

Crop	Target Diseases	Use Rate fl oz product/A (lb ai/A)	Application Instructions
Wild Rice	<p>Brown Spot (<i>Bipolaris oryzae</i> or <i>Bipolaris sorokiana</i>)</p> <p>Also known as <i>Helminthosporium oryzae</i> and <i>H. sativum</i></p> <p>Stem Rot (<i>Nakataea sigmoidea</i>)</p>	12.5-15.5 (0.20-0.25)	<p>Apply this product prior to disease development. Applications may be made by ground, air, or chemigation. For aerial application, use volumes of 5-10 GPA. An adjuvant may be added at specified rates.</p> <p>For foliar diseases, apply prior to disease development during tillering, boot, early heading, or at initial sign of disease. Under heavy disease pressure and conditions favorable for disease development, a second application may be applied.</p> <p>Do not apply more than two sequential applications of this product or other Group 11 fungicide before alternation with a fungicide that is not in Group 11.</p>

**Specific Use Restrictions:**

- 1) Do not apply more than 43 fl oz (0.70 lb ai) of this product/A/year.
- 2) Do not apply more than 0.70 lb ai/A/year of azoxystrobin-containing products.
- 3) Do exceed 2 applications of this product or other Group 11 fungicide per year.
- 4) Do not treat wild rice fields used for aquaculture of fish and crustaceans.
- 5) Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Use care in making applications near non-target aquatic habitats.
- 6) Do not allow release of irrigation or flood water for at least 14 days after the last application.
- 7) Pre-Harvest Interval (PHI): 28 days

**Tetraban Fungicide Rate Conversion Chart**

<b>Fl Ounces Product/A</b>	<b>Lb ai/A</b>	<b>Treated Acres/Gal Product</b>
4.0	0.07	32.0
5.0	0.08	25.6
5.5	0.09	23.2
6.0	0.10	21.3
6.2	0.10	21.3
7.0	0.11	18.3
8.5	0.14	15.4
9.0	0.15	14.2
9.2	0.15	14.2
10.0	0.16	13.0
11.0	0.18	11.6
12.0	0.20	10.4
12.3	0.20	10.4
13.0	0.21	9.8
14.0	0.23	9.1
15.4	0.25	8.3
15.5	0.25	8.3
18.3	0.30	6.9
18.5	0.30	6.9
20.0	0.33	6.4
20.3	0.33	6.4
24.5	0.40	5.2

## POST HARVEST APPLICATIONS

Crop	Target Diseases	Use Rate	Application Instructions								
<b>Bananas Plantains</b>  <b>Post-Harvest</b>	Crown Rot/Crown Mold ( <i>Colletotrichum musae</i> , <i>Fusarium pallidoroseum</i> , <i>Acremonium</i> spp., <i>Ceratocystis paradoxa</i> , <i>Glomerella cingulata</i> , <i>Penicillium</i> spp.)	200-400 ppm solution	<p>Apply as a single application of a 200-400 ppm solution to achieve good coverage. The application may be made as a spray, dip or may be painted onto the cut ends of the bananas. Application of the 200 ppm rate is appropriate for short distance transportation (e.g., within the USA). When a longer time in transport is expected (export), use the 300-400 ppm rate. If alum (1% w/v) is added to the spray solution, stir the suspension frequently as sedimentation and flocculation may occur. Addition of a non-ionic surfactant (0.10% v/v) may improve the compatibility of this mixture.</p> <p><b>Amount of Tetraban Fungicide to Mix 100 Gallons for Post-Harvest Banana Applications</b></p> <table><tr><th>Tetraban Fungicide Use Rate</th><th>100 gal Spray Solution</th></tr><tr><td>200 ppm</td><td>11 fl oz</td></tr><tr><td>300 ppm</td><td>15 fl oz</td></tr><tr><td>400 ppm</td><td>21 fl oz</td></tr></table>	Tetraban Fungicide Use Rate	100 gal Spray Solution	200 ppm	11 fl oz	300 ppm	15 fl oz	400 ppm	21 fl oz
Tetraban Fungicide Use Rate	100 gal Spray Solution										
200 ppm	11 fl oz										
300 ppm	15 fl oz										
400 ppm	21 fl oz										

### Specific Use Restrictions:

- 1) Do not make more than one application to bananas as post-harvest treatment.
- 2) This product may be degraded by exposure to direct sunlight. Do not store treated fruit in direct sunlight.

Crop	Target Diseases	Use Rate	Application Instructions
<b>Citrus Fruit</b> <b>Crop Group 10-10</b>  <b>Post-Harvest</b>  Calamondin Citron Citrus Hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma Mandarin Tangerine Uniq Fruit Hybrid  Including all cultivars and/or hybrids of these  See complete list of citrus fruit crops below.	Penicillium Decays Green Mold, Whisker Mold, Suppression of Blue Mold ( <i>Penicillium</i> spp.) Diplodia Stem-End Rot ( <i>Diplodia natalensis</i> ) Phomopsis Stem-End Rot ( <i>Phomopsis citrii</i> )	See Application Instructions	Use this product as a dip, drench, flood, or spray for the control of certain post-harvest diseases.  <b>For high volume (dilute) applications:</b> Mix 32-64 fl oz of this product in 25-100 gallons of an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated. Use T-Jet, flooders, or similar application systems.  <b>For low volume (concentrate) applications:</b> Mix 32-64 fl oz of this product in 7-25 gallons of water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion for the crop being treated. Apply to 250,000 lb of fruit. Use a controlled droplet type of applicator or similar system.  <b>For dip applications:</b> Mix 32-64 fl oz of this product in 100 gallons of water, wax/oil emulsion, or aqueous dilution of wax/oil emulsion. Dip for approximately 30 seconds and allow fruit to drain. For maximum decay control, treat citrus fruit once before storage and once after storage, just prior to marketing.

**Complete List of Citrus Fruit Crops:** Australian Desert Lime (*Eremocitrus glauca*); Australian Finger Lime (*Microcitrus australasica*); Australian Round Lime (*Microcitrus australis*); Brown River Finger Lime (*Microcitrus papuana*); Calamondin (*Citrofortunella microcarpa*); Citron (*Citrus medica*); Citrus Hybrids, *Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.; Grapefruit (*Citrus paradise*); Japanese Summer Grapefruit (*Citrus natsudaidai*); Kumquat (*Fortunella* spp.); Lemon (*Citrus limon*); Lime (*Citrus aurantiifolia*); Mediterranean Mandarin (*Citrus deliciosa*); Mount White Lime (*Microcitrus garrowayae*); New Guinea Wild Lime (*Microcitrus warburgiana*); Orange, Sour (*Citrus aurantium*); Orange, Sweet (*Citrus sinensis*); Pummelo (*Citrus maxima*); Russell River Lime (*Microcitrus inodora*); Satsuma Mandarin (*Citrus unshiu*); Sweet Lime (*Citrus limetta*); Tachibana Orange (*Citrus tachibana*); Tahiti Lime (*Citrus latifolia*); Tangelo (*Citrus x tangelo*); Tangerine (Mandarin) (*Citrus reticulata*); Tangor (*Citrus nobilis*); Trifoliate Orange (*Poncirus trifoliata*); Uniq Fruit (*Citrus aurantium* Tangelo group); cultivars, varieties and/or hybrids of these.

**Specific Use Restrictions:**

- 1) Do not make more than two applications to citrus fruit as post-harvest treatments.
- 2) This product may be degraded by exposure to direct sunlight. Do not store treated fruit in direct sunlight.

**Tuberous and Corm Vegetable Subgroup 1C - Post harvest**

Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Canna, Edible; Cassava, Bitter and Sweet; Chayote (root); Chufa; Dasheen; Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam Bean; Yam, True.

Use this product as a post-harvest spray for the control of certain post-harvest rots caused by Silver Scurf (*Helminthosporium solani*), *Fusarium* species, Late Blight (*Phytophthora infestans*), and Pink Rot (*Phytophthora erythroseptica*).

Application Method	Disease	Rate (fl oz)	Application Instructions
In-Line Aqueous Spray Application	Silver Scurf Fusarium Dry Rot Late Blight Pink Rot	0.6 fl oz/ton of tubers	<ul style="list-style-type: none"><li>• Ensure proper coverage of the tubers. Tumble tubers as they are treated.</li><li>• Mix the fungicide solution in an appropriate amount of water for the crop being treated.</li><li>• Use T-jet, CDA, or similar application system.</li></ul>
<b>Specific Use Restrictions:</b> <ol style="list-style-type: none"><li>1) Do not use on seed potatoes or seed pieces.</li><li>2) Ensure the solution remains in suspension by using agitation.</li><li>3) Do not make more than one post-harvest application to the tubers.</li></ol>			

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

### **Pesticide Storage**

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

### **Pesticide Disposal**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### **Container Handling [less than or equal to 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

### **Container Handling [greater than 5 gallons]**

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by stated and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call  
CHEMTREC 1-800-424-9300**

### **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Tetran ® is a registered trademark of Winfield Solutions, LLC.

Callisto®, Callisto® Xtra, and Halex® are registered trademarks of a Syngenta Group Company

Acrobat® is a registered trademark of BASF Corporation.