



12% GAMMA BHC WETTABLE POWDER

INGREDIENTS

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride

12.0%

Other isomers of benzene hexachloride

13.0%

INERT INGREDIENTS:

75.0%

Total


100%

WARNING

Keep Out of Reach of Children

See Remainder of WARNING STATEMENT on Back Panel

EPA Reg. No. 1386-27



Comments

NET WEIGHT

4 POUNDS

FC-82-B870

PROD. 1004

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

ACCEPTED WITH COMMENTS

UNICO 12% Gamma B.H.C. Wettable Powder

DIRECTIONS

CABBAGE, BROCCOLI, AND CAULIFLOWER—To control aphids, thrips, cabbage worms, and loopers, in the seedbeds, use 4 pounds (0.48 lb. gamma isomer of BHC) in 100 gallons of water per acre ($\frac{3}{4}$ tablespoonful in 1 gallon per 100 square feet), and apply spray thoroughly when insects are first noticed. Repeat as necessary.

DO NOT APPLY AFTER THE SEEDLING STAGE.

TOMATO AND EGGPLANT—To control aphids and thrips, use 3 pounds (0.36 lb. gamma isomer of BHC) in 100 gallons of water per acre ($\frac{1}{2}$ tablespoonful in 1 gallon per 100 square feet), and apply spray thoroughly when insects are first observed. Repeat in 5 to 7 days if necessary.

DO NOT APPLY AFTER EDIBLE PARTS START TO FORM.

STRAWBERRIES—To control spittlebug, use 8 pounds (0.96 lb. gamma isomer of BHC) in 100 gallons of water per acre ($1\frac{1}{4}$ tablespoonfuls in 1 gallon per 100 square feet), and apply spray early in the spring about 1 week after the first small spittle masses appear on or between the expanding leaves. Repeat treatment in about 10 days if necessary.

DO NOT APPLY AFTER FRUITS START TO FORM.

PEACHES AND PLUMS—To control plum curculio, use 3 pounds per acre (0.36 lb. gamma isomer of BHC) in 100 gallons of water or equivalent in low-volume sprayers ($\frac{1}{2}$ tablespoonful in 1 gallon) and apply thoroughly to trees. Make first application when three-fourths of petals have fallen and 3 repeated applications at 5 to 7-day intervals according to infestation.

DO NOT APPLY TO PEACHES AND PLUMS WITHIN 60 DAYS OF HARVEST.

CAUTION

MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH THE SKIN. May produce irritation of skin and eyes. Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

Do not apply under conditions which might result in residual odor or taste remaining on poultry, feed and foodstuffs. Do not store near products which may absorb the odor. **DO NOT APPLY TO CROPS WHICH ARE TO BE FOLLOWED IN ROTATION BY ROOT CROPS OR PEANUTS. DO NOT USE ON ANIMALS.**

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.