



BHC EMULSIFIABLE CONCENTRATE

INGREDIENTS

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride	11%
Other isomers of benzene hexachloride and related compounds	13%
Aromatic petroleum derivative solvent	70%
INERT INGREDIENTS	6%
Total	100%

CONTAINS 1 POUND GAMMA ISOMER OF BHC PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel
USDA Reg. No. 1386-32

Net Volume
FC-24B-6-69

1 U.S. Gallon Liquid
PROD. 205

MANUFACTURED BY

CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. May produce irritation of the skin or eyes. Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

Do not apply under conditions which might result in residual odor or taste remaining on feed and foodstuffs or crops. Do not use, spill, pour, or store near heat or open flame, or any product which will absorb odors. Keep container closed when not in use. Do not use on animals of any kind.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of containers, rinse solutions, and unused chemical in a pit on non-crop land located well away from water supplies.

This product is poisonous to bees exposed to direct treatment or to residues in sprayed areas. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

ACCEPTED

Jul 28, 1969

UNDER FEDERAL INSECTICIDE
ACT AND FEDERAL TOXIC SUBSTANCE ACT
FOYCO 210101C POISON REGISTERED
UNDER NO 1326-32

UNICO BHC Emulsifiable

DIRECTIONS FOR USE

PINE TREES—To control Southern pine beetle, mix 1 part of this product with 9 parts of water and apply this mixture as a uniformly thorough spray to the bark of all living pine trees. Begin applications early in the spring (March or April) and repeat at 2-3 week intervals through September. Apply also to injured and felled pine trees.

To control turpentine beetles, mix 1 part of this product with 9 parts of water and apply this mixture as a uniformly thorough spray to the lower 12 feet of all living pine trees. Begin applications early in the spring (usually April and May) and repeat at 2-3 week intervals through September. Apply also to injured and felled pine trees, and to stumps.

DO NOT GRAZE LIVESTOCK IN TREATED AREAS. DO NOT FEED TREATED AREAS TO LIVESTOCK. DO NOT APPLY TO PINE TREES WHICH ARE IN CONTACT WITH ROOT CROPS OR PEANUTS. BIRDS AND OTHER WILDLIFE ARE NOT KILLED.

CAUTION

HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. May produce irritation of the skin or eyes. Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

Do not apply under conditions which might result in residual odor or taste remaining on feed and foodstuffs or crops. Do not use, spill, pour, or store near heat or open flame, or any product which will absorb odors. Keep container closed when not in use. Do not use on animals of any kind.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of containers, rinse solutions, and unused chemical in a pit on non-crop land located well away from water supplies.

This product is poisonous to bees exposed to direct treatment or to residues in sprayed areas. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land well away from water supplies.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

ACCEPTED
Jul 28, 1969
UNDER FEDERAL INSECTICIDE
ACT REGISTERED UNDER NO. 1326-32

UNICO BHC Emulsifiable Concentrate

DIRECTIONS FOR USE

PINE TREES—To control Southern pine beetle, mix 1 part of this product with 50 parts of water, and apply this mixture as a uniformly thorough spray to the bark area of infested and surrounding pine trees. Begin applications early in the spring (March or April) and repeat at approximately 30-day intervals through September. Apply also to injured and felled pine trees.

To control turpentine beetles, mix 1 part of this product with 14 parts of water, and apply this mixture as a uniformly thorough spray to the lower 12 feet of the trunks of pine trees. Begin applications early in the spring (usually April and May) and repeat within 3 to 4 weeks. Apply also to injured and felled pine trees, and to stumps.

DO NOT GRAZE LIVESTOCK IN TREATED AREAS. DO NOT FEED ANY FOLIAGE FROM TREATED AREAS TO LIVESTOCK. DO NOT APPLY TO PINE TREES WHICH ARE TO BE FOLLOWED IN A ROTATION WITH ROOT CROPS OR PEANUTS. BIRDS AND OTHER WILDLIFE IN TREATED AREAS MAY BE KILLED.



ACCEPTED
7/27/72

1386-80

UNICO Chlordane Spray Powder

DIRECTIONS FOR USE

To use as a spray, fill tank about one-quarter full with water, add the correct amount of powder, agitate or stir thoroughly to obtain a good suspension. Then add remainder of water to fill spray tank and continue agitation. Some agitation is required to maintain the suspension while spraying. Proper timing of applications and thorough coverage are essential for satisfactory results.

APPLICATION ON PLANTS

POTATOES—To control Colorado potato beetle, flea beetle, serpentine and leaf miner, use 4 level tablespoonfuls per gallon of water. Apply spray to foliage when insects first appear and repeat at 7 to 10-day intervals or as necessary. Do not feed culled potatoes to dairy animals. Use at rate of 4 pounds per acre.

CABBAGE, CAULIFLOWER, AND BROCCOLI To control cabbage looper, flea beetles, green stink bug and young Harlequin bug, use 4 level tablespoonfuls per gallon of water. Apply spray when insects first appear and repeat at 7 to 10-day intervals. **DO NOT APPLY AFTER EDIBLE PARTS START TO FORM.** Use at rate of 4-pound per acre.

CANTALOUPE AND CUCUMBER To control cucumber beetles, squash bug, and squash vine borer, use 4 level tablespoonfuls per gallon of water. Apply when insects first appear and repeat as necessary. **DO NOT APPLY AFTER EDIBLE PARTS START TO FORM.** Use at rate of 4-pounds per acre.

TOMATOES To control flea beetles, tomato hornworm, and serpentine leafminer, use 4 tablespoonfuls per gallon of water. Apply spray when insects first appear and repeat at 5 to 7-day intervals as necessary. **DO NOT APPLY AFTER FRUITS BEGIN TO FORM. DO NOT GRAZE OR FEED FORAGE FROM TREATED AREAS TO LIVESTOCK OR POULTRY.** Use at rate of 4-pounds per acre.

PREPLANTING APPLICATIONS TO SOIL

POTATOES, CORN, SWEET CORN, AND STRAWBERRIES To control wireworms (does not include Southern potato wireworm) and white grubs, use 8 level tablespoonfuls per gallon of water. Apply spray to prepared soil surface at the rate of 1 gallon per 1,000 square feet, just before seeding or transplanting, and immediately disc or rake into top 3 inches of soil.

PEAS, BEANS, AND SWEET CORN To control seed maggot, use 8 level tablespoonfuls per gallon of water. Apply spray to soil surface of prepared seed bed at rate of 1 gallon per 1,000 square feet, and immediately disc or rake into top 3 inches of soil.

APPLICATIONS TO LAWN AND ORNAMENTAL TURF

To control chiggers, ants, crickets, sod webworm, and ticks, use 8 level tablespoonfuls per gallon of water. Apply spray to the grass at rate of 1 gallon per 1,000 square feet. Repeat as necessary. For imported fire ant, break open hard surface of mound and pour in 2 to 3 cupfuls per mound; spray surrounding soil as for other ants. If ants survive, repeat in 2 weeks. Do not contaminate food or feed crops.

To control white grubs and Japanese beetle grubs, use 16 level tablespoonfuls per gallon of water. Apply spray to grass in early spring or late fall, at rate of 1 gallon per 1,000 square feet.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by adequate rainfall or thorough sprinkling, and the grass has completely dried. Do not feed or graze forage from treated areas to poultry or livestock. Do not use forage from treated areas for hay. Birds feeding over treated lawn or ornamental turf may be harmed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist or vapor. In case of contact with skin, wash with soap and water. Avoid contamination of food and feedstuffs.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of containers, rinse solutions, wastes, and unused chemical in a pit or non-crop area and irrigated well away from water supplies.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

CHLORDANE SPRAY POWDER

INGREDIENTS

ACTIVE INGREDIENTS:	
Technical chlordane*	40%
INERT INGREDIENTS	60%
Total	100%

*Equivalent to 24% octachloro-4,7-methanotetrahydroindane and 16% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN
See Remainder of CAUTION Statement on Side Panel.
EPA Reg. No. 1386-80

NET WEIGHT

1 POUND

H&G-33A-5-72


PROD. 2133

DISTRIBUTED BY

**UNITED CO-OPERATIVES INC.,
ALLIANCE, OHIO 44601**



50% CHLORDANE WETTABLE POWDER



INGREDIENTS

ACTIVE INGREDIENT:	
Technical chlordane*	50%
INERT INGREDIENTS	50%
<hr/>	
TOTAL	100%

*Equivalent to 30% octachloro-4,7-methanotetrahydroindane and 20% related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 1386-92

DIRECTIONS FOR USE

To use as a spray, fill tank about one-quarter full with water, add the correct amount of powder, agitate or stir thoroughly to obtain a good suspension. Then, add remainder of water to fill spray tank and continue agitation. Some agitation is required to maintain the suspension while spraying. Proper timing of applications and thorough coverage are essential for satisfactory results.

PREPLANTING APPLICATIONS TO SOIL

POTATOES, CORN, SWEET CORN, AND STRAWBERRIES — To control wireworms (does not include Southern potato wireworm) and white grubs, use 10 to 20 pounds per acre in the quantity of water required for uniform spray coverage of the soil (7 ounces of powder per 1,000 sq. ft.). Apply spray to prepared soil surface just before seeding or transplanting and immediately disc or rake into top 3 inches of soil. Properly applied soil treatments may be effective for more than one year. Repeat when new infestations develop.

PEAS, BEANS, AND SWEET CORN — To control wireworms and white grubs, use 10 to 20 pounds per acre in the quantity of water required for uniform spray coverage of the soil (7 ounces of powder per 1,000 sq. ft.). Apply spray to prepared soil surface just before seeding or transplanting and immediately disc or rake into top 3 inches of soil. Properly applied soil treatments may be effective for more than one year. Repeat when new infestations develop.