

[1386-35]

DILUTION CHART

Recommended Dilution	Use This Amount to Make		
	1 Gallon	1 Quart	5 Gallons
1 to 9	1 3/4 Cupfuls	7 Tablespoonfuls	2 Quarts
1 to 19	7 Fl. Oz.	3 1/2 Tablespoonfuls	1 Quart
1 to 49	3 Tablespoonfuls	1 Tablespoonful	1 3/4 Cupfuls

ORNAMENTALS

Crop	To Control	Ounce of Formulation Per 100 Gallons
African Violets	Aphids	10 to 30
Asiatic	Flea Beetles	
Azaleas	Leafhoppers	
Begonias		
Camellias		
Carnations		
Chrysanthemum		

Apply when insects first appear and repeat as required to maintain control. Spray both upper and lower leaf surfaces.

VEGETABLE CROPS

Crop	To Control	Ounce of Formulation Per Acre
Asparagus	Aphids, Leafhoppers, Mexican Bean Beetle, 12-Spotted Cucumber Beetle, White Fly	10 to 30 oz.
Beans	Aphids, Leafhoppers, Mexican Bean Beetle, 12-Spotted Cucumber Beetle, White Fly	10 to 30 oz.
Broccoli, Brussel Sprouts	Aphids, Cabbage Looper, Cross-striped Cabbage Worm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug	10 to 30 oz.
Cabbage, Cauliflower	Cabbage Looper, Green Peach Aphid, Leaf Tier	10 to 30 oz.
Celery	Fireworms, Leafhopper	10 to 30 oz.
Cranberries	Blistar Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid	10 to 30 oz.
Eggplants	Cabbage Looper, Diamondback Moth Larvae, Green Peach Aphid, Imported Cabbageworm	10 to 30 oz.
Lettuce	Green Peach Aphid	10 to 30 oz.
Peppers	Aphids, Colorado Potato Beetle, Flea Beetle, Leafhopper	10 to 30 oz.
Potatoes	Aphids, Flea Beetle	10 to 30 oz.
Radish	Aphids, Cabbage Looper, Webworm	10 to 30 oz.
Spinach	Colorado Potato Beetle, Flea Beetle	10 to 30 oz.
Tomatoes	Green Peach Aphid, Stink Bug	10 to 30 oz.

Use the above amounts in sufficient water for thorough coverage (usually 100 gallons).


CAUTION

Harmful if swallowed. Protect food, milk, and milking utensils from the spray. Do not use, pour, spill, or store near heat or open flame. Avoid contamination of feed and foodstuff.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE

Use only for the purpose and in compliance with the limitations, cautions, or warnings stated on this label.



INGREDIENTS:

ACTIVE INGREDIENTS:	
Pyrethrins	1.12%
Piperonyl butoxide, technical*	11.25%
Petroleum distillate	72.63%
INERT INGREDIENTS:	15.00%
Total	100.00%

*Equivalent to 9.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.25% related compounds

®Registered Trademark F.M.C. Corp.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1386-35

Net Volume 1 U.S. Quart Liquid

FC-35A-6-73

Manufactured By

PROD. 206

Unico Pyrenone Emulsifiable Concentrate

DIRECTIONS FOR USE

This product should be diluted or mixed with water for application as a spray. It also may be used undiluted in automatic or self-actuated cattle sprayers.

DAIRY CATTLE, BEEF CATTLE, HOGS, SHEEP AND HORSES—To control horn flies and mosquitoes, use 1 quart mixed in 19 quarts of water, and apply the diluted spray at rate of 1 1/2 to 2 quarts per large animal; usually 1 quart per animal is adequate for hogs and sheep. Repeat daily as needed.

To control stable flies and horse flies, use 1 quart mixed in 9 quarts of water, and apply the diluted spray at the rate of 1 1/2 to 2 quarts per animal. Applications to control horse flies during periods when they are numerous should be made at 1 to 3-day intervals; for stable flies, repeat as needed.

To control lice, use 1 quart mixed in 49 quarts of water (12 1/4 gallons), and apply thoroughly to each animal including the head and tail at the rate of 1 gallon of diluted spray per large animal. A second application in about 10 days is usually required, after which applications should be repeated as needed.

To control sheep ticks (keds), use 1 quart mixed in 49 quarts of water, and apply spray thoroughly to entire animal. Apply after shearing or with pressure to part the fleeces.

AUTOMATIC SPRAYERS—Use this product undiluted in automatic treaters or self-actuated sprayers to control horn flies and stable flies and to protect against horse flies on dairy cattle and beef cattle. The sprayer should be placed in an area or lane so the cattle will make 2 or 3 round trips daily through it. The nozzles should be calibrated to deliver about 1.15 fl. oz. each time the sprayer is actuated. The sprayer should be hooded with canvas or burlap to prevent wind dispersion of the spray mist.

DAIRY BARN, MILK ROOMS, AND POULTRY HOUSES—To control houseflies, use 1 quart mixed in 19 quarts of water, and apply as a mist spray at the rate of 2 fl. oz. per 1,000 cu. ft. Close doors and windows before spraying and leave them closed for 10 to 15 minutes after spraying. Do not apply in dairy barns during milking. Do not apply in milk rooms when milk is being processed or handled. Do not contaminate drinking water, milk, and milk handling equipment. Do not remain in treated area. Ventilate area after treatment is completed.


GRAIN BINS AND FEED STORAGE AREAS—To control exposed stages of granary weevil, rice weevil, grain borers, saw-toothed grain beetle, confused flour beetle, red flour beetle, and Angoumois grain moth, use 1 quart mixed in 9 quarts of water, and apply the spray at the rate of 1 gallon per 1,000 square feet of surface, paying particular attention to the areas where these insects crawl, rest, or hide. Application should be made to inside and outside of grain or feed storage locations, and should be repeated as needed. As a prerequisite, bins and storage areas should be cleaned thoroughly prior to spraying. Allow the sprayed surfaces to dry before introducing grain or feed.

SPECIAL NOTE

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break empty container into previously prepared pit on non-crop land in an isolated area, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush empty container under wheels of heavy vehicle or with an axe and bury crushed container deeply in a pit on non-crop land located well away from water supplies. Do not burn empty containers.

UNIVERSAL COOPERATIVES, INC.
ALLIANCE, OHIO 44601



1386-35