

# DISINFECTANT

**ACCEPTED**  
415  
SUBJECT TO ATTACHED COMMENTS

A.O.A.C. Phenol Coef. 4

ACTIVE INGREDIENTS

INERT INGREDIENTS

**FOR USE** in schools, offices and dormitories, for disinfecting hard surfaces, such as urinals, bowl surfaces, water closets, garbage collection areas and other places where sanitation is required.

**FOR DISINFECTING TOILET BOWLS:**

1. Pour 1/2 cup of disinfectant into the bowl.

**FOR DISINFECTING GARBAGE COLLECTION AREAS:**

1. Mix 1/2 cup of disinfectant with 1 gallon of water.

**FOR OTHER HARD SURFACE DISINFECTANT PURPOSES:**

1. Add 1/2 cup of disinfectant to 1 gallon of water.

**FOR CLEANING:**

1. Add 1/2 cup of disinfectant to 1 gallon of water.

**FOR SPRAYING TO DEODORIZE:**

1. Add 1/2 cup of disinfectant to 1 gallon of water.

**WARNING: KEEP OUT OF THE REACH OF CHILDREN.** This disinfectant is recommended for use on inanimate objects only, and is not to be used as an antiseptic or germicide on the human body. Rinse empty container with water and discard it. **HARMFUL IF SWALLOWED.** Do not get in eyes. May cause eye damage or skin irritation. Avoid breathing spray mist and contaminating food.

**FIRST AID:** If swallowed, DO NOT induce vomiting. Vomiting may be harmful. Call a physician immediately. If contact with eyes or skin occurs, flush thoroughly with water. Get medical attention for eyes.

**CAUTION:** Do not use, pour, spill, or store near heat or open flame.

**NET CONTENTS: 1 Gallon KEEP FROM FREEZING**

Disinfectant, Phenol Coef. 4  
EPA Reg. No. 1383-19  
EPA Est. No. 1383 MD 1

Camerman's Note

Poor Copy

E.P.A. Reg. No. 1383-19  
E.P.A. Est. No. 1383 MD 1

1973



# STOCK PREMISES DISINFECTANT

## INGREDIENTS

### ACTIVE INGREDIENTS:

Coal tar acids .....	11%
Coal tar neutral oils .....	62%
Soap .....	17%
INERT INGREDIENTS .....	10%
Total .....	100%

ACCEPTED

6/2/72

KEEP OUT OF REACH OF CHILDREN

## DANGER

See First Aid and Other Precautions on Side Panel

EPA Reg. No. 1386-44

Net Volume

5 U.S. Gallons Liquid

FC-100A-4-72

PROD. 502

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC., ALLIANCE, OHIO 44601

ACCEPTED

6/2/72

THE FEDERAL INSECTICIDE AND FUNGICIDE ACT OF 1947  
EPA REG. NO. 1386-44

## UNICO Stock Premises Disinfectant

### DIRECTIONS FOR USE

To apply as a spray, wash or dip, mix this disinfectant with warm water in the proportions stated below, and stir thoroughly. If the water is hard, dissolve flaked or chipped soap at rate of 1/3 to 1/2 ounce per gallon in the water before adding the disinfectant.

Do not use in milking stalls, milking parlors or milk houses.

Remove all animals, poultry, and feeds from premises, cars, boats, trucks, coops, and crates.

Remove all litter and manure from floors, walls, and surfaces of barn, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.

Empty all troughs, racks, and other feeding and watering appliances.

Saturate all surfaces with accepted disinfecting solution.

Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers.

Ventilate buildings, cars, boats, coops, and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried.

All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.

**BARN AND STABLES**—To disinfect premises occupied by livestock, use 1 part to 30 parts of warm water (e.g., 2 quarts in 15 gallons; 1 pint in 3 1/4 gallons) and after thorough cleaning of area, saturate walls, floors, partitions, and stanchions with the solution. Repeat as necessary. Do not apply directly to animals. Do not apply in dairy barns or in areas where milk is processed or stored.

**HOG PENS**—Apply as a spray to ceilings, walls, stanchions, etc. Remove animals before application and keep them out until spray has dried. Cover feeders and waterers.

**SANITARY CLOSETS**—Use 4 fluid ounces to each gallon of water in the pail to aid in disposal of fecal material in sanitary closets. Repeat daily each time pail is emptied.

**SINKS, HOPPERS, AND URINALS**—Use 8 tablespoonfuls in each gallon of water for daily cleaning.

**SEED POTATO STORAGE BINS**—To disinfect empty bins, storage areas, and handling equipment for seed potatoes only, use 1 part to 30 parts of warm water (e.g., 1 quart in 7 1/2 gallons). Remove and destroy all diseased stock and after thorough cleaning, apply as a wash or as a wet spray to the bins, floors, and walls of storage areas and equipment used for handling and storage of seed potatoes.



# STOCK PREMISES DISINFECTANT

## INGREDIENTS

### ACTIVE INGREDIENTS:

Coal tar acids  
Coal tar neutral oils  
Soap

### INERT INGREDIENTS

Total

11%  
62%  
17%  
10%  
100%

## DANGER

KEEP OUT OF REACH OF CHILDREN

See First Aid and Other Precautions on Side Panel

EPA Reg. No. 1386-44

Net Volume

5 U.S. Gallons Liquid

EC 100A-4-72

PROD 502

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC., ALLIANCE, OHIO 44601

1/2/72

## UNICO Stock Premises Disinfectant

### DIRECTIONS FOR USE

To apply as a spray, wash or dip, mix this disinfectant with warm water in the proportions stated below, and stir thoroughly. If the water is hard, dissolve flaked or chipped soap at rate of 1 3 to 1 2 ounce per gallon in the water before adding the disinfectant.

Do not use in milking stalls, milking parlors or milk houses.

Remove all animals, poultry, and feeds from premises, cars, boats, trucks, coops and crates.

Remove all litter and manure from floors, walls, and surfaces of barn, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.

Empty all troughs, racks, and other feeding and watering appliances.

Saturate all surfaces with accepted disinfecting solution.

Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers.

Ventilate buildings, cars, boats, coops, and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed, set or dried.

All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.


**BARN AND STABLES**—To disinfect premises occupied by livestock, use 1 part to 30 parts of warm water (e.g., 2 quarts in 15 gallons; 1 pint in 3 1/4 gallons) and after thorough cleaning of area, saturate walls, floors, partitions, and stanchions with the solution. Repeat as necessary. Do not apply directly to animals. Do not apply in dairy barns or in areas where milk is processed or stored.

**HOG PENS**—Apply as a spray to ceilings, walls, stanchions, etc. Remove animals before application and keep them out until spray has dried. Cover feeders and waterers.

**SANITARY CLOSETS**—Use 4 fluid ounces to each gallon of water in the pail to aid in disposal of fecal material in sanitary closets. Repeat daily each time pail is emptied.

**SINKS, HOPPERS, AND URINALS** Use 8 tablespoonfuls in each gallon of water for daily cleaning.

**SEED POTATO STORAGE BINS**—To disinfect empty bins, storage areas, and handling equipment for seed potatoes only, use 1 part to 30 parts of warm water (e.g., 1 quart in 7 1/2 gallons). Remove and destroy all diseased stock and after thorough cleaning, apply as a wash or as a wet spray to the bins, floors, and walls of storage areas and equipment used for handling and storage of seed potatoes.



Repeat when bin is emptied and prior to refilling seed potatoes. Handling equipment for seed potatoes should be treated at regular intervals. Planters should be disinfected at least twice a day and between each different seed lot. Crates for seed potatoes may be disinfected by dipping in a tank of the solution. Soak used burlap bags to be used for seed potatoes in the solution for 1 hour (250 bags in 40 gallons of solution). Change the solution when dirt begins to accumulate. Do not use handling equipment, bins, crates, or bags treated until thoroughly dry and aired. Do not apply directly to seed potatoes or to bins when seed potatoes are present. DO NOT APPLY TO OR IN BINS, STORAGE AREAS, HANDLING EQUIPMENT, AND CRATES FOR EDIBLE POTATOES OR POTATOES TO BE MARKETED OR USED FOR PURPOSES OTHER THAN AS SEED FOR PLANTING.

### DANGER

**DANGER:** Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

This product is poisonous to aquatic animal life. Keep out of any body of water. Apply only for the purposes and in the manner directed on this label. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water supplies.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land in an isolated area and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land in an isolated area.

### NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

