DO NOT APPLY TO FEED OR FORAGE TO BE FED TO OR GRAZED BY LIVESTOCK OR POULTRY. DO NOT CONTAMINATE FEED OR FOOD CROPS.

SMALL QUANTITIES-Use following equivalents to mix and apply small quantities:

Amount per 100 gallons of water 1½ Pints

2 Pints 236 Pints

•

iounț	to	use	in	1	gallon	of	water
	1%	Te	asp	00	nful		
	24	Te	asp Sp	00	nful nful		

WARNING

MAY BE FATAL IF SWALLOWED! MAY BE ABSORBED THROUGH SKIN! Do not breathe soray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Do not contaminate feed and foodstuffs. Not for use or storage in or around the home. Do not use on cats, dogs or other household pets or humans. Do not use or store near heat or open flame. Not for use on poultry or in poultry houses.

This product is highly toxic to bees exposed to direct treatment or to residues on crops or foliage. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff as a result of heavy rain may occur. Clean application equipment and dispose of containers, rinse solutions, and unused chemical in a pit located on non-crop land well away from watesupplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Break empty glass container into a previously prepared pit on non-crop land and bury fragments deeply.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.



ACTIVE INGREDIENTS:

Li	indane	(Gamma	isomer	of	benzene	hexa
X	ylene	•				
INERT	INGR	EDIENTS .				

achloride) 12.7% 3.1% 100% Total

CONTAINS 1 POUND LINDANE PER GALLON

WARNING

KEEP OUT OF REACH OF CHILDREN See Remainder of WARNING Statement on Side Panel. USDA Reg. No. 1386-45

Net Volume **1 U.S. Quart Liquid**

FC-129B-11-70

MANUFACTURED BY

UNITED CO-OPERATIV ALLIANCE, OHIO 44

UNICO Lindane Emulsifiable Concentrate

DIRECTIONS FOR USE

EMULSIFIABLE CONCENTRATE

areas may be killed. PROD. 209

VES, INC.			
4601	;		

BEEF CATTLE, HOGS, SHEEP, AND GOATS—To control psoroptic and chorioptic mange mite (scabies) on beef cattle, sarcoptic mange miter on hogs, and psoroptic mites (common scabies) and chorioptic mites (foot scabies) on sheep, use 4 pints in 100 gallons of water (0.06 per cent of lindane) to spray each animal thoroughly. For animals over 3 months (calves 5 months) of age to 1 year old, use 2 pints per 100 gallons of water (0.03% lindane). Complete coverage (including inside of ears) either by spraying or dipping is essential. Treatment should be repeated in 7 to 10 days if necessary.

To control lice and ticks on beef cattle, sheep, and goats. and lice on hogs, use 2½ to 3½ pints in 100 gallons of water (0.037 to 0.05 percent of lindane) to spray each animal thoroughly. For animals over 3 months (calves 5 months) of age to 1 year old, use 2 pints per 100 gallons of water (0.03% lindane). Repeat treatment if necessary in about 10 days.

If used as a spray, apply the sprays to the kinds of animals listed with 12.7% a sprayer which develops adequate pressure. Generally, large beef cattle 84.2% require about 4 gallons of spray per head, and smaller animals, hogs, sheep 3.1% and goats require proportionally less.

DO NOT APPLY TO DAIRY CATTLE OR DAIRY GOATS. DO NOT APPLY SPRAY TO BEEF CATTLE, HOGS, SHEEP, AND GOATS WITHIN 30 DAYS OF SLAUGHTER; DO NOT DIP THESE ANIMALS WITHIN 60 DAYS OF SLAUGHTER. DO NOT APPLY TO THESE ANIMALS WITHIN 30 DAYS OF GIVING BIRTH TO YOUNG NOR FOR AT LEAST 3 WEEKS THERE-AFTER. DO NOT TREAT NURSING ANIMALS. DO NOT APPLY TO PIGS. SHEEP AND GOATS LESS THAN 3 MONTHS OLD. DO NOT APPLY TO CALVES LESS THAN 5 MONTHS OLD. DO NOT TREAT ANIMALS IN COLD, STORMY WEATHER. DO NOT TREAT OVERHEATED OR SICK ANIMALS.

VEGETATION IN NON-CROP AREAS—To control ticks and chiggers on vegetation (grass and low-growing brush) in areas where no crop is growing, use 4 pints in 100 gallons of water per acre to spray the vegetation thor-oughly. Repeat treatment as necessary. Birds and other wildlife in treated