



## UNICO Superior Miscible Spray Oil

### DIRECTIONS FOR USE

**DECIDUOUS FRUIT TREES** During the dormant or delayed dormant period to control scale insects and eggs of European red mite, clover mite, apple rust mite and fall cankerworm on apples, pears, cherries, peaches and plums, use 2 gallons per 100 gallons of water and apply spray thoroughly to twigs, branches, and trunks early in the spring on bright clear days when the temperature is above 45 F., during the time from just before growth starts until the young leaves are not more than 1/2 inch in length.

During the summer when foliage is present use 2 gallons per 100 gallons of water and apply spray thoroughly to foliage and branches to control scale insects, two spotted nut and red bug and trips. Apply when insects first appear, preferably before they cause noticeable damage and repeat at 30-day intervals, if necessary, until not more than 3 applications have been made during the growing season.

**ORNAMENTAL SHADE TREES AND HARDY SHRUBS** During the dormant or delayed dormant period to control scale insects and eggs of European red mites on ornamental deciduous shade trees and hardy ornamental shrubs, use 2 gallons per 100 gallons of water and apply spray thoroughly to twigs, branches, and trunks, early in the spring, on bright clear days when the temperature is above 45 F., just before growth starts.

During the summer period when foliage is present use 2 gallons per 100 gallons of water and apply spray thoroughly to foliage and branches to control scale insects, two spotted nut, spinybug, fruit and two spotted mites. Apply when insects first appear, preferably before they cause noticeable damage and repeat at 30-day intervals, if necessary, until not more than 3 applications have been made during the growing season.

# SUPERIOR MISCIBLE SPRAY OIL

## 60 SECONDS VISCOSITY For Dormant and Foliage Applications

### INGREDIENTS

#### ACTIVE INGREDIENTS:

Refined petroleum distillate

99%

#### INERT INGREDIENTS

1%

Total

100%

Unulfonated residue  
Class or Grade  
Viscosity at 100 F

93% Minimum  
Unclassified  
60% Seconds

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel.

EPA Reg. 1386-53

**Net Volume**

**U.S. Gallons Liquid**

FC-92A 5-72

PROD. 303

MANUFACTURED BY

**UNITED CO-OPERATIVES, INC., ALLIANCE, OHIO 44601**



# SUPERIOR MISCIBLE SPRAY OIL

## 60 SECONDS VISCOSITY

### For Dormant and Foliage Applications

#### INGREDIENTS

##### ACTIVE INGREDIENTS:

Refined petroleum distillate

99%

##### INERT INGREDIENTS

1%

Total

100%

Un sulfonated residue  
Class or Grade  
Viscosity at 100 F

93% Minimum  
Unclassified  
60% Seconds

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel.

EPA Reg. 1386-53

Net Volume

U.S. Gallons Liquid

FC-92A-5-72

PROD. 303

MANUFACTURED BY

DO NOT FEED TREATED FORAGE TO DAIRY ANIMALS. DO NOT ENSILE TREATED CORN. DO NOT FEED STOVER FROM SWEET CORN RECEIVING 3 APPLICATIONS. DO NOT APPLY TO FIELD CORN.

TO MIX—Fill spray tank about one-third full with water, pour in the required amount of Miscible Spray Oil, stir or agitate thoroughly and fill remainder of tank with water. Some agitation is required to maintain the emulsion while spraying.

The dosage rate of 2 gallons per 100 gallons of water is equivalent to 1/2 pint per 3 gallons, or 5 1/2 tablespoonsful per gallon.

## CAUTION

Harmful if swallowed. Avoid breathing of spray mist. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of containers, rinse solutions, wastes and unused mixture in a pit or non-crop land located well away from water supplies.

Do not apply in freezing weather or when temperature is likely to drop to near freezing soon after spraying.

Do not apply to broadleaved evergreens when near freezing temperatures are expected within 3 weeks. Do not apply during or immediately before, but dry weather. Do not apply to plants suffering from drought. Do not apply within 30 days before or after application of sulphur.

## NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.