

4-14-72
1320-54



NOT APPLY TO CALVES LESS THAN 5 MONTHS OLD.
DO NOT TREAT ANIMALS IN COLD, STORMY WEATHER.
DO NOT TREAT OVERHEATED OR SICK ANIMALS.

SMALL QUANTITIES: Use following equivalents for mixing and applying small quantities:

Amount in 100 Gallons of Water	Amount to Use in 1 Gallon of Water
1 Pound	2 Teaspoonfuls
1½ Pound	3 Teaspoonfuls
2 Pounds	4 Teaspoonfuls

WARNING

MAY BE FATAL IF SWALLOWED! MAY BE ABSORBED THROUGH SKIN! Do not breathe dust or spray mist. Avoid contact with skin and eyes. In case of contact, wash immediately with soap and water; for eyes, flush with water and get medical attention. Wash thoroughly after handling. Do not contaminate feed and food-stuffs.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply in any manner not specified in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff as a result of heavy rain may occur. Clean application equipment and dispose of wastes, and unused chemical in a pit on non-crop land located well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.



INGREDIENTS

ACTIVE INGREDIENTS:	
Lindane (Gamma isomer of benzene hexachloride)	25%
INERT INGREDIENTS	75%
Total	100%

WARNING

KEEP OUT OF REACH OF CHILDREN

See Remainder of **WARNING** Statement on Side Panel.
EPA Reg. No. 1386-54

NET WEIGHT

1 POUND

FC-164C-2-72

PROD. 1009

DISTRIBUTED BY

UNICO Lindane Wettable Powder

DIRECTIONS FOR USE

BEEF CATTLE, HOGS, SHEEP AND GOATS—To control lice and ticks, use 1 pound in 100 gallons of water (0.03 percent of lindane) to spray each animal thoroughly. Repeat treatment if necessary in about 10 days.

To control psoroptic and chorioptic mange mites (scabies) on beef cattle, sarcoptic mange mites on hogs, and psoroptic mites (common scabies) and chorioptic mites (foot scabies) on sheep, use 1½ pounds in 100 gallons of water (0.047 percent of lindane) to spray each animal thoroughly. Complete coverage is essential.

Treatment should be repeated in 7 to 10 days. For animals more than 3 months (calves 5 months), but less than 1 year old, the spray must be limited to 0.03% lindane (1 lb. in 100 gal. of water).

Apply the sprays to the kinds of animals listed with a sprayer which develops adequate pressure, preferably 200 to 300 p.s.i. Generally, large beef cattle require about 4 gallons of spray per head, and hogs, sheep, and goats require proportionally less spray.

For sheep and goats, a dip made by using ¾ pound in gallons of water, into which each animal is completely submerged, may be preferable to spraying. Sheep should be treated in the fall usually, although treatment in the spring after shearing may be advisable.

DO NOT APPLY TO DAIRY CATTLE OR DAIRY GOATS. DO NOT APPLY SPRAY TO BEEF CATTLE, HOGS, SHEEP, AND GOATS WITHIN 30 DAYS OF SLAUGHTER. DO NOT APPLY TO THESE ANIMALS WITHIN 30 DAYS OF GIVING BIRTH TO YOUNG NOR FOR AT LEAST 3 WEEKS THEREAFTER. DO NOT TREAT NURSING ANIMALS. DO NOT APPLY TO PIGS, SHEEP AND GOATS LESS THAN 3 MONTHS OLD. DO

**UNITED CO-OPERATIVES INC.,
ALLIANCE, OHIO 44601**