

2,4-D CERTIFICATION OF COMPLIANCE**EPA Letters Dated October 30, 1992 and January 14, 1993**

I, being an authorized representative of Universal Cooperatives, Inc., certify that all containers of Unico 2,4-D Lo-V Ester Weed Killer (1386-60) produced by October 23, 1993 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by April 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.



Ms. Nik Ramswick
Universal Cooperatives, Inc.
7801 Metro Parkway
P. O. Box 460
Minneapolis, Minnesota 55440
(612-854-0800)

March 30, 1993

2044

BEST AVAILABLE COPY

2,4-D Lo-V Ester Weed Killer

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
See Side Panel For Additional Precautionary Statements

ACTIVE INGREDIENT:

2-Ethylhexyl ester of 2,4-dichlorophenoxyacetic acid*	65.1%
INERT INGREDIENTS	34.9%
Total	100.0%

*Isomer Specific Acid Equivalent (43.20% or 3.76 lbs./gal.) by AOAC Method No. 6.D01-5.

Net Contents: 2 1/2 Gallons

EPA Reg. No. 1386-90
EPA Est. No. 1386-01-1



UNIVERSAL COOPERATIVES, INC., MINNEAPOLIS, MN 55440

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes, and clothing. When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. Persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron. For aerial applicators in enclosed cockpits and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower. Keep out of the reach of children.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Do not induce vomiting unless directed by a physician since aspiration hazard exists with this product.

If on Skin: Remove contaminated clothing and wash affected areas with soap and water. Do not reuse contaminated clothing until washed. Get medical attention if irritation persists.

If in Eyes: Flush with water for at least 15 minutes. Call a physician immediately.

If Inhaled: Remove victim to fresh air. Apply respiration if indicated.

For Medical Emergency Information call 1-800-228-6635, extension 138.

ENVIRONMENTAL HAZARDS

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

NONAQUATIC USES

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water intended for irrigation or domestic purposes. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target areas.

AQUATIC USES

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water except as specified on this label. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unpr. treated persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Read the above reentry statement and the precautionary statements to workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION: Area treated with 2,4-D Lo-V Ester Weed Killer (contains 2,4-D) on (date of application). Do not enter without appropriate protective clothing until sprays have dried. (Insert here Statement of Practical Treatment from label.)"

2,4-D Lo-V Ester Weed Killer will kill or control the following as well as many other noxious plants susceptible to 2,4-D:

- | | | | |
|-----------------------------|--------------------------|----------------------|------------------|
| *Ailanth | *Clove, Red | Lambquarters, Common | Sagebrush |
| Alligatorweed | Cocklebur, Common | Locoweed | Shepherdspurse |
| Arrowweed | Cofficeban | Medicweed | *Smartweed |
| Artichoke | Creeeping Jenny | Morningglory, Annual | Sowthistle |
| *Barnwee (Hedge, | Curly Indigo | Mustard | Stinkweed |
| Field, European) | Dandelion | Parrotleather | Sumac |
| *Bittercress, Smallflowered | *Dock | Pennycress, Field | Sunflower |
| Bowledr | Duckweed | Pennywort | Valerian |
| Buckhorn | Elderberry | *Pappergrass | Vetch, Hairy |
| Bull Thistle | Evening Primrose, Culted | Pigweed | Virginia Creeper |
| Bullnettle | *Goldenrod | Plantain | Waterhyacinth |
| Burush | *Ground Ivy | Poison Ivy | Waterlily |
| Burdock | Hemp | Poleweed | Waterprimrose |
| Bur Ragweed | *Hoary Cress | Poverlyweed | *Wild Garlic |
| Buttercup | Honeyucide | Puncturevine | Wild Lettuce |
| Catnip | Horsweed or Marestall | Purslane | Wild Onion |
| Chickweed | Indigo | Ragweed | Wild Radish |
| Chicory | Intemred | Rush | Willow |
| | Jimsonweed | *Russian Thistle | |

*These species may require repeat applications and/or use of the higher rate recommended on this product label even under ideal conditions for application.

2,4-D Lo-V Ester Weed Killer should be used as a water diluted spray, or may be applied in liquid nitrogen fertilizer (see below), for selective control of susceptible weeds growing in small grain crops, corn, grass seed crops and ornamental turf, and for non-selective control of certain weeds not in growing crops, such as roadsides, fence rows, and drainage ditch banks.

Do not use in or near a greenhouse. Crops contacted by 2,4-D Lo-V Ester Weed Killer sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.

Apply when the weeds are young and in a succulent, rapidly growing condition, since best results are obtained when soil moisture and temperature conditions are favorable for rapid growth of weed plants. Sprays applied when weeds have stopped growing rapidly, or when they are affected by a lack of moisture in the soil, are often not effective against many kinds of weeds. Spray perennial weeds after they are completely emerged, but before the bloom stage. Kill of weeds may not be evident for 2 to 3 weeks after spraying. Retreatment of areas infested with perennial weeds may be necessary.

Considerable caution must be exercised in using 2,4-D sprays to avoid injury to crops and desirable plants. Do not apply directly to vegetables, grapes, fruit trees, ornamentals, cotton, soybeans, tomatoes or other plants which are sensitive to 2,4-D and do not permit spray mist to drift onto them since even minute quantities cause severe injury.

during the growing or dormant periods. Excessive airspeeds may reduce spray coverage and cause drift. Do not apply in high air or ground surface temperatures. Avoid spraying in the immediate vicinity of water. Do not use on spraying grasses, or white clover, are usually damaged and under some conditions.

Aerial application should be used only when there is no rain or heavy dew. Many states have regulations concerning aerial application with hollow cone type insecticide or other nozzles that produce a fine spray. Aerial application may be reduced by: (1) applying in order to obtain coverage; (2) increasing the wind speed; (3) the pounds of pressure at the nozzle tip; (4) by using nozzles; (5) by not applying when wind is blowing toward the target area. Consult local regulatory authorities before making applications.

Users should note that herbicide treatment of public water supply systems will aid in securing a permit in your state.

PREPARATION OF SPRAY AND APPLICATION: Recommended 2,4-D Lo-V Ester Weed Killer should be added to water in the spray tank at 1 to 2 gallons per acre. A good mixture and continue some agitation during application to make up will depend upon the equipment to be used. Proper dosage should be applied in at least 15 gallons of water per acre. In a pressure sprayer, apply in 150 to 200 gallons of water per acre. In a backpack sprayer, apply in 5 to 10 gallons of water per acre. Always use the proper amount of water regardless of the quantity of water.

Do not use the spray equipment for other purposes unless specifically labeled for such use. **USE IN LIQUID NITROGEN FERTILIZER:** 2,4-D Lo-V Ester is compatible with some liquid nitrogen fertilizers. However, the compatibility of the mixture must be tested before combining in the spray tank.

JAR TEST

Amount of 2,4-D Lo-V Ester Rate/Acre	Level Teaspoon Volume of 28 Gall./Acre
1/2 Pint	1 1/4 Teaspoon
1 Pint	1/2 Teaspoon
2 Pints	1 Teaspoon
4 Pints	2 Teaspoons

The amount of herbicide to be tested, as indicated in the table above, should be added to 100 gallons of finished spray per acre. When using appropriate changes in the ingredients of the compatibility test.

In a quart jar add the appropriate amount of 2,4-D Lo-V Ester, to one pint of liquid nitrogen fertilizer. Cover the jar and shake for 5 minutes and again after 30 minutes.

If the mixture does not ball up or form lumps, sludge, gels, or other precipitates, the combination is compatible. If precipitates formed with agitation, the combination may be used, provided good mixing and application operations.

If incompatibility occurs, the use of a suitable compatibility agent is recommended. Add 1/4 teaspoon of a compatibility agent to the mixture. (The 1/4 teaspoon is equivalent to 2 pints of liquid nitrogen fertilizer.) If the mixture is still incompatible, DO NOT USE.

TANK MIXING SEQUENCE

If the 2,4-D Lo-V Ester/fertilizer mixture is compatible with the spray tank, fill the tank with half the amount of fertilizer to be used, and complete filling the tank with the fertilizer. Application in the spray tank during application.

If a compatibility agent must be used, add it to the spray tank before adding the fertilizer. Follow applicable recommendations and field application rate agent labeling, as well as the 2,4-D Lo-V Ester labeling.

CROPS

SMALL GRAIN CROPS (Wheat, Barley, Rye, Oats): See label for application. Do not apply during the seedling stage, late in the boot stage. Do not apply during the heading stage, late in the grain stage. To control large weeds that will interfere with harvest or to a treatment can be applied when the grain is in the dough stage.

Spring Planted Oats: Use 1/2 pint per acre in sufficient after the fully tillered stage, except during the boot to dough stage.

Fall Planted Oats: Apply 1/4 to 1 1/4 pints per acre after the boot stage. Some difficult weeds may require higher rates (3/4 to 1 1/4 pints per acre) but crop injury may result. Do not spray during or after grain harvest.

NOTE: Do not use on grain interplanted with legumes. Fields within 14 days after treatment. Do not feed treated corn. See table for recommended use rates.

Preplant (Field Corn): To control emerged broadleaf weeds prior to planting corn. Apply 7 to 14 days before planting. Do not use on corn planted in fields where soil moisture is inadequate for normal weed growth. Use high rate on cover crops.

Preemergence: Apply to soil after planting but before corn emerges or where soil moisture is inadequate for normal weed growth. Apply on loam soils and higher rate on clay soils. Use high rate on cover crops.

Postemergence: Best results are usually obtained when weeds are 2 to 4 inches tall. When corn is over 8 inches tall, use drop nozzles to apply spray as close to the soil as possible. Do not apply from heading to dough stage. If soil moisture content is high, use 1/2 pint per acre rate to control weeds.

Delay Cultivation for 8 to 10 days after application to reduce weed growth. Hybrid corn should be sprayed with 2,4-D at the recommended rate, or after corn is being grown to be tolerant to 2,4-D treatment.

4 of 4

Preharvest: After the hard dough or denting stage, apply 1 to 2 pints per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control fall weeds such as bindweed, cocklebur, dogbane, greenweed, ragweed, sunflower, velvetleaf, and vine that interfere with harvesting. Do not harrow or feed corn fields to livestock for 7 days following application. **SORGHUM (Milo):** See table for recommended use rates.

Apply when sorghum is 4 to 12 inches high with secondary roots well established. When crop is over 10 inches tall, use drop nozzles to keep spray off foliage as much as possible. Do not apply from flowering to dough stage. Do not use with all. Temporary crop injury may occur under conditions of high soil moisture and high air temperature. Hybrids should be sprayed only if the cross or line is known to be tolerant to 2,4-D at the recommended dosage or after experience has shown the particular crosses or lines being grown to be tolerant to 2,4-D treatment.

RECOMMENDED RATES OF 2,4-D LO-V ESTER WEED KILLER

Crop (See Detailed Instructions Above)	Dosage Per Acre*	
	Normal Rates (Usually Safe To Crop)	Higher Rates for Special Situations [†] (More Likely to Injure Crop)
SMALL GRAINS		
(Wheat, Barley, Rye): Annual Weeds	1/2 to 1 Pint	1 to 2 Pints
Perennial Weeds	1 Pint	1 1/4 to 2 pints
Preharvest	1 to 2 Pints	
OATS		
Spring	1/2 Pint	
Fall	1/4 to 3/4 Pint	3/4 to 1 1/4 Pints
Preharvest	1 to 2 Pints	
CORN		
Preplant (Field Corn)	1 to 2 Pints	
Preemergence	1 to 2 Quarts	
Postemergence	1/2 Pint	1/2 to 3/4 Pint
Preharvest	1 to 2 Pints	
SORGHUM (Milo)		
Preemergence	1/2 Pint	1/2 to 3/4 Pint

*The higher rates as recommended above may be necessary to control difficult weed problems such as under dry conditions in the Western states. They should not be used, however, unless possible crop injury is acceptable. Consult State Agricultural Experiment Station or Extension Service Weed Specialists for recommendations or suggestions to fit local conditions.

[†]If band treatment is used, base the dosage rate on the actual area sprayed.

FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS IN SOYBEANS
(Preplant Application Only)

2,4-D Lo-V Ester Weed Killer may be used for postemergence control of many susceptible annual and perennial broadleaf weeds. This product may be applied prior to planting soybeans to provide foliar burn-down control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. Make only preplant applications to emerged weeds prior to planting soybeans grown in reduced tillage production systems. Apply only according to instructions given below.

Do not use any tillage operations between herbicide application and planting of soybeans. **Mixing Instructions:** Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may be added to spray mixture to increase the herbicidal effectiveness of 2,4-D Lo-V Ester Weed Killer on certain weeds. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

Application Precautions: Apply using air or ground equipment in a spray volume sufficient to provide uniform coverage of weeds. Use 2 or more gallons of total spray volume per acre for aerial application and 10 or more gallons per acre for ground equipment.

APPLICATION TIMING AND USE RATES

PRODUCT	BROADCAST APPLICATION RATE	WHEN TO APPLY
		(Days Prior To Planting Soybeans)
2,4-D Lo-V Ester	1 Pint/Acre	Not Less Than 7 Days
	2 Pints/Acre	Not Less Than 30 Days

For best weed control results, application should be made when weeds are small, actively growing and free of stress caused by temperature extremes, moisture stress, diseases, or insect damage. The control of individual weed species may be variable. Consult your local county agent or State Agricultural Extension Specialist or Crop Consultant for advice.

Use Precautions and Restrictions:

Important Notice — Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of such injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present at the time of application. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

- Do not use on sandy soils with less than 1.0% organic matter.
- Do not make more than one application per season regardless of the application rate used.
- Do not apply when weather conditions such as atmospheric temperature inversion or when wind direction favors drift from the treated area to susceptible plants.
- Do not allow livestock grazing or harvest hay, forage or fodder from treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.
- In treated fields, plant soybean seed as deep as practical, but not less than 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is adequately covered.
- Do not apply 2,4-D Lo-V Ester Weed Killer as described unless you are prepared to accept the results of soybean injury, including possible stand loss and/or yield reduction.
- During the growing season following application, do not replant treated fields with crops other than those labeled for use with 2,4-D Lo-V Ester Weed Killer.

SUGARCANE: Use 2 pints per acre as a preemergence application before cane appears or 2 quarts per acre as a blanket spray after cane emerges and through July, to aid in the control of J and other seedlings and susceptible broadleaf weeds.

LAWN AND ORNAMENTAL TURF: Use 1 to 3 pints of 2,4-D Lo-V Ester Weed Killer in one-half water to give 50% coverage to one acre on established stands of perennial grasses. Do not apply to creeping grasses such as bent grass, etc. Spraying newly seeded turf should not be treated until after the second mowing and the turf is well established (6 to 8 weeks). Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall with fall application, reseed in the spring. Legumes are severely damaged or killed, therefore, do not reseed where the legumes are desired. Deep-rooted perennial weeds may require repeated treatments in the same season or in subsequent years. The maximum number of broadcast applications per treatment site is 2 per year.

GRASS SEED CROPS: Apply 1 to 4 pints of 2,4-D Lo-V Ester Weed Killer in the spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five leaf stage, using 3/4 to 1 pint per acre to control small seedling weeds. After the grass is well established, higher rates of 1 to 4 pints can be used to control hard to kill annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on turf unless injury can be tolerated. Do not graze dairy animals nor cut forage for hay within 7 days of application.

FALLOW LAND: On established perennial species such as Canada thistle and field bindweed, apply up to 2 quarts per acre of 2,4-D Lo-V Ester Weed Killer for annual broadleaf weeds, apply 1 to 2 quarts per acre. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from the soil.

PASTURE AND RANGELAND: NOTE: Do not graze dairy animals on treated areas within 7 days after application. Do not harvest grass for hay within 30 days of application. Do not graze meat animals on treated areas within 3 days of slaughter. Do not use on bent grass, alfalfa, clover, or other legumes or on newly seeded pastures. Do not apply after heading begins or when grass is in the boot to milk stage where grass seed production is desired.

Bitterweed, Greenweed, Crotch, Dock, Kachla, Marshmallow, Mustard, and Other Broadleaf Weeds: Use 4 pints of 2,4-D Lo-V Ester Weed Killer per acre in the amount of water needed for uniform application. If the weeds are young and growing actively, 2 pints per acre will provide control of some species. Deep-rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

Wild Garlic and Wild Onion: Apply 4 pints per acre, making three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring.

Weed Control in Newly Sprigged Coastal Bermudagrass: Apply 2 to 4 pints per acre preemergence and/or postemergence. **Sand Shrewy Oak and Sand Sagebrush:** On the oak, use 2 pints in 5 gallons of oil or in 4 gallons of water plus 1 gallon of oil per acre. Apply by aircraft between May 15 and June 15. On the sagebrush, use 2 pints in 3 gallons of oil per acre and apply by aircraft when foliage is fully expanded and the brush is actively growing.

Big Sagebrush and Rabbitbrush: Use 4 pints per acre in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. Brush should be leafed out and growing actively when treated. Retreatment may be needed.

Ceanothus, Manzanita, Buckbrush, Coastal Sage, Coyotebrush, and Certain Silver Cholla Species: Use 4 pints per acre in 5 to 10 gallons of water. One gallon of fuel oil may be included in the spray mixture for added effectiveness. Make applications by aircraft or ground equipment to obtain uniform spray coverage. For effective control, the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed.

General Weed Control (Airsifts, Needles, Vacant Lots, Drainage Ditch Banks, Fence Rows, Industrial Sites, and Similar Areas): Use 1 to 2 quarts per acre. Usually 2 quarts per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 30 days or until 2,4-D has disappeared from soil. The maximum number of broadcast applications per treatment site is 2 per year.

Woody Plant Control: To control woody plants susceptible to 2,4-D, such as Alder, Buckhorn, Elderberry, Sumac, and Willow on non-crop areas, use 2 quarts in 100 gallons of water. Wet all parts of the plants thoroughly, including stem and foliage, to the point of run off. Higher volumes of up to 400 gallons per acre are necessary where the brush is very dense, and over 6 to 8 feet high. Applications are more effective when made on actively growing plants. Treatment should not be made during time of severe drought or in early fall when leaves lose their green color. Hard to control species may require retreatment next season.

USES IN FOREST MANAGEMENT

Ceanothus Release: For control of alder, apply 1 1/2 to 2 quarts of product per acre in 8 to 25 gallons of water, and apply as a foliage spray between mid-May and mid-June.

For control of madrone, manzanita, oak, tan oak, and similar species to release hemlock, spruce, and fir, apply 2 quarts of product per acre in 8 to 25 gallons of water, just prior to or during budbreak of Douglas fir.

After northern conifers, jack pine, red pine, black spruce, and white spruce cease growth and "harden off" in late summer, a spray of 1 1/2 to 2 quarts in 8 to 25 gallons of water per acre may be applied by air to control certain competing hardwood species such as alder, aspen, birch, hazel, and willow. Since this treatment may cause occasional conifer injury, do not use if such injury cannot be tolerated. Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

For control of hazel brush and similar species in the Lake states area, apply 2 quarts of product per acre in 8 to 25 gallons of water, when new shoot growth of hazel is complete.

Site Preparation:
(As Budbreak Spray)—For control of alder prior to planting seedlings, apply 2 quarts of product per acre in 8 to 25 gallons of water, after alder budbreak but before foliage is 1/4 full size.

(As Foliage Spray)—For control of alder prior to planting seedlings apply 2 quarts of product per acre in 8 to 25 gallons of water, after most alder leaves are full size.

Aquatic Applications:
For Aquatic Weeds in Lakes, Ponds, Drainage Ditches, and Marshes: Use 2 1/2 to 4 pints of product in 50 to 100 gallons of water per acre. Spray to wet foliage thoroughly. Application should be made when leaves are fully developed above waterline and plants are actively growing. Your State Conservation Department or Game and Fish Commission will assist you in determining the best time and rate for application under local conditions.

DO NOT APPLY to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen content of water and kill fish.

Perennial and other hard to control weeds may require a repeat application to give adequate control.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

1050193

CO-OP® IS A REGISTERED TRADEMARK OF UNIVERSAL COOPERATIVES, INC.

BEST AVAILABLE COPY