LOW VOLATILE ESTER 600

Directions for Use

GENERAL INFORMATION [illegible]

[illegible] **See Caution Statement.**

[illegible]

CORN—Pre-Emergence: [illegible]

CORN—Post-Emergence: [illegible]

Early Spraying: [illegible]

Lay-By Spraying: [illegible]

High Clearance Spraying: [illegible]

SORGHUM—Post-Emergence Only: [illegible]

HARD TO KILL WEEDS in NON-CROP LAND [illegible]

NOTICE

[illegible]

Distributed By
FARMERS REGIONAL COOPERATIVE
Fort Dodge, Iowa

NET CONTENTS 1 GALLON

ACCEPTED
MAY 23 1968
UNDER [illegible] DE
FUNGI[illegible] CT
FOR ECO[illegible] ED
UNDER NO. 1586 11
Comment

FELCO

LOW VOLATILE ESTER 600

CONTAINS 6 POUNDS 2,4-D ACID EQUIVALENT PER GALLON

2,4-D WEED KILLER

EMULSIFIABLE — USE IN AERIAL OR GROUND SPRAYERS

INGREDIENT STATEMENT

ACTIVE INGREDIENTS:	
Isooctyl ester of 2,4-Dichlorophenoxyacetic acid	94.62%
INERT INGREDIENTS:	5.38%
Total	100.00

*Equivalent to 2,4-Dichlorophenoxyacetic acid 62.82%

CAUTION: KEEP OUT OF THE REACH OF CHILDREN
SEE RIGHT PANEL FOR ADDITIONAL WARNINGS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

LOW VOLATILE ESTER 600

Directions for Use

GENERAL INFORMATION ON APPLICATION

Crop		Amount Per Acre	App. Acres 1 Gallon Will Cover
CORN	Pre-Emergence	[illegible]	[illegible]
	Post-Emergence	[illegible]	[illegible]
SMALL GRAIN	Spring Wheat, Oats, Barley, Winter Wheat	[illegible]	[illegible]
	Rye	[illegible]	[illegible]
PASTURES AND TURF		[illegible] pts.	6
NON-CROP AREAS AND SPOT TREATMENT		[illegible] to 2 pts. in 20 gals. water	[illegible] to 6

SUSCEPTIBLE WEEDS

Yellowbeed	Lambsquarter	Sunflower
Buckhorn	Mustard	Wild Carrot
Cocklebur	Pigweed	Wild Radish
Dandelion	Morning Glory	Wild Sweet Potato
Dock	Plantain	
Knotweed	Ragweed	

WEEDS MORE DIFFICULT TO KILL
(Use higher dosage rate)

Bitterweed	Horse Nettle	Sheep Sorrel
Buckbrush	Locoweed	Hoary Cres.
Buckwheat	Milkweed	Willow
Canada Thistle	Poison Ivy	Wild Onion

SMALL GRAIN Wheat, Barley, Rye, Oats: For weed [illegible] **Winter Wheat** and **Rye**, treat only in the spring. Use [illegible] to [illegible] pt. in sufficient water to cover one acre. Apply from the tillered to the jointing stage. In **Spring Wheat, Oats,** and **Barley,** use ½ to [illegible] pt. per acre. Use higher dosage for hard to kill weeds. Treat when crop is fully tillered (4 to 8 inches high) up until early boot stage. Do not spray during seeding stage or boot to dough stage. Do not graze or grade treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.

PASTURES AND TURF: Use [illegible] pt. in 20 gallons water per acre. Apply early in summer when plant are actively growing. Treat again in bud stage before weeds go to seed. For lawns use [illegible] tablespoon per gallon to [illegible] sq. ft. Repeat as needed for control. Do not spray until after first cutting. Injury may occur to bentgrass, St. Augustine grass, Dichondra, carpetgrass and cover.

WOODY PLANTS: For plants such as Sumac, Willow, Buckbrush in fence rows, pastures and ditches, use [illegible] pts. in [illegible] gallons water with [illegible] of spreader-sticker. Wet parts thoroughly. Apply after [illegible] and treat the foliage [illegible] for best results.

CAUTION

CAUTION: DON'T OVER DOSE! [illegible]

[illegible]

[illegible]

[illegible]