

ACCEPTED

Aug. 26, 1969

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 1386-73



# ROSE and FLORAL Spray Powder

Sulphur may cause injury to foliage if applied during periods of high daytime temperature. Do not apply during periods when the daytime temperature is 90 degrees F. or above, or within two days of the time such temperatures may be expected.

### CAUTION

May be absorbed through skin. Harmful if inhaled or swallowed. May cause irritation of nose and throat. Do not breathe dust or spray mist. Do not take internally. Skin contact may be harmful. Avoid contact with skin and eyes. Wash thoroughly after handling and before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Avoid contamination of feed and foodstuffs.

For the protection of honeybees, avoid use, if possible, during periods when bees are visiting the plants or crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

This product is poisonous to fish and wildlife. Keep out of any body of water. Apply only in the manner and for the purposes specified in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty container (paper) in a hot fire, outdoors, in an isolated area. The smoke is dangerous. Stay well away from the smoke.

### NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.



### INGREDIENTS

|  |       |
|--|-------|
| ACTIVE INGREDIENTS:                                  |       |
| Carbaryl (1-naphthyl N-methylcarbamate) .....        | 12.5% |
| Lindane (gamma isomer of benzene hexachloride) ..... | 3.0%  |
| Zineb (zinc ethylene bisdithiocarbamate) .....       | 12.5% |
| Sulphur .....  | 25.0% |
| INERT INGREDIENTS .....                              |       |
| Total .....  |       |
| 100.0%   |       |

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

USDA Reg. No. 1386-73

NET WEIGHT

(H&G-111)

1 POUND

DISTRIBUTED BY

Prod. 2144

## UNICO ROSE and FLORAL SPRAY POWDER

### DIRECTIONS FOR USE

Mix 5 tablespoonfuls in 1 gallon of water (8 pounds in 100 gallons per acre). Apply as a spray to ROSES and to ANNUAL and PERENNIAL ORNAMENTAL FLOWERING PLANTS listed below to control the insects and plant diseases indicated. Begin applications before diseases are apparent and before insects become numerous, which is usually when new growth is just starting in the spring on roses and perennial flowering plants, or when seedlings of annual flowering plants appear above ground or young plants are transplanted into the garden. Spray the foliage evenly and thoroughly, and be sure to spray the underside of all leaves. Repeat applications at 7 to 10-day intervals and after heavy rain to maintain protection through the growing season.

Use on:

|                      |                  |                     |
|----------------------|------------------|---------------------|
| <u>Aster</u>         | <u>Dogwood</u>   | <u>Lilac</u>        |
| <u>Azalea</u>        | <u>Euonymus</u>  | <u>Peony</u>        |
| <u>Arborvitae</u>    | <u>Forsythia</u> | <u>Rhododendron</u> |
| <u>Boxwood</u>       | <u>Gardenia</u>  | <u>Rose</u>         |
| <u>Camellia</u>      | <u>Gladiolus</u> | <u>Snapdragon</u>   |
| <u>Chrysanthemum</u> | <u>Hemlock</u>   | <u>Spirea</u>       |
| <u>Dahlia</u>        | <u>Holly</u>     | <u>Yew</u>          |
| <u>Delphinium</u>    | <u>Iris</u>      | <u>Zinnia</u>       |

to control aphids, ants (attending aphids), certain caterpillars such as tent caterpillar, flea beetles, fleahoppers, Fuller's rose weevil, Japanese beetle adults, leafhoppers, lace bug, leaf miners, plant bugs, rose chafer, thrips (exposed), spittlebug, and white fly among the insects, and black spot, powdery mildew, Botrytis flower blight and certain other flower blights, leaf scorch, spot anthracnose, Didymellina leaf spot, Septoria leaf spot, and rust among the plant diseases.

For Botrytis flower blight and similar flower blights on azalea and camellia, apply liberally to flowers (while open) and to soil and litter around the plants. For Botrytis blight, spot anthracnose, and leaf spot on dogwood, apply at bud break and repeat as previously directed.

UNITED CO-OPERATIVES, INC.  
ALLIANCE, OHIO 44601