

ACCEPTED

UNICO

# FRUIT SPRAY Powder

**GRAPES**—Grape berry moth, Japanese beetle adults, red banded leaf roller, leafhopper, two spotted spider mite, black rot, downy mildew

Apply (1) when buds or shoots show 1/2 inch green, (2) when shoots are 10 to 12 inches long, (3) one week after bloom, and (4) thereafter at 7 to 10 day intervals for one or more applications as needed. **Do not apply within 7 days of harvest.**

**STRAWBERRIES** — Spittlebug, strawberry leaf roller, strawberry weevil, spider mites, Botrytis fruit rot, leaf spot

Apply (1) when new growth starts in the spring, (2) at petal fall, and (3) at subsequent 7 to 10-day intervals as needed. **Do not apply within 2 days of harvest.**

**RASPBERRIES**—Japanese beetle adults, leaf roller, leafhoppers, raspberry aphid, spider mites, anthracnose. Post harvest applications to raspberry canes may be necessary to combat anthracnose.

Apply (1) when new growth starts in the spring, (2) at petal fall, and (3) at subsequent 7 to 10-day intervals as needed. **Do not apply within 7 days of harvest.**

Do not graze or feed forage from treated areas to livestock.

### SPECIAL NOTES

Not effective if used with alkaline materials such as Bordeaux, lime, lime-sulfur, and casein-lime spreaders. Some injury may occur on tender foliage in the presence of rain or high humidity of several days duration following application. For protection of honey bees, avoid use, if possible, during periods when honey bees are visiting the crop; when necessary to use during such periods, warn beekeepers, well in advance, to locate hives at a safe distance until one week after application.

### CAUTION

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid breathing spray mist. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothes. Change to clean clothing daily. Avoid contamination of food and foodstuffs.


**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

This product is poisonous to fish. Keep out of any body of water. Apply only in the manner and for the purposes directed on this label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land well away from water sources.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. Burn empty container (papers) in a hot fire, outdoors, in an isolated area. The smoke may be hazardous. Stay well away from the smoke.

### NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.



### INGREDIENTS

<b>ACTIVE INGREDIENTS:</b>	
Carbaryl (1-naphthyl N-methylcarbamate)	19.6%
1, 1-Bis(chlorophenyl)-2,2,2-trichloroethanol	3.6%
Captan [N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide]	19.6%
<b>INERT INGREDIENTS</b>	
	57.2%
Total	100.0%

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

See Remainder of CAUTION Statement on Side Panel

USDA Reg. No. 1386-74

**NET WEIGHT**

(H&G-117)

**1 POUND**

Prod. 2i41

DISTRIBUTED BY

**UNITED CO-OPERATIVES, INC.**

**ALLIANCE, OHIO 44601**

### APPLES AND PEARS

Apply (1) when buds show pink, (2) just before blossom buds open, (3) when most of the petals have fallen, (4) ten days after petal fall, and (5) at seven to ten day intervals for two or more applications as needed, according to variety. **Do not apply within 14 days of harvest.**

Apply (1) when buds show pink, (2) just before blossom buds open, (3) when most of the petals have fallen, (4) ten days after petal fall, and (5) at seven to ten day intervals for two or more applications as needed, according to variety. **Do not apply within 7 days of harvest.**

May cause fruit thinning if applied during and within 30 days after full bloom. To avoid this, delay use until 30 days after bloom. May cause foliage injury on York and McIntosh apples if used before second cover spray. If required, use another suitable insecticide fungicide during these periods, in accordance with directions on its label. Foliage injury may result from combination with summer oils. Do not use on D'Anjou pears.

**PEACHES**—Oriental fruit moth, curculio, red banded leaf roller, tarnished plant bug, Japanese beetle adults, European red mite, two-spotted spider mite, brown rot, scab, and blossom blight (for which applications at 3 to 4 day intervals during bloom may be necessary).

Apply (1) when buds show pink, (2) just before blossom buds open, (3) when most of the petals have fallen, (4) ten days after petal fall, and (5) at seven to ten day intervals for two or more applications as needed, according to variety. **Do not apply within 14 days of harvest.**

**CHERRIES**—Black cherry aphid, leaf roller, slugs, European red mite, plum rust mite, two spotted mite, brown rot, and leaf spot.

Apply (1) at petal fall, (2) thirty days after petal fall, and (3) to trees after harvest at 30 day intervals for leaf spot. **DO NOT REPEAT APPLICATIONS ON CHERRIES WITHIN 30 DAYS.** Use another suitable insecticide fungicide according to directions on its label, particularly when plum curculio, cherry fruit fly, and blossom blight are present, in applications required at full bloom, shuck fall, and in the first cover spray subsequent to petal fall when this material cannot be used. **Do not apply within 7 days of harvest.**