



GARDEN SPRAY Powder



INGREDIENTS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	12.50%
Rotenone	2.50%
Other cube resins	4.00%
Zineb (zinc ethylene bisdithiocarbamate)	6.25%
Ziram (zinc dimethyl dithiocarbamate)	6.25%
INERT INGREDIENTS	68.50%

Total 100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1386-75

NET WEIGHT

H&G 109

1 POUND

Prod. 2142

DISTRIBUTED BY

UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

UNICO GARDEN SPRAY POWDER

Each 1 Pound Makes 8 1/2 Gallons of Spray

DIRECTIONS FOR USE

Mix 1 1/2 level tablespoons in 1 gallon of water. Mix 1 1/2 pounds in 100 gallons of water applied at the rate of 100 gallons per acre, thoroughly and evenly as a spray on the crops listed below to control the insects and plant diseases indicated. Begin applications before diseases are apparent and before insects become numerous which is usually when seedlings appear above ground. Young plants are transplanted into the garden. Spray foliage and fruit evenly and thoroughly, both upper and lower surfaces. Repeat applications at 7 to 10 day intervals and after heavy rains to maintain protection through growing season.

CROPS, INSECTS AND PLANT DISEASES

BEANS - Aphids, anthracnose, white mold, Mexican bean beetle, bean leaf beetle, trifoliate worm, cutworm, alfalfa weevil. WITHIN 7 DAYS OF HARVEST.

BROCCOLI, CABBAGE, CAULIFLOWER AND KALE - Imported cabbage worm, diamond-back moth, cabbage looper, lequin bug, and downy mildew. DO NOT APPLY WITHIN 14 DAYS OF HARVEST.

CARROTS - Flea beetles, carrot root fly, carrot root borer. DO NOT APPLY WITHIN 14 DAYS OF HARVEST. CARROT TOPS ARE TO BE USED AS FEED FOR LIVESTOCK. DO NOT APPLY WITHIN 14 DAYS OF HARVEST.

CUCUMBER, MELONS AND SQUASH - Aphids, hoppers, melon worm, pickle worm, squash vine borer, downy mildew, angular leaf spot, and leaf curl. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

EGGPLANT - Aphids, flea beetles, tobacco hornworm, Alternaria leaf spot, and early blight. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

GARDEN BEETS - Flea beetles, cutworm, and root borer. 14 DAYS OF HARVEST. LEAF TOPS ARE TO BE USED AS FOOD FOR PETS, OTHERWISE TO BE DESTROYED. WITHIN 3 DAYS OF HARVEST.

PEAS - Leafhopper, pea weevil, and pea root borer. WITHIN 10 DAYS OF HARVEST.

PEPPERS - Flea beetles, tarnished plant bug, and downy mildew. Apply 7 to 10 days before harvest. WITHIN 1 DAY OF HARVEST.

SPECIAL NOTES

Not effective if used with alkaline materials. Avoid spray when severe wind conditions are prevalent to minimize drift and possible damage.

BEEES - This product is highly toxic to bees exposed to direct treatment. Avoid spraying bees. Protective information by the Beekeepers National Cooperative Association Extension Service.

CAUTION

WARNING: KEEP OUT OF REACH OF CHILDREN. This product is highly toxic. Avoid contact with eyes. If contact occurs, wash thoroughly with plenty of water. If on skin, wash with soap and water. If inhaled, get fresh air. If swallowed, do not induce vomiting.

Do not eat, drink, or smoke. Keep out of body of water. Do not use in areas where children play. Do not use in areas where weather conditions are such that spray may be carried to other areas. Do not apply to plants that are stressed. Do not apply to plants that are in contact with water. Wash hands and clothing after use. Do not use in areas where children play.

NOTE: This product is highly toxic to bees. Do not use in areas where bees are present.

DO NOT REUSE CONTAINERS. Do not reuse containers for any purpose. Empty containers should be disposed of properly. Do not reuse containers for any purpose.

NOTICE

This product is highly toxic. Do not use in areas where children play. Do not use in areas where weather conditions are such that spray may be carried to other areas.