

300751-10
 JUN 14 1967
 UNDER THE FEDERAL INSECTICIDE ACT
 REGISTERED
 1386-79



**Water Soluble
 RAT AND MOUSE
 KILLER**

INGREDIENTS

ACTIVE INGREDIENTS:
 Sodium salt of 2-pivalyl-1,3-indandione* 0.14%
 INERT INGREDIENTS _____ 99.86%
 Total _____ 100.00%

*Equivalent to 0.13% 2-pivalyl-1,3-indandione

CAUTION

Keep Out of Reach of Children

**See Remainder of CAUTION
 Statement on Back Panel**

USDA Reg. No. 1386-79

NET WEIGHT

**1.5 Ounces Advp.
 Prod. 907**

Distributed By

UNITED CO-OPERATIVES, INC.
 Alliance, Ohio

DIRECTIONS FOR USE

Empty contents of this package into 1 quart of plain water to make liquid bait.

To control rats, place 1 pint to 1 quart of liquid bait in shallow containers or chicken founts and expose in locations where rats are known to frequent. At least two or more placements of liquid bait give best results.

To control mice, expose liquid bait in same manner as for rats, except smaller quantities in placements distributed at 5 to 10-foot intervals should be used in areas where mice or their signs have been seen.

Containers in which liquid bait is exposed must be made inaccessible to or protected from children, dogs, cats, livestock and poultry, if necessary by means of boxes or cages through which rodents can enter but which will exclude larger animals. Dogs and cats may actually search for and consume the liquid bait. It must be placed so as to positively prevent access by these pets.

Effective use of liquid bait requires CONTINUOUS exposure over a period of 12 to 14 days. Replenish bait solution before completely consumed, and replace if water becomes dirty. Baiting should be continued until all rodent activity ceases. Where re-infection is likely, permanent liquid bait stations should be used, and liquid bait placed in them as often as needed.

CAUTION

If taken internally, this rodenticide reduces the clotting ability of the blood and may cause serious hemorrhage. Humans, livestock, poultry, and pets must be protected against accidentally swallowing this powder or its solutions. Store powder and place liquid bait so as to be inaccessible to all animals except rodents. If taken internally, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. **CALL A PHYSICIAN IMMEDIATELY.**

NOTE FOR PHYSICIANS: Administration of blood transfusions combined with intravenous injections and oral doses of Vitamin K. is indicated.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on the face hereof.