

# **CAUTION**

## **KEEP OUT OF REACH OF CHILDREN**

EPA Reg. No. 1386-92

### **DIRECTIONS FOR USE**

To use as a spray, fill tank about one-quarter full with water, add the correct amount of powder, agitate or stir thoroughly to obtain a good suspension. Then, add remainder of water to fill spray tank and continue agitation. Some agitation is required to maintain the suspension while spraying. Proper timing of applications and thorough coverage are essential for satisfactory results.

#### **PREPLANTING APPLICATIONS TO SOIL**

**POTATOES, CORN, SWEET CORN, AND STRAWBERRIES** — To control wireworms (does not include Southern potato wireworm), and white grubs, use 10 to 20 pounds per acre in the quantity of water required for uniform spray coverage of the soil (7 ounces of powder per 1,000 sq. ft.). Apply spray to prepared soil surface just before seeding or transplanting, and immediately disc or rake into top 3 inches of soil. Properly applied soil treatments may be effective for more than one year. Repeat when new infestations develop.

**PEAS, BEANS, AND SWEET CORN** — To control seed corn maggot, use 3 to 4 pounds per acre in the quantity of water required for uniform spray coverage of the soil (1-1/2 ounces per gallon per 1,000 sq. ft.). Apply spray to soil surface of prepared seed bed and immediately disc or rake into top 3 inches of soil. Repeat annually.

#### **APPLICATIONS TO LAWNS AND ORNAMENTAL TURF**

To control chiggers, ants, crickets and ticks, use 3 to 4 pounds per acre in the quantity of water required for uniform spray coverage of the grass (1-1/2 ounces per gallon to 1,000 sq. ft.). Apply spray thoroughly to the grass and repeat as necessary. For imported fire ants in lawns and ornamental turf, break open hard surface of mound and pour in 8 ounces of dust per mound; spray surrounding soil as for other ants. If ants survive, repeat in 2 weeks. Do not contaminate food or feed crops.

**EARTHWORMS** — Apply at the rate of 20 to 25 pounds per acre. Do not contaminate food or feed crops.

**SODWEBWORMS** — Apply at rate of 6 to 7 pounds per acre. Do not contaminate food or feed crops.

To control white grubs and Japanese beetle grubs, use 10 to 20 pounds per acre in the quantity of water required for uniform spray coverage of the grass (7 ounces of powder per 1,000 sq. ft.). Apply spray to grass in early spring or late fall. Properly applied treatments at this rate may be effective for more than 1 year. Repeat when new infestations develop.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by adequate rainfall or thoroughly sprinkling, and the grass has completely dried. Do not feed or graze grass from treated areas to livestock and poultry. Do not use grass from treated areas for hay. Birds feeding over treated lawn or ornamental turf may be harmed.

#### **CAUTION**

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist or dust. In case of contact with skin wash with soap and water. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish and wildlife. Keep out of any body of water. Do not apply to crops or areas other than those for which directions are stated in this label. Do not apply when weather conditions favor drift of spray from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land located well away from water supplies.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke is poisonous. Stay well away from the smoke.

#### **NOTICE**

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

## **NET WEIGHT**

## **50 POUNDS**

FC-251-A872

PROD. 1067

**DISTRIBUTED BY**  
**UNITED CO-OPERATIVES, INC.**  
ALLIANCE, OHIO 44601