



5% CHLORDANE DUST

10-11-72
THE FEDERAL INSECTICIDE
ACT AND
1386-223

UNICO 5% Chlordane Dust

CAUTION

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing dust or vapor. In case of contact with skin, wash with soap and water. Do not contaminate feed and foodstuffs. This product is toxic to fish, birds and wildlife. Keep out of any body of water. Do not apply in any manner not specified in this label. Do not apply when weather conditions favor drift of dust from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land located well away from water supplies.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS Do not reuse this container for any purpose. Burn empty containers in a hot fire, outdoors, in an isolated area. The smoke is poisonous. Stay well away from the smoke.

DIRECTIONS FOR USE

LAWNS AND ORNAMENTAL TURF

ANTS, CHIGGERS, SOD WEBWORMS, CRICKETS, CUTWORMS, AND EARWIGS—Apply 25 to 30 lb. per acre (6 oz. per 500 sq. ft. of area) directly into the grass. Repeat as necessary.

For imported fire ant in lawns, break open hard surface of mound and pour in 2 to 4 cupfuls per mound; sprinkle surrounding soil as for other ants. If ants survive, repeat in 2 weeks.

WHITE GRUBS AND JAPANESE BEETLE GRUBS—Apply 150 lb. per acre (2 lb. per 500 sq. ft. of area) in early spring or late fall. Repeat annually if necessary.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by adequate rainfall or thorough sprinkling, and the grass has completely dried. **DO NOT** graze or feed forage from treated areas to livestock and poultry.

Birds feeding over treated areas may be harmed.

PREPLANTING SOIL APPLICATIONS

CUTWORMS—Apply 25 to 30 lb. per acre (6 oz. per 500 sq. ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper, or tobacco transplants. If desired, place 2 to 3 teaspoonfuls in a ring around each transplant but not in direct contact with plant.

WIREWORMS—Apply 75 to 100 lb. per acre (1 to 1 1/4 lb. per 500 sq. ft.) uniformly over surface of prepared land, work into top 3 to 4 inches of soil prior to planting of corn, sweet corn, tobacco, or strawberries.

CABBAGE ROOT MAGGOT—Apply 75 to 100 lb. per acre (1 to 1 1/4 lb. per 500 sq. ft.). Work into top 3 inches of soil just before setting of cabbage, broccoli, or cauliflower transplants.

SEED CORN MAGGOT—Apply 25 to 30 lb. per acre (6 oz. per 500 sq. ft.). Work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS—Uniformly apply 150 to 200 lb. per acre (2 lb. per 500 sq. ft.) of prepared soil and work into top 4 to 6 inches of soil just before setting of strawberry or ornamental transplants.

Repeat preplanting soil treatments annually, if necessary.

INGREDIENTS

ACTIVE INGREDIENT		
Technical Chlordane*	5.0%
INERT INGREDIENTS	95.0%
Total	100.0%

*Equivalent to 3.0% octachloro-4,7-methanotetrahydroindane and 2.0% related compounds.
EPA Reg. No. 1386-223

CAUTION

Keep Out Of Reach Of Children

See Remainder of CAUTION Statement on Back Panel

NET WEIGHT

5 POUNDS

FC-545-A672

PROD. 1041

DISTRIBUTED BY
UNITED CO-OPERATIVES, INC.
ALLIANCE, OHIO 44601

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

FC-545-A672

PROD. 1041