

DO NOT USE ON LAWNS OR GRASS INTERPLANTED WITH CLOVER UNLESS PROBABLE INJURY TO THE CLOVER CAN BE TOLERATED. DO NOT APPLY TO GRASS SEED CROPS DURING THE FLOWERING OR POLLINATING STAGE. DO NOT APPLY TO GRASS IN THE SEEDLING STAGE; WAIT UNTIL GRASS IS ROOTED AND WELL TILLERED.

**CAUTION**

Harmful if swallowed. Avoid contact with skin, eyes or clothing, avoid breathing spray mist. If contact with eyes occurs, flush with plenty of water. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate any body of water, especially domestic and irrigation water supplies.

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Use only lawn sprayers for this product. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. Bury unused spray and rinse solutions well away from desirable plants and drip zones of shade trees.

**DESTROY EMPTY CONTAINERS:** Do not reuse this container for any purpose. Break empty container into a previously prepared pit on non-crop land in an isolated area, and bury fragments deeply.

**KEEP FROM FREEZING NOTICE**

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

**SPECIMEN LABEL**

**UNICO**

**LAWN WEED KILLER**

Kills Many Common Broadleaf Weeds in Lawns. Contains 2,4-D Amine.

**INGREDIENTS**

ACTIVE INGREDIENT:	
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid*	13.2%
INERT INGREDIENT	86.8%
Total	100.0%

\*EQUIVALENT TO 11.0% 2,4-dichlorophenoxyacetic acid. Isomer specific by AOAC Method No. 6 D01-5

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1338-109  
Net Volume

EPA Est. No. 1388-OH-1  
1 U.S. Quart Liquid

021015-8 80

MANUFACTURED BY PROD. 2101



**universal cooperatives, inc.**  
Minneapolis, Minnesota

**UNICO  
DIR**

Mix with water for application to the water in the tank or when weeds are young and best results. Kill of weeds spraying. Retreatment of necessary.

**LAWNS**—To control ragweed, lambsquarter, cock fleabane, pigweed, wild rag, stinging nettle, and sunflower water and apply this mixture (20x50).

To control certain jimsonweed, prickly lettuce, henbit and buttercup, use 1 and apply this mixture to wet (20x50).

**VACANT LOTS AND NO** weeds such as those listed in lots, drainage ditch-banks, useful or ornamental plants (1/4 cupful) in 1 gallon of water thoroughly.

Repeated application to maintain control. 2,4-D is not plants and vines to control the honeysuckle may be controlled.

**WILD GARLIC AND WIL** in lawns, use 6 tablespoonful to the actively growing plants in successive years until no will be killed by this treatment.

**UNICO**®

**LAWN WEED KILLER**

UNIVERSAL COOPERATIVES, INC.  
1226-101  
EPA Reg. No. 1386-109

Kills Many Common Broadleaf Weeds in Lawns. Contains 2,4-D Amine.

INGREDIENTS	
ACTIVE INGREDIENT:	
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid*	13.2%
INERT INGREDIENT	86.8%
Total	100.0%

\*EQUIVALENT TO 11.0% 2,4-dichlorophenoxyacetic acid. Isomer specific by AOAC Method No. 6 D01-5

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Remainder of CAUTION Statement on Side Panel  
EPA Reg. No. 1386-109      EPA Est. No. 138C-CH-1  
Net Volume      1 U.S. Quart Liquid  
921015-8-80      MANUFACTURED BY PROD. 2101



**universal cooperatives, inc.**  
Minneapolis, Minnesota 55420

WNS OR GRASS INTERPLANTED WITH DESIRABLE INJURY TO THE CLOVER CAN BE APPLIED TO GRASS SEED CROPS DURING POLLINATING STAGE. DO NOT APPLY TO GRASS STAGE; WAIT UNTIL GRASS IS ROOTED

**CAUTION**

Avoid contact with skin, eyes or clothing. If contact with eyes occurs, flush with water. Product is toxic to fish. Keep out of lakes, streams, rivers, ponds, and other bodies of water. Do not contaminate any body of water, including irrigation water supplies.

Do not apply to vegetables, flowers, ornamental plants, or desirable plants. Do not store in contact with insecticides, or fungicides. Use only lawn mowers. Coarse sprays are less likely to drift. Do not apply to shade trees. Do not apply to shade trees. Do not apply to shade trees.

REUSING CONTAINERS: Do not reuse this container for any other purpose. Empty container into a previously prepared pit in an isolated area, and bury fragments deeply.

**KEEP FROM FREEZING NOTICE**

Use and in compliance with the limitations, directions, and cautions stated on this label.

**KEEP FROM FREEZING NOTICE**

**UNICO Lawn Weed Killer DIRECTIONS FOR USE**

Mix with water for application as a spray. To mix, add this weed killer to the water in the tank or container and stir or agitate thoroughly. Apply when weeds are young and in a succulent, rapidly growing condition for best results. Kill of weeds may not be evident for about 2 weeks after spraying. Retreatment of areas infested with perennial weeds may be necessary.

**LAWNS**—To control susceptible broadleaf weeds such as mustard, ragweed, lambsquarter, cocklebur, plainfain, morning glory (annual), daisy fleabane, pigweed, wild radish, buckhorn, bull thistle, burdock, dandelion, stinging nettle, and sunflower, use 6 to 7 tablespoons (3/8 cupful) in 1 gallon of water and apply this mixture to wet uniformly the weed plants on 1,000 square feet (20'x50').

To control certain less susceptible broadleaf weeds such as jimsonweed, prickly lettuce, mallow, purslane, smartweed, wild carrot, henbit and buttercup, use 12 tablespoonfuls (1/2 cupful) in 1 gallon of water and apply this mixture to wet uniformly the weed plants on 1,000 square feet (20'x50').

**VACANT LOTS AND NON-CROP AREAS**—To control certain broadleaf weeds such as those listed above as occurring in lawns, on unused vacant lots, drainage ditch-banks, roadsides, and other uncared for areas, where useful or ornamental plants are not growing, use 8 to 12 tablespoonfuls (1/3 cupful) in 1 gallon of water and apply as a coarse spray to wet weeds thoroughly.

Repeated application may be necessary if new weed growth occurs to maintain control. 2,4-D is not sufficiently active against many kinds of woody plants and vines to control them satisfactorily. Poison oak, poison ivy, and honeysuckle may be controlled if applications are repeated several times.

**WILD GARLIC AND WILD ONION**—To control wild garlic and wild onion in lawns, use 6 tablespoonfuls (3/8 cupful) in 1 gallon of water and apply spray to the actively growing plants in the early spring and again in late fall. Repeat in successive years until no new growth appears. Legumes (clover) probably will be killed by this treatment.

84A