

1386-143



## UNICO Livestock and Barn Fogging Spray

### DIRECTIONS FOR USE

UNICO Livestock and Barn Fogging Spray, a ready-to-use spray for barn foggers and sprayers which dispense an atomized mist, is recommended for control of house flies, stable flies, and horn flies which enter the barn on livestock including dairy cattle, and for house flies which congregate in the barn between milkings when applied over the animals in the barn as a fog or finely atomized mist, and for protection of dairy cattle, beef cattle, horses, and hogs, against horn flies, house flies, stable flies, and mosquitoes when applied directly to the animals as a fine spray.

**BARN FOGGING**—To control house flies, stable flies, and horn flies on the animals and in the barn, apply as a fog or finely atomized mist over the livestock in stanchions or in holding pens. Use one fogger or spray unit for each 2,500 sq. ft. of barn area, placed above the animals and preferably near the center of the area. Doors and windows should be closed prior to fogging; openings may cause drafts and permit escape of the fog. Operate the unit until spray mist or fog is thoroughly distributed throughout the entire area; spray out ½ fl. oz. uniformly per 1,000 cu. ft. Do not ventilate the barn for 10 to 15 minutes after the unit has been stopped. Permit the barn to aerate before milking. **REPEAT DAILY, MORNING AND AFTERNOON, OR AS NEEDED.** Keep sprayer, container and nozzle clean to ensure proper delivery of fog or spray mist.

The sprayer or fogger should be one made to spray or fog oil or water based insecticide sprays. However, the important point to keep in mind is that a definite amount of toxicant, uniformly distributed per unit of space, is necessary in order to perform an efficient job.

# LIVESTOCK AND BARN FOGGING SPRAY

Contains Pyrenone®

### INGREDIENTS

|                                     |       |
|-------------------------------------|-------|
| ACTIVE INGREDIENTS .....            | 100%  |
| Technical piperonyl butoxide* ..... | 1.0%  |
| Pyrethrins .....                    | .1%   |
| Petroleum base oil .....            | 98.9% |

\*Equivalent to 0.8% (Eutylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

Technical piperonyl butoxide and pyrethrins are constituents of Pyrenone Reg. U.S. Pat. Off., F.M.C.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel.

EPA Reg. No. 1386-143

Net Volume

FC-706A-7-72

5 U.S. Gallons Liquid

PROD. 310

MANUFACTURED BY

ACCEPTED  
Aug-14-19  
1386-14

# LIVESTOCK AND BARN FOGGING SPRAY

Contains Pyrenone®

## INGREDIENTS

|                                     |       |
|-------------------------------------|-------|
| ACTIVE INGREDIENTS .....            | 100%  |
| Technical piperonyl butoxide* ..... | 1.0%  |
| Pyrethrins .....                    | .1%   |
| Petroleum base oil .....            | 98.9% |

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

Technical piperonyl butoxide and pyrethrins are constituents of Pyrenone Reg. U.S. Pat. Off., F.M.C.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

See Remainder of CAUTION Statement on Side Panel.

EPA Reg. No. 1386-143

Net Volume

5 U.S. Gallons Liquid

FC-706A-7-72

PROD. 310

MANUFACTURED BY

UNITED CO-OPERATIVES, INC., ALLIANCE, OHIO 44601

**DIRECT SPRAYING OF LIVESTOCK**—To protect dairy cattle, beef cattle, horses and hogs against horn flies, house flies, stable flies, and mosquitoes, apply  $\frac{1}{8}$  to  $\frac{1}{2}$  fl. oz. (approximately 2 to 3 teaspoonfuls) as a fine spray uniformly distributed over each animal. Use a good hand sprayer or electric sprayer to spray each animal, preferably from the rear, directing the spray along the sides, including the belly, flanks, and withers, down each leg, over the back, and around the poll. Hold nozzle of sprayer 2 to 3 feet away from animal so as to deposit droplets on the hair. **DO NOT WET THE HIDE.** For milking dairy animals, apply the spray 20 to 30 minutes before beginning to milk. **DO NOT SPRAY UDDERS OF LACTATING DAIRY ANIMALS.** Repeat daily, morning and afternoon, as needed.

To control lice on cattle and horses, apply 5 teaspoonfuls (approximately  $\frac{5}{6}$  fl oz.) as a fine spray to each animal, with particular attention to infested areas. Rub spray by hand into the coat. Repeat at intervals of 2 to 3 weeks as needed.

## CAUTION

Harmful if swallowed. Avoid contamination of feed, foodstuffs, and milking utensils or equipment.

Do not wet the skin of animals as this may cause blistering. Do not apply to plants or vegetation. Do not store or use near heat or open flame.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. If container is glass, break into previously prepared pit on non-crop land well away from water supplies and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe and bury crushed container in a pit on non-crop land well away from water supplies.

## NOTICE

Use only for the purpose and in compliance with the limitations, cautions or warnings stated on this label.

