



MALATHION-4

CATTLE and POULTRY DUST

INGREDIENTS

ACTIVE INGREDIENTS:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)

INERT INGREDIENTS

Total

USDA Reg. No. 1386-144

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, wash thoroughly after use. Avoid contamination of feed and foodstuffs.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Burn empty containers in a hot fire, outdoors, in an isolated area. The smoke is hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

CATTLE To control hornflies on cattle, including dairy cattle, apply 2 ounces (4 tablespoonfuls) per animal. Sprinkle the powder on poll, neck (top and sides), back, and upper portion of each side; rub in lightly by hand to distribute powder beneath the hair. Repeat applications at intervals of 10 to 14 days as necessary to maintain control of horn flies. Treat dairy animals in stanchions within an hour after milking to avoid contamination of the milk. **DO NOT APPLY TO LACTATING DAIRY CATTLE WITHIN 5 HOURS BEFORE MILKING OR DURING MILKING. AVOID CONTAMINATION OF FEED, FEEDING TROUGHS, WATERING CUPS OR DEVICES, AND MILKING EQUIPMENT.**

POULTRY (Direct Applications. To control Northern fowl mite (feather mite) and poultry lice, and as a supplement to the treatment of poultry houses for chicken red mite, apply dust directly to birds by means of rotary or plunger hand duster, or by sprinkling from a sifter-top can or other container. Care must be taken to assure thorough coverage of each bird and uniform distribution of the dust through the feathers down to the skin. On caged birds, apply the dust to the birds and cages, as well as to the adjacent structure. To treat individual birds, hold by legs with head downward and work the dust through feathers to the skin, beginning at the vent and working down each side to the head. Repeat in 4 to 8 weeks or when necessary.

POULTRY HOUSES To control chicken red mite, Northern fowl mite (feather mites), and poultry lice apply 1 pound of powder per 50 to 60 square feet of litter and floor surface and to nests, roosts, and adjacent areas. Dust may be applied with a rotary or plunger hand duster, or by sprinkling from a sifter-top can or other container, provided care is taken to assure thorough coverage and uniform distribution of the dust over the surfaces and areas specified. Repeat treatment as necessary.

AVOID CONTAMINATION OF FEED, FEEDING TROUGHS, AND WATER FOUNTAINS, WATER CUPS OR WATERING TROUGHS WITH THE DUST.

NOTICE

Use only for the purposes and in compliance with the limitations, cautions or warnings stated on this label.

NET WEIGHT

5 POUNDS

FC 513 C1268

PROD 972

DISTRIBUTED BY:
UNITED CO-OPERATIVES, INC.
ALLIANCE OHIO

ACCEPTED
May - 26 1971
1386-144